

# Metal-free selective mono-halodecarboxylation of heteroarenes under mild conditions

Scott H. Henderson,<sup>a</sup> Ryan A. West,<sup>a</sup> Simon E. Ward<sup>b</sup> and Mark A. Honey<sup>a,c,†\*</sup>

<sup>a</sup>Sussex Drug Discovery Centre, University of Sussex, Brighton, BN1 9RH, UK

<sup>b</sup>Medicines Discovery Institute, Park Place, Cardiff University, CF10 3AT, UK

<sup>c</sup>School of Pharmacy, Faculty of Science & Engineering, University of Wolverhampton, Wolverhampton, WV1 1LY, UK

**Keywords:** Halodecarboxylation; Selective Halogenation; Mild Conditions

## 1. Summary

The halodecarboxylation of heteroarene carboxylic acids by treatment with N-bromosuccinimide (NBS) or N-chlorosuccinimide (NCS) was performed. This procedure provides a convenient route to synthetically useful mono-halogenated heteroarene intermediates such as halo-indoles, -aza-indoles, -indazoles and -aza-indazoles. The mild conditions employed and simple protocol provides an advantage over traditional halodecarboxylation procedures that require expensive and toxic metal catalysts, basic conditions, time-consuming intermediate isolation and elevated reaction temperatures.

## 2. Introduction

Haloheteroarenes serve as versatile building blocks in organic chemistry, most notably as coupling partners in transition metal-catalysed cross-coupling reactions for the formation of C-C and C-N bonds. Due to their utility, haloheteroarenes have been used in the construction of molecules of biological interest including natural products<sup>1,2,3</sup> and small molecules in drug discovery programs such as Merck's recent LRRK2 inhibitor, MLI-2,<sup>4</sup> and Vertex's PKC  $\theta$  inhibitor (Fig 1).<sup>5</sup>

Figure 1

Few methods are documented that describe the selective monohalogenation of heteroaromatic scaffolds.<sup>6,7,8</sup> A traditional approach to the regiospecific installation of a halogen atom on a substrate is the thermal halodecarboxylation of a silver(I) carboxylate salt in the presence of a halide source known as the Hunsdiecker-Borodin reaction (HBR) (Fig 2).<sup>9,10</sup> Substrates for the HBR are often aliphatic carboxylates with limited aromatic and heteroaromatic carboxylates being employed successfully.<sup>11</sup> The preparation of bromopyridines under basic aqueous conditions at elevated temperatures offers a rare example of the HBR being successfully applied to heterocyclic acid substrates.<sup>12,13</sup> The major shortcoming of the HBR is the requirement to isolate the dry silver carboxylate salt in sufficient purity to undergo decarboxylation and afford the corresponding halide in good yield.<sup>14</sup> Accordingly, modifications of the HBR such as the Cristol-Firth modification (CFM) and Kochi modification (KM) avoid the need to isolate the silver salt by employing more stable metal salts such as red mercury<sup>15</sup> and lead acetate<sup>16</sup> that undergo halodecarboxylation on heating *in situ*. The CFM and KM suffer from the same major limitation that they both employ toxic metal reagents to initiate the decarboxylation. The Barton modification (BM) avoids the use of toxic and expensive metals yet requires the preparation of the respective thiohydroxamate ester (Barton ester). Halodecarboxylation of the Barton ester results in the formation of both the desired halide and the undesired 2-(alkylthio)pyridine side-product, limiting the atom efficiency of the reaction (Fig 2).<sup>17</sup> Recent advances in the field include the metal-free decarboxylative halogenation of unsaturated carboxylic acids with *N*-halosuccinimides in the presence of TBA-TFA,<sup>19</sup> LiOAc<sup>18</sup> or photolysis with bromine and  $\text{PhI}(\text{OAc})_2$ .<sup>20</sup> Hypervalent iodine has also been utilized by Hamamoto and Miki<sup>21</sup> for

\*Author for correspondence ([m.honey@sussex.ac.uk](mailto:m.honey@sussex.ac.uk) / [m.honey@wlv.ac.uk](mailto:m.honey@wlv.ac.uk)).

†Present address: School of Pharmacy, Faculty of Science & Engineering, University of Wolverhampton, Wolverhampton, WV1 1LY, UK

the synthesis of polybromoindoles. However, this methodology is seemingly unable to produce mono-halogenated products and requires the use of alkali metal salts as the halide source. Since preparing this manuscript, a recent report by Larrosa<sup>22</sup> has described the decarboxylative iodination of benzoic and heterobenzoic acids through the use of iodine and potassium phosphate. Although this protocol is very diverse, it requires the use of stoichiometric amounts of base, elevated temperatures (100 °C), and currently suffers from over bromination (when bromine is used in place of iodine) of substrates due to the increased electrophilicity of bromine in comparison to that of iodine.

Figure 2

With the above in mind, current methods of halodecarboxylation suffer from the use of highly toxic and expensive metals employed in stoichiometric amounts, requires the use of corrosive elemental halide sources or suffer over halogenation. It is therefore desirable to establish a more reliable and economical method for mono-halodecarboxylation of heteroarene carboxylic acids to furnish synthetically useful haloheteroarenes. Presently, we describe a simple base-free method for the regiospecific mono-halodecarboxylation of substituted and unsubstituted heteroarene carboxylic acids under mild and metal-free conditions at room temperature with NBS and NCS.

### 3. Results and Discussion

Our interest in the synthesis of haloheteroarene precursors peaked when exploring indazoles as a potential scaffold for an in-house kinase inhibitor project. Our studies began somewhat fortuitously with the bromodecarboxylation of indazole **1** with NBS. The reaction proceeded with two equivalents of NBS in *N,N*-dimethylformamide (DMF) at room temperature affording product **2** in 35% yield (Table 1, entry 1) after just 1 h. Inspection of the reaction mixture by LC-MS analysis appeared to show that a dibrominated product had also formed as a major impurity. Gratifyingly, by decreasing the equivalents of NBS it was observed that no measurable amount of dibrominated product was formed. Despite the reaction being facile it was found that by extending the reaction time from 1 h to 16 h the reaction reliably went to completion. Under the optimised conditions product **2** was furnished in 77% isolated yield (Table 1, entry 2).

Reactions involving NBS can be exothermic and can pose a risk when undertaken on a large scale,<sup>23</sup> therefore we sought to employ a solvent that was predicted to be safe to use on a large scale and in addition would furnish the desired product in high yield. A small solvent screen was undertaken with solvents that were reported to cause negligible exotherms when used with NBS (Table 1, entries 3-5).<sup>23</sup> Unfortunately none of these solvents facilitated the halodecarboxylation reaction in as high yield as observed when the reaction was carried out in DMF (Table 1).

Table 1

With the halodecarboxylation procedure in hand, the indazole carboxylic acid scope was investigated (Table 2). A variety of indazole carboxylic acids were subjected to the reaction with NBS in DMF (products **2-13**). Examples of indazole acids bearing electron withdrawing (products **3-5**) and electron donating substituents (products **8** and **9**) underwent decarboxylative bromination in excellent yields under the reaction conditions. However, when the electron donating methoxy group was moved to the 4, 6 and 7 positions on the indazole (**10**, **11** and **12**), no product could be detected. *N*-methylation of the indazole to afford compound **13** appeared to be unsuccessful, with no halodecarboxylated product observed by LC-MS - it appears as though the presence of the free indazole N-H is a requirement for decarboxylation when using the current set of reaction conditions. Halodecarboxylation to afford the 5-amino indazole analogue (product **7**) also proved unsuccessful. Initial mechanistic investigations reveal that the reaction possibly proceeds through a non-radical pathway, as the reaction continued to proceed when performed in the absence of light. However, detailed mechanistic investigations are continuing within our laboratory, and so a clearer insight into the lack of reactivity of some substrates will be detailed in due course.

Table 2

We now turned our attention to chlorodecarboxylation and iododecarboxylation reactions using the optimised conditions but employing *N*-chlorosuccinimide (NCS) and *N*-iodosuccinimide (NIS) as halide sources (Table 3, entries 2 and 3). Although the chlorodecarboxylation product (Table 3, entry 2) was obtained in lower yield than the bromodecarboxylation product (Table 3, entry 1) using similar conditions, the current procedure represents one of the few ways to synthesise chloroheteroarenes under mild conditions. Unfortunately, the iododecarboxylation was not facilitated under the current reaction conditions.

Table 3

To demonstrate the versatility of the halodecarboxylation procedure for heteroarenes, further reactions with NBS were investigated (Table 4). Although halo-indoles (products **14-16**) have been reported in the literature to be problematic to isolate,<sup>24,25</sup> under the current procedure we have been able to furnish **15** and **16** in reasonable yields.

Table 4

Aza-indole and aza-indazole acids underwent halodecarboxylation smoothly to furnish the corresponding haloheteroarenes **17** and **18** in good yield, implying that more electron deficient systems undergo halodecarboxylation efficiently under the current conditions. Attempts to further expand this methodology to simple benzoic acids have currently proved unsuccessful, although this remains an active area of research within our group.

## Conclusions

In summary, a mild and efficient protocol for the mono-halodecarboxylation of heteroarene carboxylic acids has been developed. It is noteworthy that this halodecarboxylation is metal-free and displays significant advantages over traditional methods of halodecarboxylation that commonly require harsh reaction conditions, toxic reagents or suffer over bromination. The proposed method extends the substrate scope of traditional halodecarboxylations to heteroarene substrates. This methodology provides a novel and direct route to haloheteroarene building blocks that can be used in transition-metal catalysed cross-coupling reactions to construct molecules of scientific importance. Investigations into the site selective halodecarboxylation of other heteroarene acid precursors is currently underway in our laboratory.

## Acknowledgements

We have no further acknowledgements to report

## Ethical Statement

We have no ethical interests to report

## Funding Statement

S. H. H. is funded by the BBSRC (BB/L017105/1); R. A. W is funded by the BBSRC (BB/L017180/1); S. E. W. is funded by the Wellcome Trust, MRC, BBSRC and the European Structural Funding via Sêr Cymru scheme; M. A. H. was funded by the University of Sussex.

## Data Accessibility

Experimental data associated with this article can be found as part of the Supplementary Material

## Competing Interests

We have no competing interests.

## Authors' Contributions


S. H. H. was involved in the conception and design, acquisition of data, analysis and interpretation of data and drafting and revising of the article. R. A. W. was involved in the conception and design, acquisition of data, analysis and interpretation of data and drafting and revising of the article. S. E. W. was involved in the conception of the project and revising the article. M. A. H. was involved in the conception and design, acquisition of data, analysis and interpretation of data and drafting and revising of the article. All Authors approved the final version to be published.

## References

- 1) Joules, J.A.; Mills, K.; *Heterocyclic Chemistry Fifth Edition*. John Wiley and Sons, Ltd. Chichester, UK. 2010. (ISBN: 978-1-405-13300-5)
- 2) Gribble, G. W. *J. Chem. Educ.* 2004, **81**, 1441 (DOI: 10.1021/ed081p1441)
- 3) Carbain, B., Hitchcock, P.B., Streicher, H. *Tetrahedron Lett.* 2010, **51**, 2717-2719 (DOI: <https://doi.org/10.1016/j.tetlet.2010.03.044>)
- 4) Scott, J.D; *et al. J. Med. Chem.* 2017, **60**, 2983-2992 (DOI: 10.1021/acs.jmedchem.7b00045)
- 5) Jimenez J-M; *et al. J. Med. Chem.* 2013, **56**, 1799-1810 (DOI: 10.1021/jm301465a)
- 6) Ganguly, N.C.; De, P.; Dutta, S. *Synthesis*. 2005, **7**, 1103-1108 (DOI: 10.1055/s-2005-861866)
- 7) Grimmitt, R. *Halogenation of Heterocycles: III. Heterocycles Fused to Other Aromatic or Heteroaromatic Rings, Advances in Heterocyclic Chemistry*, 1994, **59**, 245-369. (DOI: 10.1016/S0065-2725(08)60009-4)
- 8) Eisch, J.J. *Halogenation of Heterocyclic Compounds, Advances in Heterocyclic Chemistry*, 1967, **7**, 1. ([https://doi.org/10.1016/S0065-2725\(08\)60588-7](https://doi.org/10.1016/S0065-2725(08)60588-7))
- 9) a) Borodin, B. *Liebigs Ann.* 1861, 119, 1221. b) Hunsdiecker, H. C. *Chem. Ber.* 1942, **75**, 291. (DOI: 10.1002/cber.19420750309)
- 10) Kürt, L.; Czakó, B. *Strategic Applications of Named Reactions in Organic Synthesis: Background and Detailed Mechanisms*. Elsevier Academic Press.



- Burlington, MA, 2005. (eBook ISBN: 9780080919164)
- 11) R. G. Johnson and R. K. Ingham, *Chem. Rev.* 1956. **56**, 219-269 (DOI: 10.1021/cr50008a002)
  - 12) Kuffner, F.; Russo, C. *Monatsh.* 1954. **85**, 1097-1103. (DOI: <https://doi.org/10.1007/BF00899858>)
  - 13) Mariella, R. P.; Belcher, E. P. *J. Am. Chem. Soc.* 1952. **74**, 1916-1919. (DOI: 10.1021/ja01128a013)
  - 14) Laue, T.; Plagens, A. *Named Organic Reactions, Second Edition.* John Wiley and Sons, Ltd. Chichester, UK. 2005 (ISBN: 978-0-470-01041-9)
  - 15) Cristol, S.; Firth, W. *J. Org. Chem.* 1961. **26**, 280. (DOI: 10.1021/jo01060a628)
  - 16) Kochi, J.K. *J. Org. Chem.* 1965. **30**, 3265-3271. (DOI: 10.1021/jo01021a002)
  - 17) Barton, D.H.R.; Lacher, B.; Zard, S.Z. *Tetrahedron Lett.* 1985. **26**, 5939-5942. ([https://doi.org/10.1016/S0040-4039\(00\)98266-2](https://doi.org/10.1016/S0040-4039(00)98266-2))
  - 18) Roy, S. *J. Org. Chem.* 1997. **62**, 199-200. (DOI: 10.1021/jo951991f)
  - 19) Roy, S. *Tetrahedron.* 2000. **56**, 1369-1377. ([https://doi.org/10.1016/S0040-4020\(99\)01035-2](https://doi.org/10.1016/S0040-4020(99)01035-2))
  - 20) a) Concepcion, J. I.; Francisco, C. G.; Freire, R.; Hernandez, R.; Salazar, J. A.; Suarez, E. *J. Org. Chem.* 1986. **51**, 402-424. (DOI: 10.1021/jo00353a026) b) Moriarty, R.M. *J.C.S. Chem. Comm.* 1987, **0**, 675-676 (DOI: 10.1039/C39870000675)
  - 21) Hamamoto, H.; Umamoto, H.; Umamoto, M.; Ohta, C.; Fujita, E.; Nakamura, A.; Maegawa, T.; Miki, Y., *Heterocycles*, 2015, **91**, 561-572 (DOI: 10.3987/COM-14-13162)
  - 22) Perry, G. J. P.; Quibell, J. M.; Panigrahi, A.; Larrosa, I., *J. Am. Chem. Soc.* 2017, **139**, 11527-11536 (DOI: 10.1021/jacs.7b05155)
  - 23) Shimizu, S.; Imamura, Y.; Ueki, T. *Org. Process Res. Dev.* 2014. **18**, 354-358. (DOI: 10.1021/op400360k)
  - 24) Piers, K.; Meimaroglou, C.; Jardine, R.V.; Brown, R.K. *Canadian Journal of Chemistry.* 1963. **41**, 2399-2401 (doi.org/10.1139/v63-353)
  - 25) Weissgerber, R. *Ber.* 1913. **46**, 652. (DOI: 10.1002/cber.19130460188)

## Tables


**Table 1** Screening of optimal conditions


Entry	Solvent	NBS (equiv.)	Reaction time (h)	Isolated yield (%)	% conversion by LCMS
1	DMF	2	1	35	35
2	DMF	1	16	77	86
3	DCM	1	16	-	Trace product
4	EtOAc	1	16	-	50
5	MeCN	1	16	-	75

Reaction conditions: 1 (0.2 mmol), NBS, solvent (1.5 mL) at room temperature (rt). NBS = *N*-bromosuccinimide, DMF = *N,N*-dimethylformamide



**Table 2** Halodecarboxylation of indazoles

Reaction conditions: heteroarene carboxylic acid (0.2 mmol), NBS (0.2 mmol), solvent (1.5 mL) at room temperature. NBS = *N*-bromosuccinimide, DMF = *N,N*-dimethylformamide

**Table 3** Screening of halodecarboxylation conditions


Entry	NXS	Isolated yield (%)	% conversion by LCMS-
1	NBS	77	86
2	NCS	39	35
3	NIS	-	0

Reaction conditions: 1 (0.2 mmol), NXS, DMF (1.5 mL) at room temperature. NBS = *N*-bromosuccinimide, DMF = *N,N*-dimethylformamide

**Table 4** Halodecarboxylation of heteroarene carboxylic acids

Reaction conditions: heteroarene carboxylic acid (0.2 mmol), NBS (0.2 mmol), solvent (1.5 mL) at room temperature. NBS = *N*-bromosuccinimide, DMF = *N,N*-dimethylformamide


## Figures




**Figure 1.** Chemical bonds in red can be constructed through cross coupling with haloheteroarene precursor.

## Previous work:

### Hunsdiecker-Borodin




### Barton




R = alkyl, in general limited aryl and heteroaryl


### Larrosa



### Miki and Hamamoto



### This work:



R' = EWG, EDG, H

Figure 2. Strategies for halodecarboxylation<sup>8,11</sup>