

The Impact of Individual Health Education on Health Literacy: Evaluation of the Translated Version (Sinhala) of Health Education Impact Questionnaire in Type 2 Diabetes

Bulathsinghalage Poornima Reshamie Cooray, Hana Morrissey, Eisha Indumani Waidyarathnec, Patrick Anthony Ball

Bulathsinghalage Poornima Reshamie Cooray, School of Psychological and Clinical Sciences, Faculty of Engineering, Health, Science and the Environment, Charles Darwin University, Australia
Hana Morrissey, School of Pharmacy, Faculty of Science and Engineering, University of Wolverhampton, Wolverhampton, WV1 1LY, the United Kingdom

Eisha Indumani Waidyarathnec, Department of Anatomy, Faculty of Medicine, University of Ruhuna, Galle, Sri Lanka

Patrick Anthony Ball, School of Pharmacy, Faculty of Science and Engineering, University of Wolverhampton, Wolverhampton, WV1 1LY, the United Kingdom

Conflict-of-interest statement: The author(s) declare(s) that there is no conflict of interest regarding the publication of this paper.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Hana Morrissey, School of Pharmacy, Faculty of Science and Engineering, University of Wolverhampton, Wulfruna Street, Wolverhampton, WV1 1LY, the United Kingdom.

Email: patrick.ball@wlv.ac.uk

Telephone: +44 (0) 1902 322173

Received: February 15, 2018

Revised: May 6, 2018

Accepted: May 8, 2018

Published online: May x, 2018

ABSTRACT

INTRODUCTION: The prevalence of Type 2 diabetes is globally on the rise, in both developed and developing countries. Type 2

diabetes is a major public health issue in Sri Lanka. This study aims to investigate the effect of structured self-management health education intervention based on ‘PITS model’ (Pathophysiology, Indications, Treatment and Specifics) would result in a clinically significant improvement in glycaemic control of type 2 Diabetes Mellitus (T2DM) patients.

METHODS: Patients who were diagnosed with T2DM at two tertiary care hospitals in Sri Lanka, comply with the selection criteria were enrolled to the study. The intervention consisted with two repeated one to one education sessions followed up in six and twelve months. HbA1c, lipid profiles, waist circumference, BMI and other biomedical measurements were done in both groups. Analysis of covariance between groups were conducted to determine the effectiveness of the intervention.

RESULTS: Mean HbA1c level in both intervention and usual care group was 8.6% with deviation from their target glycaemic level (6.5%, 48 mmol/mol) at baseline. At six months, there was a significant reduction ($p < 0.001$; size of effect = 0.69) in HbA1c between the intervention and the usual care group controlling the baseline values.

CONCLUSION: The results demonstrate the effectiveness of one to one diabetes self-management intervention among the adults with T2DM.

Key words: Diabetes self-management; Type 2 Diabetes; One to one patient education; HbA1c level

© 2018 The Author(s). Published by ACT Publishing Group Ltd. All rights reserved.

Cooray BPR, Morrissey H, Waidyarathnec EI, Ball PA. The Impact of Individual Health Education on Health Literacy: Evaluation of the Translated Version (Sinhala) of Health Education Impact Questionnaire in Type 2 Diabetes. *International Journal of Diabetes Research* 2018; 1(1): 1-6 Available from: URL: <http://www.ghrnet.org/index.php/ijhr/article/view/2272>

INTRODUCTION

Type 2 diabetes (T2DM) is a common chronic condition which, if not

controlled, can cause complications such as nephropathy, retinopathy and neuropathy. In recent years T2DM prevalence increased globally. There were 1.16 million cases of all types of diabetes were recorded in Sri Lanka in 2016 as stated in an International Federation of Diabetes Atlas^[1-3]. A significant reduction in diabetes-related complications has been recorded with the diabetes education in primary care setting, with the consideration of health beliefs and their cultural behaviours^[2-4].

However, a range of indices are used to measure health education in individuals and populations. Due to use of different, new or non-validated tools by researchers to assess the health education impact; there were few available evidence of a direct relationship between health education and improvement of patients' health outcomes^[5-7].

The health education impact questionnaire (heiQTM) is a validated tool that has been in use to evaluate the impact of patient education interventions on patient knowledge, behaviours, ability to act in an emergency, to navigate the health services and resourcefulness^[8,9]. The questionnaire has been used to measure health education impact on patients with a broad range of chronic diseases and validated in different settings, by a range of health professionals, which has been adapted to 20 languages^[9-11].

The heiQTM was derived from a grounded approach, with its most prominent feature being its breadth and capability to evaluate individual's ability to manage their condition irrespective of which chronic disease they have. This questionnaire covers eight main domains in 40 questions: health directed behaviour, positive and active engagement in life, emotional wellbeing, self-monitoring and insight, constructive attitudes and approaches, skills and technique acquisition, social integration and support and health services navigation and additionally with another section to evaluate the progress of the education program^[10-12].

Jayasuriya *et al.*, 2015 study concluded that improvement in glycaemic control was significant among Sri Lankan patients if they were modified through diabetes self-management interventions as in many other countries^[4]. However, Sri Lankans patients have poor understanding of the importance of dietary control, physical activity and adherence to therapy in the management of diabetes. This might be due to insufficient information they receive from their treating professionals^[2,13-15]. Additionally, Sri Lanka has remote and rural locations where government-specialist clinics are scarce. Those government specialist medical clinics available in urban areas are often overcrowded with patients. Thus, most T2DM patients are managed in primary care system, by general practitioners^[13-15].

However, Jayasuriya *et al.*, 2015 and a systemic review of Cooray *et al.*, 2017 revealed that there is a gap between the patient knowledge, management of diabetes and the effectiveness of patient education measured by validated questionnaires in Sri Lanka^[4,16,17].

Therefore, the aim of this study was to evaluate the impact of structured health education program on patients' knowledge by utilizing the heiQ questionnaire on their health-related behaviours, capabilities and motivations.

DESIGN AND METHODS

This study was designed as a prospective, non-experimental, step-up controlled study, with health education as the main intervention (Figure 1). To reduce samples cross contamination between the non-intervention and intervention groups cluster design method has been adopted. Thus, sample was recruited from two main independent tertiary care facilities in Western and Southern Provinces of Sri Lanka.

Both non-intervention and intervention groups participants continued to receive their usual diabetes care and treatment as indicated by their treating practitioners and the local guidelines for each of the two hospitals.

Potential participants were informed verbally and in writing about the study objectives and processes, their rights, and what was expected of them if they decided to participate. Participation was voluntary, with written consent obtained in Sinhala, their native language prior to participation.

The heiQTM instrument was translated (forward-backward translation) to Sinhala by professional translation services contracted by the developer, Osborne *et al.* (2007) at the University of Melbourne. The heiQTM was used to measure participants' knowledge and attitude towards their self-care, pre- and post- the delivery of a structured health-education session and compared with non-intervention group. The health education session was carried out one-on-one, verbal script-based consultation, aided by PowerPoint presentation by the primary investigator. The education program followed the Pathophysiology, Indication, Treatment and Specifics (PITS) patient education model^[17]. The model presents information in an organised and logical format that enhances the receiver's ability to recall information. Also, this model enables the receiver to easily follow the educators' thoughts and direction of information flow, which may be contingent on the emotional state of the patient or other constraints^[17].

Sample selection, control and randomisation

All genders with T2DM patients aged 18 years and over were approached regardless of level of education, socioeconomic level and employment status. As people under the age of 18 and those who are pregnant or diagnosed with polycystic ovarian syndrome or gestational diabetes or have severe diabetes complications who require different type of educational material and higher level of consent, they were excluded from the study.

Participant was enrolled if they were diagnosed with T2DM by the physicians who was not on haemodialysis and able to speak and understand Sinhala, who is not identified as having cognitive, hearing or vision impairment.

The study participants were approached on the basis of every third T2DM patient from the clinic registry for both non-intervention and intervention groups in two different tertiary care facilities in Sri Lanka, at the beginning of the study. Then, intervention group further categorised into subgroups (Subgroup A and Subgroup B) and it was carried out using a computer-generated randomising algorithm in Microsoft ExcelTM.




Figure 1 Sample Design.

Being a low-budget preliminary study, a total of 166 participants was considered as a manageable sample that would provide sufficient data to inform future studies on the trends, variances and relationships, between self-care behaviour, level of health education and diabetes control. The estimated withdrawal rate for the study was expected to be approximately 10%.

Data collection and analysis

Demographic data were collected using a socio demographic survey questionnaire designed by the investigator. The research method was based on two main aspects the clinical data collection and the diabetes health-education intervention. Clinical data collection took place on three occasions for all intervention and non-intervention participants – at the enrolment day, six-months and 12-months follow up – while diabetes education was carried out on the following basis: (1) At the enrolment day for intervention group (Subgroups A and B); (2) At the six-month follow up for participants in Subgroup B only.

Clinical and biomedical data collection was confined to measuring weight, height, waist circumference, BP, and blood sample collection for HbA1c and lipid profile tests.

Change in medications was not in the scope of this study; however patient adherence to therapy and self-care behaviours were monitored during the two follow up visits at six and 12 months.

The heiQ™ validated instrument used under the licensed agreement with Deakin University, Australia and printed heiQ™ baseline questionnaires were administered at the data collection sites; supported (explained without input) by the investigator, to measure the knowledge about their disease condition, and medication adherence before the intervention. Patients were asked to tick the most applicable answer describing their level of knowledge. The participants viewed the questions without the domain subtitle, allowing them to answer the individual questions based on their own perspective without being influenced by the domain title.

Following the completion of the questionnaire, the intervention group received the structured health-education program on one-on-one basis by the principal investigator. The program comprised of a presentation covering the pathogenesis, progression and complications of T2DM; importance of self-management and physicians follow-up. The session then progressed to focus on the participant prescribed medication, including the use of blood glucose monitors and insulin pens, how the medication work, their side effects and the benefit from adherence to medication on disease prognosis and development of complications. The same procedure was used for the data collection process of the non-intervention-group with the exclusion of the diabetic health-education intervention.

All enrolled participants (Subgroup A and B of intervention group and the non-intervention group) completed the heiQ™ follow up at 6 months and at twelve months, which included patient feedback on the education program and the way it was delivered. Completed questionnaires with their ID numbers were analysed and scored based on a Likert-type scale from 1 to 4, corresponding to <strongly disagree>, <disagree>, <agree> and <strongly agree>. Each domain has four to six questions. The scores were summated within each domain to obtain a scale score for each domain based on guidelines adopted by the questionnaire developer.

The participants' domain mean scores achieved in the baseline were compared to the follow-up mean scores and with non-intervention group to determine if their knowledge improved after the delivery of the patient education intervention using developers' guideline on effect size calculation and Analysis of Covariance (ANCOVA) using IBM® SPSS® (version 23) and R-Studio® (version 3.2.2) software packages.

Ethical approval

The study was approved by the Human ethics committee (Ref. No: HI4082) of Charles Darwin University, Australia and Ethics Review Committee (Ref No: 17.11.2014:3.32) of Faculty of Medicine, University of Ruhuna, Sri Lanka.

RESULTS

Overall analysis of demographics of 150 patients who were retained (out of 166 participants) at the end of study (retained rate 92%) indicate that large percentage of participants were women (72.3%, 120) with mean age of 56.2 years, with a standard deviation [SD] of 8.95. Most participants (85%) were managing their diabetes with oral hypoglycaemic agents. Among these patients, three-quarters of the T2DM patients were treated with lipid lowering agents, while half of the T2DM patients also received treatment for hypertension. In addition to dyslipidaemia and hypertension, a few participants were treated for other conditions such as thyroid disorders (three patients), arthritis (three patients) and one patient had cancer.

Table 1 summarises the comparison of categorical and important biomarker baseline values for intervention and non-intervention groups. Detailed baseline values for each sub group levels publish in a separate manuscript.

The analysis of heiQ™ based on the developers' guidelines indicate that one hundred and sixty-four heiQ™ forms were completed at baseline, 152 (92.7%) at the six months' follow-up and 150 (91.5%) at the end of 12 months' follow-up.

The results of this study based on analysis of covariance (ANCOVA) indicated significant improvement of the knowledge in all other domains between baseline to six months, six to twelve months and baseline to twelve months except for skills and technique acquisition between baseline to six months among intervention group in comparison to the non- intervention group (control group) (Table 2).

Further analysis results of sub groups A and B of intervention group and non- intervention group summarised in table 3 indicate the net group change effect size of the knowledge improvement calculated based on questionnaire developers' guidelines.

The effect size in each domain after the repeated education intervention (at 12 months) is summarised in Table 3. Domain 2: Positive and active engagement in life and Domain 1: Health directed behaviour showed greater improvement following the second follow up than the other domains. Domain 5: Constructive attitudes and approaches, Domain 7: Social integration and support and Domain 3: Emotional wellbeing showed improvement in the small to medium range. Although a positive effect was evident in the answers for Domain 4: Self-monitoring and insight, it was small.

As summarised in Table 3, Domain 8: Health services navigation and Domain 6: Skills and technique acquisition showed small and medium effect size changes while majority of the domains with quite low or negative effect.

Analysis of the second follow up, after 12 months, revealed a reduction of net positive reliable change in domains in the non-intervention group.

The assessment of health education program based on nine follow-up questions from those who completed the education interventions, mean score was calculated. Figure 2 shows that Question 45 "It was worth my time to take part in this study" scored the highest, with a score of 5.55, while all the questions received an average score of more than five. This finding confirms the success of the education intervention.

Table 1 One-Way ANOVA results for comparing the mean measurement values of intervention group vs non-intervention group at baseline analysis.

Measurement	Intervention		Non-intervention		F value	Sig.
	Mean	Standard Deviation (SD)	Mean	Standard Deviation (SD)		
Age (yr)	56	8.9	56	9.2	0.669	0.514
Weight (kg)	59	10.3	61	9.1	0.954	0.387
Height (cm)	154.4	8.4	157.3	7.6	3.185	.044*
Waist Circumference (cm)	95.5	9.1	96.8	7.9	0.598	0.551
HbA1c level (%)	8.56	1.62	8.65	1.53	0.279	0.757
Systolic Pressure (mmHg)	126.3	16.8	129.4	16.7	0.651	0.523
Diastolic Pressure (mmHg)	79.6	9.4	84.9	9.1	5.97	0.003**
Heart rate (bpm)	78.4	11.6	81.1	11	1.899	0.153
Total Cholesterol (mg/dL)	189.8	44.1	181.3	38.9	0.908	0.406
HDL Cholesterol (mg/dL)	45.1	8.5	45	8.9	0.097	0.908
Triglycerides (mg/dL)	126.3	64.4	129.1	46.7	0.25	0.779
LDL Cholesterol (mg/dL)	100.1	35.2	109.5	35.9	1.33	0.267
BMI Value (kg/m ²)	24.8	4.3	24.7	3.5	0.075	0.928

*Between group effect is statistically significant at the $p < 0.05$ level, ** Between group effect is statistically significant at the $p < 0.01$ level.




Figure 2 Evaluation of the health-education program.

DISCUSSION

The findings from the baseline analysis indicated that all participants, intervention and non-intervention, shared a similar starting point with their health and physical measurements and diabetes knowledge.

The results of this study reveal that all heiQ™ domains showed at least low to moderate correlations with the follow ups. The initial mean values in the current study were comparative to the results presented in the French translated heiQ™ data^{18,19}. This difference may indicate that the Sri Lankan participants already had limited knowledge of their disease. However, this exploration of the positive improvement in all domains in the intervention group during the

follow-up analysis compared to the non-intervention group, strongly evidences the effect of the education intervention on knowledge improvement in the study participants. Domains 1, 3 and 6 showed higher results in this study than that of the previous studies validated in the French translated heiQ™. However, results were in a middle range between the two studies for Domains 2, 4, 5, 7 and 8.

A study by Osborne *et al.* (2011), the developer of the heiQ™, concluded that it is common for people who accumulate a high heiQ™ score in Domain 3: Emotional wellbeing to also have high scores in Domain 2: Positive and active engagement in life and a low score in Domain 4: Self-monitoring and insight. This current study

validated this finding. The study sample revealed that Domains 8 and 7: Health services navigation and Social integration and support respectively, had the lowest scores. This result might be due to the poor development of technology, or less priority given to personal conditions than family matters as per cultural beliefs in adults in Sri Lanka. These findings evidence by the Amarasekara et al, 2014 study of health beliefs and practices in Sri Lanka.

The greatest effect size was found in Domain 1: Health directed behaviour, Domain 2: Positive and active engagement in life and Domain 8: Health services navigation, between the baseline score and the follow ups in the intervention group. Domain 5: Constructive attitudes and approaches, Domain 6: Skills and technique acquisition, Domain 7, Social integration and support and Domain 3 Emotional wellbeing were also notable in the way intervention group individuals participated in the education session in the follow ups compared to

that delivered at baseline. Intervention group participants also showed an improvement in their ability to confidently interact with a range of health organisations and health professionals (the researcher, nurses, GPs and the pathology laboratory). These findings are critical for patients in Sri Lanka, as they have high dependence on physicians' instructions and on medical healthcare professionals to manage their health conditions. This could increase the overcrowded issues in most state sector hospitals in Sri Lanka as revealed by the Medagama et al., 2015 study. In addition, these findings reveal that the participants were encouraged to become proactive in managing their health.

However, there was a slight reduction in skill and technique acquisition and health service navigation at six months, after twelve months. These findings seem reasonable as people with worse health may have had more experience or challenge with these areas and been more engaged with health professionals to control their

Table 2 Summary of between-intervention and non-intervention group effects for eight domains at baseline, 6 months and 12 months.

Domains	Time interval (months)	Groups	Mean value	F Statistic	Significance	Size of Effect
Health directed behaviour	0-6 months	Intervention	2.82	24.25	<0.001**	0.138
		Non-intervention	2.73			
	6-12 months	Intervention	3.09	65.75	<0.001**	0.309
		Non-intervention	2.65			
	0-12 months	Intervention	3.35	80.78	<0.001**	0.355
		Non-intervention	2.68			
Positive and active engagement in life	0-6 months	Intervention	2.84	15.88	<0.001**	0.095
		Non-intervention	2.8			
	6-12 months	Intervention	3.09	13.89	<0.001**	0.086
		Non-intervention	2.83			
	0-12 months	Intervention	3.2	23.07	<0.001**	0.135
		Non-intervention	2.86			
Self-monitoring and insight	0-6 months	Intervention	2.97	13.27	<0.001**	0.081
		Non-intervention	2.96			
	6-12 months	Intervention	3.14	23.09	<0.001**	0.137
		Non-intervention	2.89			
	0-12 months	Intervention	3.21	28.16	<0.001**	0.161
		Non-intervention	2.82			
Constructive attitudes & approaches	0-6 months	Intervention	2.92	45.11	<0.001**	0.229
		Non-intervention	2.96			
	6-12 months	Intervention	3.16	7.83	0.006**	0.05
		Non-intervention	2.81			
	0-12 months	Intervention	3.23	14.83	<0.001**	0.091
		Non-intervention	2.84			
Skills and technique acquisition	0-6 months	Intervention	2.78	n.s	n.s.	n.s.
		Non-intervention	2.75			
	6-12 months	Intervention	3.1	29.14	<0.001**	0.165
		Non-intervention	2.95			
	0-12 months	Intervention	3.23	32.98	<0.001**	0.182
		Non-intervention	2.68			
Social integration & support	0-6 months	Intervention	2.86	13.21	<0.001**	0.08
		Non-intervention	2.84			
	6-12 months	Intervention	3.11	14.55	<0.001**	0.089
		Non-intervention	2.86			
	0-12 months	Intervention	3.17	21.39	<0.001**	0.126
		Non-intervention	2.9			
Health services navigation	0-6 months	Intervention	2.66	11.28	0.001**	0.092
		Non-intervention	2.64			
	6-12 months	Intervention	3.15	24.22	<0.001**	0.141
		Non-intervention	2.85			
	0-12 months	Intervention	3.14	31.39	<0.001**	0.176
		Non-intervention	2.74			
Emotional wellbeing	0-6 months	Intervention	2.4	15.28	0.001**	0.092
		Non-intervention	2.41			
	6-12 months	Intervention	2.13	17.11	<0.001**	0.107
		Non-intervention	2.45			
	0-12 months	Intervention	2.01	21.91	<0.001**	0.133
		Non-intervention	2.44			

** The mean difference is significant at the 0.05 level, n.s. not significant

Table 3 Summary of between-intervention subgroups (A and B) and non-intervention group effects for eight domains at baseline to 12 months.

Domain	Intervention group					Non-intervention group		
	Baseline mean	Sub group A with one intervention		Sub group B with two interventions		Baseline mean	12 months mean	Group change effect size
		12 months mean	Group change effect size	12 months mean	Group change effect size			
Health directed behaviour	2.82	3.09	0.50**	3.35	0.99***	2.73	2.65	-0.22
Positive and active engagement in life	2.84	3.09	0.54**	3.2	0.81***	2.8	2.83	0.07
Self-monitoring and insight	2.97	3.14	0.36*	3.21	0.48*	2.96	2.89	-0.17
Constructive attitudes & approaches	2.92	3.16	0.40*	3.23	0.50**	2.96	2.81	-0.33
Skills and technique acquisition	2.78	3.1	0.59**	3.23	0.73**	2.75	2.95	0.43*
Social integration & support	2.86	3.11	0.44*	3.17	0.54**	2.84	2.86	0.05
Health services navigation	2.66	3.15	0.85***	3.14	0.88***	2.64	2.85	0.55**
Emotional wellbeing	2.4	2.13	-0.45*	2.01	-0.68**	2.41	2.45	0.06

* small effect size, ** medium effect size, *** greater effect size.

conditions, whilst they become less responsive when they get used to their disease condition.

There was no revision made to the translated version of the heiQ™ as all items precisely promote health and behavioural change. However, based on verbal communication with the study participants, the terminology used the items translated from “I feel hopeless because of my health problems” and “I feel like I am actively involved in my life” were shown not to be fully understood, and they were answered after clarification by the investigator. Some clarification was made to reduce variability in results. Therefore, further research is encouraged using the modifications to the translation of those two statements.

The follow-up questions were based on the fundamental environmental and personal determinants of a person’s opportunities to access the research intervention and provide their feedback. Question 45: “It was worth my time to take part in this study” scored the highest, with a score of 5.58. It was one of the main outcomes of the education intervention that had positive feedback and highlights the success of the education intervention.

CONCLUSION

The heiQ™ translated into Sinhala was well accepted by participants suffering from T2DM, revealing the usefulness of the questionnaire as an instrument to evaluate the impact of health-education intervention for any chronic disease, not just diabetes. Sinhala heiQ™ serves as a proximal goal for self-management programs to advance outcome assessment in this field in Sri Lanka.

Practical implications and recommendations

The method used in this study through the use of the heiQ™ can identify individual patient education needs and address relevant aspects to their education status and health education gaps. Thus, current study intervention could be used as a model of an integrated program to improve T2DM patient awareness and medication self-management by incorporating it to the current T2DM management program. Additionally, the heiQ™ could be used as a tool for further studies to determine the value of its application in other health conditions, to compare outcomes across multiple cultures and languages in Sri Lanka.

ACKNOWLEDGEMENTS

I would like thank all medical, nursing and laboratory staff; and all participants from Karapitiya Teaching Hospital and Colombo South Teaching Hospital, Kalubowila. Special thanks to Charles Darwin

University, Australia and University Grant Commission Sri Lanka for providing scholarship and grants to support for first author’s study.

Registration of study

This study was registered under the Sri Lanka Clinical Trial Registry, No: SLCTR/2015/014, which is a primary registry for clinical trials involving human subjects conducted in Sri Lanka that is linked to the Registry Network of the International Clinical Trials Registry Platform of the WHO. The site approval was obtained from Directors of both tertiary care facilities in Sri Lanka.

REFERENCES

- Katulanda P, Constantine GR, Mahesh JG, Sheriff R, Seneviratne RD, Wijeratne S, Wijesuriya M, McCarthy MI, Adler AI, Matthews DR. (2008) Prevalence and projections of diabetes and pre-diabetes in adults in Sri Lanka--Sri Lanka Diabetes, Cardiovascular Study (SLDCS). *Diabetic Medicine* 25(9): 1062-1069. doi: 10.1111/j.1464-5491.2008.02523.x
- Amarasekara AA, Fongkaew W, Turale S, Wimalasekara SW, Chanprasit C. (2014) An ethnographic study of diabetes health beliefs and practices in Sri Lankan adults. *International Nursing Review* 61: 507-514. doi: 10.1111/inr.12136.
- International Diabetes Atlas 8th Edition Online 2017. Brussels, Belgium International Diabetes Federation. ISBN 978-2-930229-87-4. Access date 09/05/2018. available at: http://diabetesatlas.org/IDF_Diabetes_Atlas_8e_interactive_EN/
- Jayasuriya R, Pinidiyapathirage MJ, Jayawardena R, Kasturiratne A, de Zoysa P, Godamunne P, Gamage S, Wickremasinghe AR. (2015). Translational research for Diabetes Self-Management in Sri Lanka: a randomized controlled trial. *Primary Care Diabetes* 9(5): 338-345. doi: 10.1016/j.pcd.2015.01.014
- Osborne RH, Batterham R and Livingston J. (2011) The evaluation of chronic disease self-management support across settings: the international experience of the health education impact questionnaire quality monitoring system. *Nurs Clin North Am* 46(3): 255-270. doi: 10.1016/j.cnur.2011.05.010
- Choi TST, Walker KZ and Ralsto RA. (2015) Diabetes education needs of Chinese Australians: A qualitative study. *Health Education Journal* 74(2): 197-208. doi: 10.1177/0017896914523276
- Jordan JE, Osborne RH and Buchbinder R. (2011) Critical appraisal of health literacy indices revealed variable underlying constructs, narrow content and psychometric weaknesses. *Journal of Clinical Epidemiology* 64(4):366-379. doi: 10.1016/j.jclinepi.2010.04.005.
- Osborne RH, Elsworth GR and Whitfield K. (2007) The Health Education Impact Questionnaire (heiQ): an outcomes and evaluation measure for patient education and self-management

- interventions for people with chronic conditions. *Patient Education and Counselling* 66(2):192-201. doi: 10.1016/j.pec.2006.12.002
9. Morita R, Arakida M, Osborne RH, Nolte S, Elsworth GR and Mikami H. (2013) Adaptation and validation of the Japanese version of the Health Education Impact Questionnaire (heiQ-J) for the evaluation of self-management education interventions. *Japan Journal of Nursing Science* 10(2): 255-266. doi: 10.1111/j.1742-7924.2012.00224.x.
 10. Epstein J, Osborne RH, Elsworth GR, Beaton DE and Guillemin F.(2015) Cross-cultural adaptation of the Health Education Impact Questionnaire: experimental study showed expert committee, not back-translation, added value. *Journal of Clinical Epidemiology* 68(4): 369-369. doi: 10.1016/j.jclinepi.2013.07.013.
 11. Osborne RH, Batterham RW, Elsworth GR, Hawkins M and Buchbinder R. (2013) The grounded psychometric development and initial validation of the Health Literacy Questionnaire (HLQ). *BMC Public Health* 13: 658-675. doi: 10.1186/1471-2458-13-658
 12. Elsworth GR, Nolte S and Osborne RH. (2015) Factor structure and measurement invariance of the Health Education Impact Questionnaire: Does the subjectivity of the response perspective threaten the contextual validity of inferences? *SAGE Open Medicine* 3(1):1-13. doi: 10.1177/2050312115585041.
 13. Pinidiyapathirage MJ, Kasturiratne A, Ranawaka UK, Gunasekara D, Wijekoon N, Medagoda K, Perera S, Takeuchi F, Kato N, Warnakulasuriya T, Wickremasinghe AR. (2013). The burden of diabetes mellitus and impaired fasting glucose in an urban population of Sri Lanka. *Diabetic Medicine* 30(3): 326-332. doi: 10.1111/dme.12013.
 14. Ranasinghe DC, Ranasinghe P, Jayawardena R, Matthews DR and Katulanda P. (2014). Evaluation of physical activity among adults with diabetes mellitus from Sri Lanka. *International Archives of Medicine* 7(1): 15-19. doi: 10.1186/1755-7682-7-15
 15. Medagama AB, Bandara R and Wijethunga R. (2015). The high burden of in-hospital diabetes mellitus at a tertiary care hospital in Sri Lanka: a case control study. *Journal of Diabetes & Metabolism* 06(03):502-506. doi: 10.4172/2155-6156.1000502
 16. Cooray BPR, Ball PA, Morrissey H and Waidyaratne EI. (2017) The effect of health education in the management of Type 2 Diabetes and the Sri Lankan perspective: A Review. *Sri Lanka Journal of Diabetes, Endocrinology and Metabolism* 7(1):33-38. doi: 10.4038/sjdem.v7i1.7325
 17. Stewart NM (2012) Practical Patient Literacy, The Medagogy Model. McGraw-Hill Education / Medical; 1 edition (16 Feb. 2012) pp230 ISBN 978-0071761314 available online at: <https://smy64gw01.storage.googleapis.com/MDA3MTc2MTMxNA==01.pdf> access date 09/05/2018
 18. Morita, R, Arakida, M, Osborne, RH, Nolte, S, Elsworth, GR, Mikami, H. (2013). Adaptation and validation of the Japanese version of the Health Education Impact Questionnaire (heiQ-J) for the evaluation of self-management education interventions. *Japan Journal of Nursing Science*, 10(2), 255-266. doi: 10.1111/j.1742-7924.2012.00224.x
 19. Belanger, A, Hudon, C, Fortin, M, Amirall, J, Bouhali, T, Chouinard, MC. (2015). Validation of a French-language version of the health education impact Questionnaire (heiQ) among chronic disease patients seen in primary care: a cross-sectional study. *Health & Quality of Life Outcomes*, 13(1), 64. doi: 10.1186/s12955-015-0254-0