

Revisiting Innovation Practices in Subsistence Farming: The Net Effects of Land Management, Pesticide, Herbicide and Fungicide Practices on Expected Crop Harvest in Ethiopia

Adah-Kole Emmanuel Onjewu

Wolverhampton Business School, University of Wolverhampton, UK.

a.onjewu@wlv.ac.uk

Vahid Jafari-Sadeghi

Newcastle Business School, Northumbria University, UK.

vahid.jafari-sadeghi@northumbria.ac.uk

Sundas Hussain

Nottingham Business School, Nottingham Trent University, UK.

sundas.hussain@ntu.ac.uk

Abstract

To settle inconsistent findings in the farming innovation and productivity nexus, this inquiry examines the land management practices of 7,625 households in rural Ethiopia. Specifically, the net effects of (1) improved seeds, (2) mixed cropping and (3) row planting on the use of (4) pesticides, (5) herbicides and (6) fungicides are assessed. Using a structural equation technique, the study probes how these six practices predict households' expected harvest. It is found that while improved seeds increase pesticide, herbicide and fungicide use, mixed cropping and row planting generally reduce these practices. Moreover, mixed cropping moderately increases expected harvest while improved seeds and row planting have the reverse effect. The interrelations of these factors increase knowledge in contingency-driven agronomics, and provoke reflection on the sustainability of land management practices. Particularly, opposed to prevailing views, it is demonstrated that sowing traditional seeds will reduce households' reliance on pesticides, herbicides and fungicides. The inherent findings speak to policy-makers tasked with supporting peasant life in rural Ethiopia and similar contexts.

Keywords: *Improved Seeds; Mixed Cropping; Row Planting; Pesticides; Herbicides; Fungicides; Expected Harvest; Farming Innovation; Subsistence Farming; Structural Equation Modelling; Ethiopia.*

Biographical Notes:

Dr Adah-Kole Emmanuel Onjewu is a Senior Lecturer in International Business at the University of Wolverhampton. He holds a PhD in Business and Management from the University of Plymouth and his research interests are entrepreneurial behaviour, entrepreneurial education, family businesses, export performance and tourism. His publications investigate how firm and individual behaviours yield positive household, firm and socio-economic outcomes for development in the African, Middle-East and similar less developed contexts. He has published in several journals including *Annals of Tourism Research*, *Journal of Business Research*, *Studies in Higher Education*, *Technological Forecasting & Social Change*, *Critical Perspectives on*

International Business and Industry & Higher Education. He is a fellow of the UK's Higher Education Academy.

Dr Vahid Jafari-Sadeghi is a Senior Lecturer in International Business Strategy at Newcastle Business School, Northumbria University. He holds a PhD in International Entrepreneurship from the University of Turin where he was also a post-doctoral fellow. Dr Jafari-Sadeghi has been a visiting researcher at the University of Regina and collaborates with colleagues on several projects in various universities. He has published in several reputable journals such as *International Business Review*, *Journal of International Entrepreneurship*, *Research in International Business and Finance*, and the *International Journal of Entrepreneurship and Small Business*. He has also served as guest editor and reviewer for several journals and chaired a number of tracks in international conferences.

Dr Sundas Hussain is a Senior Lecturer in Management at Nottingham Business School, Nottingham Trent University. Her PhD examined entrepreneurship support for socially disadvantaged women in collaboration with a Birmingham-based housing association. Dr Hussain also leads a small team as a Climate Director with Woodfarm Education Centre, a Scottish registered charity. Since 2015, the centre has successfully delivered several Climate Challenge Fund projects financed by the Scottish government. Grants received range from £100,000 to £250,000 for climate change projects aiming to reduce CO₂ emissions and the carbon footprint of the local community. Her other affiliations include membership of the Gender and Enterprise as well as Entrepreneurship in Minority special interest groups at the Institute for Small Business and Entrepreneurship, and fellowship of the Higher Education Academy.

1.0 Introduction

In Africa, as in other underdeveloped settings, suitable conditions and access to resources for subsistence farming are key to household survival (Baipheti and Jacobs, 2009). Local practices for food production and preservation ensure availability of nutrition and the reduction of poverty (Rankoana, 2017). In spite of mounting urban migration, the United Nations' Food and Agriculture Organisation [FAO] (2015) still cites subsistence farming by families as a key determinant for well-being and a source of livelihood in rural Africa. Similarly, Jayne *et al.* (2019) indicate that small-scale holdings managed by families comprise the vast majority of Africa's rural dwellers, and Bjornlund *et al.* (2020) estimate this to be in the range of 50-70% of the population. To somewhat quantify their contribution, smallholdings engaging family labour feed and employ two-thirds of the African population and cultivate 62% of the land (FAO, 2021a).

Notwithstanding households' immense endeavour, agricultural production across sub-Saharan Africa has lagged behind global output since the 1960s (FAO, 2009a), and crop harvests remain significantly lower than in other developing regions to the extent that food security is presently threatened (Bjornlund *et al.*, 2020). To be sure, there is no shortage of enterprise on the part of households but general underperformance can be explained by poor harvests arising from factors including crop phenology, pests, herbivores, poor soil, rodents,

wildlife trespassing and unpredictable rainfall (Runo *et al.*, 2011; Stokstad, 2017; Gross *et al.*, 2018). Recently, *Spodoptera Frugiperda* (or fall armyworm), a pest previously found only in North and South America, has been sighted in Africa (Sisay *et al.*, 2018). If left uncontrolled, fall armyworms can destroy maize and other staple crops up to a value of US\$3 billion in one year (Stokstad, 2017).

Strikingly, fall armyworms are only one of several invasive alien species [or IAS hereafter] causing crop damage in Africa. Other IAS on the continent are *Zonocerus Variegatus* [variegated grasshopper], *Spodoptera* [black armyworm], *Cryphonectria Parasitica* [chestnut blight], *Phytophthora Cinnamomic* [root rot], *Odontotermes Obesus* [fungus-farming termites], *Acacia Mearnsii* [black wattle] desert locust, brown streak and wheat rust (Banjo *et al.*, 2010; Graziosi *et al.*, 2020; FAO, 2021b). In East Africa, the five leading microbes plaguing maize farms are *Chilo Partellus* [spotted stem borers], Maize Lethal Necrosis Disease [MLND], *Parthenium Hysterophorus* [flowering plant], *Liriomyza* [leaf miner flies] and *Tuta Absoluta* [tomato leaf miner] (Loha *et al.*, 2018). In 2016, these five IAS were jointly responsible for a combined crop loss valued at US\$0.9-1.1 billion across Ethiopia, Kenya, Malawi, Rwanda, Tanzania and Uganda (Pratt *et al.*, 2017). This was a direct squeeze on the productivity and sustenance of rural households.

To manage IAS pitfalls, subsistence households resort to the use of pesticides, herbicides and fungicides [or xenobiotics hereafter] to increase harvests (Tolera, 2020). This practice follows other low-tech and pragmatic choices made to optimise crop production such as improved seedlings [versus traditional seedlings] (Ahmed *et al.*, 2017), mixed cropping [versus monocropping] (Demlew *et al.*, 2019) and row planting [versus broadcast planting] (Vandercasteelen *et al.*, 2018). These routines are of particular import because of the increasing scarcity of fertile ground as 'over 75% of arable land in Africa is considered degraded' (Orchard *et al.*, 2017, p. 46). Furthermore, to heed Emerton and Snyder (2018), it is vital for scholars to understand households' sustainable land management practices and the economic effects that arise as a consequence.

Ensuing from the above, the aim of this study is to test the effect of improved seeds, mixed cropping and row planting on the use of xenobiotics and the extent to which these correlations explain expected harvest. Along these lines, we offer a novel link to advance scholars' understanding of land management practices in a setting where crop damage is an issue of serious concern to world bodies not limited to the FAO. Our consequent contributions are fourfold. First, we provide original evidence on the influence of improved seeds, mixed cropping and row planting on the use of xenobiotics. To the best of our knowledge, no prior

studies have assessed the interplay of these practices. Second, we examine how improved seeds, mixed cropping and row planting distinctly impact on expected harvest. In spite of the prevalence of these land practices, their association with expected harvest has also not been conceptualised in extant work. Third, to explain households' behaviour, we leverage the contingency and sustainability lens to add theoretical perspective to the interpretation of the findings. Last, we examine rare first-hand data from a large sample of household farmers in Ethiopia. The granular nature and size of this sample yields both direct and representative insights into land management practices to better steer policy-making.

To press forward, the overarching question addressed in this inquiry is: 'How do improved seeds, mixed cropping and row planting affect the use of xenobiotics and expected harvest in turn?'. The rest of this article is arranged as follows: Section 2 presents the research context while section 3 appraises the six land management practices currently conceptualised to predict expected harvest. Subsequently, section 4 explains the measurement variables, items and scales before findings are offered in section 5. In section 6, the findings are then compared with prior findings by way of a discussion. We conclude with theoretical contributions, practical implications and areas for future research in section 7.

2.0 The Ethiopian Context

This study is set in a landlocked country in the horn of Africa formally recognised as the Federal Democratic Republic of Ethiopia. It is bordered by Eritrea to the north, Sudan to the north-west, South Sudan to the west, Djibouti and Somaliland to the northeast, Somalia to the east and Kenya to the south. Its largest city and capital is Addis Ababa and, with 12 cities in total, Ethiopia covers a total land area of 420,000 square miles (Ayalew *et al.*, 2016). It is home to 112 million inhabitants (World Bank, 2021a), making it the second most populous African country after Nigeria. Its gross domestic product (GDP) was US\$95.9 billion in 2019 and the economy is forecast to grow by 8.7% in 2022 (World Bank, 2021a). The agriculture, fishing and forestry sector generated US\$870 million in 2018, eclipsing the previous high of US\$790 million in 2009 (World Bank, 2021b).

Ethiopia warrants attention for a number of reasons. Notably, the vast majority of the 112 million Ethiopians reside in rural areas and remoteness is still a dominant characteristic of life (Abate *et al.*, 2020). Schmidt *et al.* (2018) estimated that, in 2015, over 20% of Ethiopians lived in areas that required more than five hours of travel to a city of 50,000 people. Ethiopia is also interesting to study because African observers and commentators view the country as the archetypal developmental state and a model for social mobilisation on the continent (Berhanu and Poulton, 2014; Clapham, 2018). Jayne *et al.* (2019) affirm that Ethiopia is the

closest example of smallholder-led agricultural growth in Africa with a 6% real annual average increase in production from 2000 to 2015. Yet, while there is evidence of technological solutions aiding the delivery of social services in Ethiopia, agriculture has not kept pace with this advancement (Berhane *et al.*, 2018). Indeed, numerous multi-country studies have shown that the adoption rate of improved technology is much lower in rural areas where the need is greatest (Jacoby, 2000; Christiaensen *et al.*, 2003; Deichmann *et al.*, 2009; Stifel *et al.*, 2017; Damania *et al.*, 2017). This is a plausible assumption in the context of Ethiopia's largely rural population.

Turning to xenobiotics, although the use of pesticides in Ethiopia was historically low, recent growth in food production and the expansion of the floriculture industry (Getu, 2009) have resulted in higher utilisation to the extent that the country now has 'the largest accumulation of pesticides in Africa' (Dawud *et al.*, 2019, p. 197). According to the Ministry of Agriculture and Rural Development [MoARD], in 2007, there were an estimated 250 sites holding 1,500 tons of pesticides (MoARD, 2007). More recently, Shita *et al.* (2018) enumerated common areas of farming innovation in Ethiopia including seed varieties, water irrigation, inorganic fertilizers and pesticides. On the latter, 'the total pesticide applied area reached 3.2 million hectares [22.32% of the total farmed land] in 2014/15 which has increased over time' (Shita *et al.*, 2018, p. 101). Kebede *et al.* (2017) also cited row planting as a popular farming innovation supported by the Ethiopian government.

Thus, the above-mentioned undercurrents aggregate to make Ethiopia a fertile ground for investigating land management practices that will advance understanding of households' productivity. The resulting analysis and findings will support scholarly reflections on national output from the ground up in a country believed in some quarters to be Africa's beacon of prosperity (Demiessie, 2021).

3.0 Theoretical Background and Hypotheses Development

Schoar (2010) draw attention to the distinctiveness of subsistence entrepreneurship as activities undertaken mainly to sustain family units. Indeed, Venugopal *et al.* (2015, p. 235) assert that 'more than a billion entrepreneurs worldwide live in subsistence contexts and run microenterprises to meet life's basic consumption needs'. Accordingly, there is a view that subsistence farmers in Africa do not exhibit the entrepreneurial propensity that is needed for growth or more transformational entrepreneurship (Cieslik and D'Aoust, 2018). Therefore, Schoar (2010) believes that the motivations for persisting with subsistence entrepreneurship, like household farming, are (1) capital constraints in terms of the resources available for production and (2) the labour market or manpower that can be committed to such activities.

These dimensions and related factors can be further understood through the contingency theory and sustainability perspective.

3.1 Contingency Theory and Sustainability

Recognising the scale of damage that could be caused by IAS, households' use of xenobiotics, in spite of added costs and other perceived disadvantages, can be explained by pressing circumstances causing them to seek satisfactory rather than optimal results. To understand this behaviour, there is a time-honoured belief that it is normal for economic agents to incessantly assess their situation and adapt, discard and replace their practices to increase rent (Sargent, 1993; Arthur, 1994; Holland *et al.*, 1986). This evokes the contingency perspective which suggests that the 'most effective technique depends on the set of circumstances at a particular point in time' (Luthans and Stewart, 1977, p. 182). Luthans and Stewart (1977, p. 183) added that contingency is generically situational in orientation and warrants 'identifying and developing functional relationships between environmental, management and performance variables'. In this vein, households' use of xenobiotics to reduce crop damage and increase harvest upholds contingency logic. Thus, espousing Luthans and Stewart's (1977) general contingency framework, the current inquiry views households as entities with limited resources operating within an environment where sustenance is dependent on maximising the performance of the land.

Furthermore, Luthans and Stewart's (1977, p. 186) general contingency framework embraced environmental variables 'with which the manager must interact and operate'. To some degree, this conjures Elkington's (1994) sustainability theory for ensuring the welfare of people, protecting the natural environment and raising profitability/performance. Although there are myriad definitions of sustainability, sustainable communities scholars describe the domain as economic growth that simultaneously solves environmental and local problems faced by people (Hempel, 1999). On this basis, Slaper and Hall (2011) expand Elkington's (1994) sustainability stance as having economic, environmental and social dimensions. According to them [Slaper and Hall, 2011], the measures of the economic dimension are personal income, firm size, employment distribution and revenue generation. For the environmental dimension, the measures comprise pollutants, solid and hazardous waste management, and change in land use. In terms of social measures, Slaper and Hall (2011) specified employment rate, female labour force participation, average household income, relative poverty and health-adjusted life expectancy. Altogether, these three dimensions and their measures correspond with Sissons *et al.*'s (2019) understanding of inclusive growth. Even for rural households, meeting current needs without compromising the ability of future generations to meet theirs is imperative.

Moreover, bearing in mind improved seeds, mixed cropping, row planting and the use of xenobiotics as contingent antecedents, it may be possible to explain findings on the extent to which these practices predict expected harvest through the economic, environmental and social dimensions of sustainability. This follows precedent in previous research investigating similar outcomes (e.g. van der Meulen *et al.*, 2014; Armanda *et al.*, 2019). We now proceed to appraise the antecedents in view of hypotheses development.

3.2 Improved Seeds

Seeds are a basic agricultural input and access to this commodity in its traditional or adapted form is a prerequisite for sustainable production (Abebe and Alemu, 2017). Abebe and Alemu (2017) add that the availability of seeds combined with other inputs is important for guaranteeing household food security, and improved seeds are essential for raising crop productivity. The widespread adoption of improved seeds stemmed from the 'technical package' of the green revolution first conceived in the United States in the late 1960s (Lakshman, 1993, p. 259). The FAO (2009) has since defined improved seeds as the genetic modification of seedlings aimed at increasing the quality and production of crops through increased drought tolerance, high yields and early maturity. Indeed, genetically improved seeds are 'widely understood to be a beneficent technology that dramatically increases agricultural output' (Lakshman, 1993, p. 255). Thus, 'in the last decade, the number of African countries researching and growing genetically modified seeds has increased fourfold' (Rock, 2018, p.15). The two kinds of improved seeds are hybrid and open-pollinated (Bachewe *et al.*, 2018). Although hybrid seeds increase yields considerably, the gains rapidly decline after the first year and this induces farmers to acquire new seeds. However, open-pollinated seeds do not exhibit such sharp decline in yield and are recommended to be replenished once every 3-4 years (Bachewe *et al.*, 2018).

In Ethiopia, the adoption of improved seeds is generally low (Abay *et al.*, 2017; Gebre *et al.*, 2019), but national initiatives like the Tropical Legume Development Program have nevertheless increased the rate of use (Verkaart *et al.*, 2019). The basis for investigating the effect of improved seeds on exigent land management practices is supported by McGuire and Sperling's (2011) contention. They stress that 'seed is a vital input but farmers' production and food security are likely to be affected more by ownership of assets, use of other inputs, or climate than by small fluctuations in seed availability' (McGuire and Sperling, 2011, p. 498). Furthermore, consistent with complementarity effects, agricultural yields could be higher when inputs are adopted simultaneously rather than separately (Feder, 1982). Besides, 'in the context of Ethiopia, agricultural input supply strategies in the last decade encouraged farmers to adopt chemical fertilisers and improved seeds as a package' (Bachewe *et al.*, 2018, p. 288),

signalling the interrelatedness of farming innovation. Prior research including Dorfman (1996), Gebremariam and Tesfaye (2017) and Bachewe *et al.* (2018) have modelled farmers' adoption of improved seeds alongside chemical fertilisers and irrigation inputs. Mirroring their precedent, the first set of hypotheses seek to estimate whether households' seed selection practices predict specific xenobiotic use:

H1. *Improved seeds are positively associated with the use of pesticides*

H2. *Improved seeds are positively associated with the use of herbicides*

H3. *Improved seeds are positively associated with the use of fungicides*

3.3 Mixed Cropping

Mixed cropping is the established practice of inter-planting crops in smallholdings (Maxwell and Fernando, 1989; Omamo, 1998). Proponents believe that mixed cropping systems reduce exposure to pests and the probability of total crop damage (Ezulike and Igwatu, 1993; Fondong *et al.*, 2002). Other reported benefits of mixed cropping are an improvement in soil cover (Howeler and Cadavid, 1990), the suppression of weeds (Olasantan *et al.*, 1994), improved soil fertility (Lusembo *et al.*, 1998) and greater labour efficiency (Odurukwe and Ikeorgu, 1994). Subsistence farmers may also gain from a more diverse diet and the possibility of additional income (Gold, 1993). In spite of these claims, there is a counter argument that 'yields of cash-crops appearing in mixed stands often fall well below those in pure stands' (Omamo, 1998, p. 155). In fact, Daellenbach *et al.* (2005) showed that mixed cropping systems reduce total production when compared to mono cropping. Yet, Yigezu *et al.* (2019) think otherwise and assert that, at least in legume-cereal mixed stands, there are clear economic advantages over pure stands as mixed stands provide higher yields and gross margins. All things considered, it would seem that these outcomes and overall performance are farm specific and contingent on other inputs like specialised machinery (Cook and Weller, 2004).

Acknowledging the heterogeneity of mixed versus pure stands, the correlations of alternate stand types with other ancillary practices have long drawn scholars' interest. Theoretically, Rhoades and Bebbington (1990) took an ecological and economic view to explain farmers' intercropping decisions. First, for ecology, they considered the pursuit of environmental reliance, stability and diversity as motivators. Second, for the economic stance, they cited risk reduction and the efficient use of land and labour as reasons for tending single or multiple crops. There is also a nutritional view, albeit from an intensive farming context, extolling an increase in 'nutrient productivity due to beneficial interactions between neighbouring plants' (Zhang *et al.*, 2019, p. 2). Jacques and Jacques (2015) concur that there are nutritional implications and biodiversity advantages that may accrue from making appropriate stand [monocropping versus mixed cropping] choices. In terms of the measurement of this practice, previous works have operationalised stand type to understand its effects on seed germination

(Dubey and Fulekar, 2011), pests and weeds (Li *et al.*, 2019) and pesticides (Gockowski and Ndoumbe, 2004). Accordingly, there is empirical basis to test the effect of stand type on all forms of xenobiotic use:

H4. *Mixed Cropping is positively associated with the use of pesticides*

H5. *Mixed Cropping is positively associated with the use of herbicides*

H6. *Mixed Cropping is positively associated with the use of fungicides*

3.4 Row Planting

Row planting, as opposed to broadcast planting, is one of two sowing techniques for placing seeds in the soil for germination (Aguilar and Jacobsen, 2003). Beginning with the latter, broadcast planting is performed, first of all, by preparing seedbeds and, secondly, dispersing seeds over the seedbeds by hand or using broadcast spreading equipment (Orth *et al.*, 2009). To complete the broadcast process, soil surface is swept over the seedlings to improve soil contact, fend off seed-eating pests and reduce solar radiation which preserves vitality and aids the uniform emergence of crops (Aguilar and Jacobsen, 2003). On the other hand, row planting requires the placement of seeds in furrows that are centimetres apart in defined rows which are then covered with surface soil (Fahong *et al.*, 2004). This method is often preferred for even distribution of plants which makes weeding and hoeing less of an ordeal down the line (Aguilar and Jacobsen, 2003). Selecting an ideal planting method is of genuine essence in arid regions (like Ethiopia) where water conservation and irrigation are real issues (Lalitha and Chhabra, 2020). Making appropriate decisions in this regard not only affects crop yield and quality, but also impacts on soil health (Iqbal *et al.*, 2019). Also, sowing technique has an apparent bearing on labour requirement (Vandercasteelen *et al.*, 2018), with row planting said to be more labour-intensive than broadcast planting.

Reverting to the theoretical links, during the sowing process, there is a common practice of coating seeds with pesticides and/or fungicides to control pests irrespective of broadcast or row planting (Nuyttens *et al.*, 2013). In one study, Reddy (2003) stated that row planting has the potential to reduce herbicide inputs. Yet, in another study, Gessesse (2020) wrote that since row planting allows more space for the growth of weeds compared to broadcasting, farmers are advised to apply herbicides. These claims are a clear contradiction that warrants investigation for a definitive perspective on the link between row planting and xenobiotic use. Devi *et al.* (2018) and Mahajan *et al.* (2019) offer empirical precedent of association between row planting and xenobiotic use. Althoother, these factors are considered within a package of farming innovation (Kebede *et al.*, 2017). Thus, the next hypotheses evaluate whether:

H7. *Row planting is positively associated with the use of pesticides*

H8. *Row planting is positively associated with the use of herbicides*

H9. *Row planting is positively associated with the use of fungicides*

3.5 Pesticide

Pesticides are bioactive substances that are applied in farmlands to deter, incapacitate or discourage pests while increasing crop yield, quality and economic returns (Imfeld and Vuilleumier, 2012). In order of kilograms per hectare, Japan recorded the highest annual pesticide use between 2010 and 2014, followed by China, Mexico, Brazil, Germany, France, the UK, the United States and India (Zhang, 2018). There are records dating back to 1921 on the regulation of pesticide use in Kenya (Wandiga, 2001), suggesting that this method of pest management has been practiced in Africa for more than a century. Post-independence, there have been several government programmes promoting the use of pesticides to African smallholders and commercial farmers, leading to a pesticide dependency (Williamson, 2003). For 28 years between 1966 and 1994, cotton farmers in Ivory Coast, had free [unpaid] access to pesticides because their produce was deemed to be of national importance (Ajayi *et al.*, 2009). The use of pesticides on the continent continued to grow to a point where Africa accounted for 2-4% of the \$31 billion global pesticide market (Agrow, 2006). In West Africa alone, figures from the FAO showed that pesticide imports exceeded \$800 million in 2012/2013 (Haggblade *et al.*, 2021). Just now, the global pesticides market is forecast to grow from \$75 billion in 2017 to \$90 billion by 2023 (TechSci, 2021).

In Ethiopia, common pesticides procured by small scale farmers include permethrin, cypermethrin, deltamethrin, chlorpyrifos ethyl, malathion and diazinon (Mekonen *et al.*, 2014). Theoretically, there is a plethora of papers associating pesticide use with expected harvest. Cooper and Dobson (2007, p. 1340) affirm that 'the use of pesticides has undoubtedly played a very significant role' in the spectacular increase in crop yields in the United States. Williamson *et al.* (2008) indicate the same trend in Ghana, as Bonner and Alavanja (2017, p. 89) contend that 'for the near and foreseeable future, pesticides may be an important component of a comprehensive strategy to increase crop yield'. Hence pesticide adoption rate is still seeing an upswing in Ethiopia (Negatu *et al.*, 2017), to the extent that it now poses a risk to surface water and aquatic life; following concerns raised by the Ethiopian Pesticide Risk Reduction Programme (Teklu *et al.*, 2015). Similar to Hossard *et al.*'s (2014) premise that pesticide use affects crop yield, a tenth hypothesis is formulated:

H10. *The use of pesticides is positively associated with expected harvest*

3.6 Herbicide

Herbicides, often called weed killers, are chemical substances sprayed to control unwanted plants without injuring cultivated crop (Davies and Caseley, 1999, Kebede and Anbasa, 2017). Weed management in this way began in the latter part of the 1940s, and glyphosate, first introduced in 1974, emerged as the herbicide of popular choice (Rüegg, 2007; Duke and Powles, 2008). In order of kilograms per hectare, Japan recorded the highest annual herbicide

use between 2010 and 2014, followed by Brazil, the United States, Mexico, Germany, Canada, France, the UK and India (Zhang, 2018). Worldwide, herbicides now represent 60% of all xenobiotic use in farmlands (Dayan, 2019). This is mainly because glyphosate, in particular, is said to be toxicologically and environmentally safe (Duke and Powles, 2008). Other obtainable herbicides are phenoxy hormone, triazines, amides, carbamates, dinitroanilines, urea derivatives, sulfonyl ureas, bipiridils and uracil. Zhang (2018) noted that triazines, urea derivatives, uracil and bipiridils use increased between 1990 and 2007, but the application of bipiridils has declined since 2007. In Africa, claims have been made of the obvious but overlooked 'opportunity for herbicides' (Gianessi and Ashley, 2011, p. 211), because 'smallholder farms do not use herbicides and adoption rates are less than 5%' (Gianessi, 2013, p. 1102). It seems that the vast majority of farmers manage the problem by hand weeding (Rodenburg *et al.*, 2019).

In Ethiopia, Tamru *et al.* (2017) have observed the rapid adoption of herbicides among smallholders since 2004, following a rise in the importation of glyphosate and other substances mainly from China and India. This trend is theoretically explained by herbicides being damage control agents that can minimise crop losses and maximise yield potential. Thus, Kebede and Anbasa (2017) have demonstrated the efficacy of herbicides for boosting maize yields in Ethiopia, similar to Tessema *et al.* (2018). In other crops, the effectiveness of herbicides for increasing teff yield has been cited (Tamru *et al.*, 2017), while Mitiku and Dalga (2014) reached the same conclusion for wheat. Scholars have mostly associated herbicide use with labour productivity (Gebissa *et al.*, 2019; Bouwman *et al.*, 2021). However, following the suggestion by Hailu *et al.* (2016) that, as a production input, herbicides are also an explanatory variable for crop productivity, a further hypothesis is contemplated:

H11. The use of herbicides is positively associated with expected harvest

3.7 Fungicide

Fungicides are chemical compounds diffused for the control of parasitic fungi or their pathogens during crop cultivation (Oliver and Hewitt, 2014). The history of fungicide use predates 1755 when it was discovered that a seed-borne fungi [*Tilletia Laevis*] in wheat could be controlled by treating seeds with lime and salt (Morton and Staub, 2008). Fungicides have since become more sophisticated in their composition and include common names such as Benomyl, Iprodione, Lime Sulfur, Mancozeb and Propineb (Kang and Jung, 2017). In order of kilograms per hectare, between 2010 and 2014, the highest annual fungicide use by country was in Japan followed by Mexico, France, the UK, Germany, Brazil, the United States and India (Zhang, 2018). Zhang (2018) adds that the use of fungicides increased from 1990 to 2007, but has oscillated ever since. This fluctuation could, in part, be explained by the increasing resistance of pathogens to fungicides for which 'new anti-fungal compounds need

to be discovered' (Steinberg and Gurr, 2020, p. 1). Notwithstanding, the global fungicide market was worth approximately \$13.4 billion in 2018 (Çıldır and Liman, 2020). Where they are used, fungicides are deemed to be effective and low in toxicity (Zhang *et al.*, 2018). Although fungicide use is generally low in Africa (Ochieng *et al.*, 2019), they are sprayed in potato plantations in South Africa (Muzhinji *et al.*, 2018) and Uganda (Namuga *et al.*, 2017). They are also mainly used by vegetable farmers in Benin, Ethiopia, Ghana and Senegal (Williamson *et al.*, 2008).

One of the main attractions to fungicides in Ethiopia is the mitigation of wheat rust fungus which, according to Allen-Sader *et al.* (2019), poses one of the greatest threats to global food security. Ethiopia harvests 1.7 million hectares of wheat annually and this produce is relied on by 4.2 million households (Central Statistics Agency, 2018). Yet in the 2013-2014 period, an estimated 40,000 hectares were diseased by wheat rust (Olivera *et al.*, 2015). To offset such outcomes, fungicides are also leveraged in the production of garlic (Endalew *et al.*, 2020) and sugarcane in Ethiopia (Kecha, 2020). Relatedly, Kassaw *et al.* (2021) indicate that greater yields accrue from the use of fungicide in Ethiopian potato plantations. In fact, there is no shortage of theoretical links between fungicide use and improved farm harvests (Danso-Abbeam and Baiyegunhi, 2017). Inspired by previous tests probing how fungicides correlate with crop yield (Haverkort and Bicomumpaka, 1986; Dalla Lana *et al.*, 2015), the penultimate hypothesis checks the extent to which:

H12. The use of fungicides is positively associated with expected harvest

3.8 Expected Harvest, Improved seeds, Mixed cropping and Row Planting

There are extensive claims that improved seeds (e.g. Lakshman, 1993), mixed cropping (e.g. Yigezu *et al.*, 2019) and row planting (e.g. Vandercasteelen *et al.*, 2018) separately enhance farm productivity. As a proxy for productivity, expected harvest is depicted here as the estimated units of produce predicted by households for forthcoming harvest. The inclusion and review of this factor harps on the expectancy-value theory of motivation. First conceived by Atkinson (1957), the expectancy-value perspective postulates that the degree of success envisaged by individuals and their inner beliefs will occasion subsequent behaviour (Wigfield, 1994). The theory also helps to explain individuals' level of persistence during the performance of tasks (Eccles and Wigfield, 2020). There is an argument that because of the relatively long lead time between planting and harvest seasons, households pre-plan their land management practices in order to maximise expected harvest. In other words, the expected value of crop yield will instigate the performance of specific land management practices from the outset. Thus, McMaster *et al.* (2000) pondered the relationship between expected harvest resulting from soil and water conservation. Also, recalling that expectancy-value also encompasses inner beliefs, Dennis (1989) considered individual characteristics as a function of expected

timber harvest. In the same spirit, drawing on the farming innovation techniques appraised in this study, the last set of hypotheses test the extent to which:

H13. *Improved Seeds are directly and positively associated with expected harvest*

H14. *Mixed Cropping is directly and positively associated with expected harvest*

H15. *Row Planting is directly and positively associated with expected harvest*

4.0 Method

4.1 Data and Measures

The data examined in this study were collected from the 2018-2019 living standards measurement study of Ethiopian households conducted by the World Bank (2021c) using stratified random sampling. This follows Peet *et al.* (2015), Aregbeshola and Khan (2018) and Gebremariam and Tesfaye's (2018) precedent. The scope of the data covered household characteristics, community infrastructure, livestock information, post planting and post harvest information. Out of 16,914 cases in the sample, only 7,625 households that indicated the use of all three xenobiotics were extrapolated. For measurement, seven main variables were examined namely improved seeds (IMPSEED), mixed cropping (MIXCROP), row planting (ROWPLAN), use of pesticides (PEST), use of herbicides (HERB), use of fungicides (FUNG) and expected harvest (EXPHARV). The control variables were type of farm (FARMTYP), name of town (REGION) and type of crop (CROP). All factors were measured using single item questions as outlined in table 1 below.

Table 1: Measurement Details

Variable	Items	Scales
IMPSEED	What type of seed/seedling was used for crop on the field?	Traditional/Improved
MIXCROP	Was the area planted with crop on the field pure stand or mixed?	Pure Stand/Mixed Stand
ROWPLAN	What type of crop sowing techniques was used for crop on the field?	Broadcast Planting/Row Planting
PEST	Did you use any pesticide to prevent damage of crop on this field?	Yes/No
HERB	Did you use any herbicide to prevent damage of crop on this field?	Yes/No
FUNG	Did you use any fungicide to prevent damage of crop on this field?	Yes/No
EXPHARV	How much of crop [units] do you expect to harvest from this field?	Continuous
FARMTYP	What is the holder's farm type?	Crop Production Only/Both Livestock and Crop Production
REGION	Name of Town	Continuous
CROP	Type of Crop	Continuous

4.3 Sample Characteristics

Of the 7,625 households in the sample, 89% were both crop and livestock smallholders while 11% cultivated only crops. Geographically, 29% of the households resided in the northern Amhara area followed by 20.8% in the Southern Nations, Nationalities and People's (SNNP) region. Other territories were Oromia spanning from the mid-west to the south of the country

(15.9%), Benishangul-Gumuz in the north-west (12.7%) and Tigray in the northernmost part (10.7%). The remaining territories each represented less than 4% of the sample but nine out of the Ethiopia's twelve regions and chartered cities have been surveyed. The crops cultivated were mostly maize in 19.2% of households, followed by sorghum and teff in 12.6% and 11.5% respectively. Table 2 presents a full list and proportion of other produce including wheat, barley and kale.

Table 2: Sample Characteristics

Farm Type		
	Frequency	Percent
Both Livestock and Crop production	6790	89.0
Crop Production Only	835	11.0
Total	7625	100.0
Region		
	Frequency	Percent
Amhara	2213	29.0
SNNP	1586	20.8
Oromia	1214	15.9
Benishangul-Gumuz	972	12.7
Tigray	817	10.7
Dire Dawa	282	3.7
Harar	279	3.7
Gambela	235	3.1
Afar	27	0.4
Total	7625	100.0
Crop		
	Frequency	Percent
Maize	1464	19.2
Sorghum	957	12.6
Teff	936	12.3
Wheat	540	7.1
Barley	476	6.2
Kale	276	3.6
Horse Beans	271	3.6
Red Kidney Beans	224	2.9
Godere	211	2.8
Red Pepper	215	2.8
Millet	196	2.6
Sweet Potato	167	2.2
Garlic	138	1.8
Groundnuts	123	1.6
Pumpkins	101	1.3
Potatoes	102	1.3
Soya Beans	99	1.3
Field Peas	99	1.3
Other Root Crops	77	1.0
Sesame	67	0.9
Lentils	69	0.9
Rice	65	0.9
Vetch	72	0.9
Chickpeas	70	0.9
Green Pepper	58	0.8
Rapeseed	63	0.8
Nueg	63	0.8
Linseed	52	0.7

Onion	51	0.7
Cassava	52	0.7
Ginger	29	0.4
Mung Bean/Masho	30	0.4
Other Vegetables	20	0.3
Sunflower	26	0.3
Cabbage	22	0.3
Spinach	19	0.2
Tomatoes	12	0.2
Beer Root	18	0.2
Carrot	19	0.2
Oats	12	0.2
Other spices	10	0.1
Cardamom	8	0.1
Coriander	5	0.1
Tobacco	9	0.1
Other Pulses	5	0.1
Lettuce	5	0.1
Turmeric	9	0.1
Other Cereal	4	0.1
Cotton	2	0.0
Cauliflower	3	0.0
White Cumin	2	0.0
Gibto	2	0.0
Total	7625	100.0

5.0 Analysis

A non-linear regression-based partial least squares structural equation modelling [PLS-SEM] technique was adopted to analyse the household data. The specific software and version used was WarpPLS 7.0 (Kock, 2019). This approach was deemed appropriate because of its greater predictive power over covariance-based structural equation modelling [CB-SEM]. To explain, ‘in a direct comparison with CB-SEM, the variance explained in the dependent variable is substantially higher [in PLS-SEM]’ (Hair *et al.*, 2017, p. 119). WarpPLS is also suitable for analysing constructs with binary data (Sajid *et al.*, 2020).

5.1. Measurement Model

Preparatory to path analysis, it is important to assess the reliability and validity of variables. However, all variables in the current study are single item constructs for which reliability and validity tests do not suffice. In terms of collinearity, variance inflation factor [VIF] scores were generated for all variables to confirm the absence of linear combinations in the structural model. As demonstrated in table 3, all VIF values were lower than the 5 limit recommended by Hair *et al.* (2011).

Table 3: Collinearity Diagnostic

	<i>IMPSEED</i>	<i>MIXCROP</i>	<i>ROWPLAN</i>	<i>PEST</i>	<i>HERB</i>	<i>FUNG</i>	<i>EXPHARV</i>	<i>REGION</i>	<i>CROP</i>	<i>FARMTYP</i>
VIF	1.120	1.152	1.231	1.080	1.129	1.105	1.068	1.194	1.086	1.006

5.2 Structural Model and Hypothesis Testing

The structural model is examined through the path coefficients (β) and p -values of the associations hypothesised. Figure 1 shows the obtained results.




Figure 1. The Structural Model

From figure 1, beginning with land management practices, the path analysis of improved seeds indicates that the factor positively correlates with the use of pesticides ($\beta = 0.18$), herbicides ($\beta = 0.07$) and fungicides ($\beta = 0.20$). However, mixed stands have a negative association with pesticide ($\beta = -0.04$), herbicide ($\beta = -0.19$) and fungicide ($\beta = -0.06$) use. There are mixed findings in how row planting affects household's use of xenobiotics. It [row planting] has a weak but positive influence on the use of pesticides ($\beta = 0.02$) but a negative impact on herbicide ($\beta = -0.13$) and fungicide ($\beta = -0.04$). In turn, all types of xenobiotic use reduce expected harvest [pesticides ($\beta = -0.00$), herbicides ($\beta = -0.10$) and fungicides ($\beta = -0.04$)]. Furthermore, the path analysis revealed that both broadcast planting ($\beta = -0.15$) and improved seeds ($\beta = -0.06$) decrease expected harvest, while mixed stands ($\beta = 0.06$) moderately increases the outcome. All control variables, region ($\beta = -0.06$), crop ($\beta = -0.07$) and type of farm ($\beta = -0.00$) are negatively correlated to expected harvest. Overall, the model explains 6% of households' expected harvest in rural Ethiopia.

Table 4 below further summarises the hypotheses testing and results.

Table 4: Hypothesis Testing

Hypothesised Relationships	Path Coefficients	P-Values	Test
H1. IMPROVED SEEDS \Rightarrow PESTICIDES	0.18	<.01	Significant

Hypothesised Relationships	Path Coefficients	P-Values	Test
H2. IMPROVED SEEDS ⇒ HERBICIDES	0.07	<.01	Significant
H3. IMPROVED SEEDS ⇒ FUNGICIDES	0.20	<.01	Significant
H4. MIXED CROPPING ⇒ PESTICIDES	-0.04	<.01	Not Significant
H5. MIXED CROPPING ⇒ HERBICIDES	-0.19	<.01	Not Significant
H6. MIXED CROPPING ⇒ FUNGICIDES	-0.06	<.01	Not Significant
H7. ROW PLANTING ⇒ PESTICIDES	0.02	0.02	Significant
H8. ROW PLANTING ⇒ HERBICIDES	-0.13	<.01	Not Significant
H9. ROW PLANTING ⇒ FUNGICIDES	-0.04	<.01	Not Significant
H10. PESTICIDE ⇒ EXPECTED HARVEST	-0.00	0.35	Not Significant
H11. HERBICIDE ⇒ EXPECTED HARVEST	-0.10	<.01	Not Significant
H12. FUNGICIDE ⇒ EXPECTED HARVEST	-0.04	<.01	Not Significant
H13. IMPROVED SEEDS ⇒ EXPECTED HARVEST	-0.06	<.01	Not Significant
H14. MIXED CROPPING ⇒ EXPECTED HARVEST	0.06	<.01	Significant
H15. ROW PLANTING ⇒ EXPECTED HARVEST	-0.15	<.01	Not Significant

6.0 Discussion

By way of interpretation, this study demonstrates that the higher the use of pesticides, herbicides and fungicides, the lower households' expected harvest will be. Of the six land management practices examined, households' best chance of increasing expected harvest is by tending mixed crops. To be sure, (1) cultivating more than one crop moderately increases expected harvest, and (2) sowing improved seeds using row planting will decrease expected harvest. It has also been shown that using improved seeds increases households' tendency to apply pesticides, herbicides and fungicides in their smallholdings. Likewise, row planting very slightly increases households' use of pesticides but reduces the application of herbicides and fungicides. Based on these findings, Lakshman's (1993, p. 225) claim that improved seeds 'dramatically increase agricultural output' is disproved in the context of rural households in Ethiopia. Also, Daellenbach *et al.*'s (2005) assertion that mixed cropping systems reduce total production is challenged, while Yigezu *et al.*'s (2019) stipulation that mixed cropping provides higher yields is somewhat upheld. Furthermore, Reddy's (2003) argument that row planting reduces herbicide use is sustained, negating Gessesse's (2020) recommendation that farmers practicing row planting should apply herbicides. Instead of increasing crop yield as claimed by Cooper and Dobson (2007), Williamson *et al.* (2008) and Bonner and Alavanja (2017), this study finds that spraying pesticides reduces crop yield. Similarly, the view that herbicides increase yield in maize (Kebede and Anbasa, 2017; Tessema *et al.*, 2018), teff (Tamru *et al.*, 2017) and wheat (Mitiku and Dalga, 2014) does not seem to be true when the expected harvest of all crops is considered. Kassaw's (2021) finding that fungicides increase yields is also contested by the current findings. It is now opportune to reflect on these findings through the contingency and sustainability lens.

By and large, Luthans and Stewart (1977) believe that pressing environmental, resource and management influences are at the core of how economic agents operate. Beginning with

environmental factors beyond households' control, in Ethiopia, agricultural extension programmes compel the use of improved seeds and other inputs as a condition for government support. Unequivocally, Bachewe *et al.* (2018, p. 288) affirm that 'in the context of Ethiopia, agricultural input supply strategies in the last decade encouraged farmers to adopt chemical fertilizers and improved seeds as a package, at times bundled with input credit, making adoption of these two inputs an inherently simultaneous decision or one between sets of possible technology bundles'. There is also evidence that loyalty to the defunct political party, the Ethiopian People's Revolutionary Democratic Front (EPRDF), has been a condition for farmers receiving seeds, credit and other inputs (Berhanu and Poulton, 2014). Characteristically, cooperative societies are organised to work with extension programs to cascade government support to grassroot farmers. Consequently, Abebaw and Haile's (2013, p. 87) indication that 'membership of agricultural cooperative [in Ethiopia] has a significant positive and larger impact on pesticide adoption' as well as improved seeds is hardly surprising.

In terms of the influence of resources or the lack of in rural households, there is immense poverty among this social group (Demissie and Kasie, 2017). A large proportion of farming families in Ethiopia are barely able to supply the labour required for subsistence activity (Vandercasteelen *et al.*, 2018). Being in a dire situation, these households are heavily reliant on external support for inputs such as seeds and other inputs (Abebaw and Haile, 2013). Invariably, the seeds on offer are of the improved variety which, as this study demonstrates, increases the likelihood of pesticide, herbicide and fungicide use. In turn, these practices reduce expected harvest, further constraining the resources available to rural households.

The above environmental and resource pressures converge to determine the quality of farm management decisions made by rural households. In the face of environmental and resource limitations, they may be obliged to obtain improved seeds and use different kinds of xenobiotics without regard to expected harvest because the alternative to this could be a lack of subsistence activity. By contrast, it can be expected that rural households that are resistant to the overtures of extension programmes and non-reliant on farming inputs from external sources will exhibit greater expected harvest. Therefore, self-sufficiency or the delivery of 'technical packages' with no strings attached is deemed to be an important criterion for raising rural households' productivity.

Revisiting Elkington's (1994) sustainability theory, consideration of Slaper and Hall's (2011) economic, environmental and social dimensions is appropriate here. In relation to economic measures, this study maintains that the land management practices examined do not yield

sufficient output nor do they optimise the labour of rural households. Take row planting for example, this sowing technique requires greater effort from households yet it reduces expected harvest significantly more than other farming innovations. This result concurs with Zhai *et al.*'s (2018) verdict that broadcasting does indeed increase grain yield. It is also worth acknowledging evidence that pesticide use correlates more positively with the cultivation of cash crops for commercial purposes (Matthews *et al.*, 2003), rather than for domestic consumption. Household farmers in Ethiopia also deem herbicides to be ineffective (Gessesse, 2020). According to Mariyono *et al.* (2018), xenobiotics are only needed if and when pest and disease infestations occur otherwise they are mostly a wasteful insurance policy. Overall, the six land management practices inherent in this review are suboptimal and do not make economic sense for rural households in Ethiopia.

Taking the environment into account, the ecological hazards posed by pesticides, herbicides and fungicides have been widely reported in different countries (De la Cruz *et al.*, 2014; Tamru *et al.*, 2017; Kapsi *et al.*, 2019). In Africa, there are serious shortcomings in the handling practices of these toxic substances (Williamson *et al.*, 2008). Even when used proportionately, the leaching of xenobiotics into the soil and groundwater causes long-term pollution (Arias-Estévez *et al.*, 2008). Above ground, xenobiotics also disrupt biodiversity and much needed ecological balance. For example, Ethiopian farmers have been alleged to handle pesticides very indiscriminately without consideration of the effect on honeybees, threatening the essential pollination cycle (Fikadu, 2020). In fact, over 90% of Ethiopian farmers are unaware that pesticides could damage water bodies (Mengistie *et al.*, 2017). Through oblivion, land quality is compromised by the toxic content of xenobiotics such as sulphur in herbicides.

To conclude, to be sustainable, land management practices and xenobiotic use ought to improve rural households' quality of life but this does not appear to be the case. For instance, there are doubts surrounding the positive effect of fungicides on the nutritional composition of crops (Mbah *et al.*, 2018). Worse still, Banjo *et al.* (2010) report that 25 million land workers in developing countries suffer from pesticide poisoning every year. This leads to diagnoses such as respiratory difficulty, memory loss, various cancers, skin conditions and other acute diseases that cause 18,000 annual deaths (Banjo *et al.*, 2010). These statistics are inconsistent with the social health and life expectancy principles of sustainability. Land management practices and xenobiotic use are only beneficial if they have, as Green (2012, p. 1324) best puts it, 'the ability to bring benefits to growers and satisfy society's expectations for safe, abundant and affordable food...in an environmentally sustainable way'.

7.0 Implications, Future Research and Limitations

7.1 Theoretical Contributions

This paper is one of the first to conceptualise multiple land management practices vis-à-vis expected harvest for empirical validation. Specifically, it breaks new ground by linking row planting to the use of pesticides, herbicides and fungicides. Its core contribution to the literature is the isolation of specific practices that will increase expected harvest in rural Ethiopia and, possibly, similar contexts. It reconciles contradictory findings in the farming innovation and productivity nexus and reduces theoretical ambiguity by offering definitive perspectives in two areas:

First, it has been determined that the ingredients for increasing expected harvest are, in order of coefficient strength, mixed cropping, use of traditional seeds and broadcast planting. Moreover, the use of traditional seeds will reduce reliance on xenobiotics, which also bodes well for maximising harvests. Accordingly, supported by expectancy-value theory, the evidence of mixed cropping, traditional seeds and broadcast planting as viable predictors of expected harvest introduces fresh antecedents to the farming innovation and productivity nexus. Second, by isolating and separately examining pesticides, herbicides and fungicides, we offer exactitude and theoretical specificity on the distinctive effect of different xenobiotics. This is an improvement on previous studies that assess herbicides and fungicides as a composite construct of pesticides. Undoubtedly, the use of composite constructs when multi dimensions can be interrogated leads to the underestimation of correlations (Bracken, 1996; Craven *et al.*, 2003; O'Mara *et al.*, 2006). The current structural model and the ensuing findings effectively address this problem.

7.2 Practical Implications

For rural households, the chief contribution of this inquiry is that, from the outset, (1) they are better off sowing traditional seeds using broadcasting method. Subsequently, (2) the tending of crops sprouting from traditional seeds will reduce the inclination to use pesticides, herbicides and fungicides. This insight should incentivise households to spurt xenobiotics more consciously and less habitually. This point is especially pertinent as Genet *et al.* (2020) identify pesticides as one significant variable costs incurred by Ethiopian farmers. The findings also refute the wisdom of government support programmes encouraging rural households to sow improved seeds and dispense xenobiotics. On the current evidence, these factors reduce households' expected harvest. In fact, their increased use in Ethiopia and Africa as a whole may partly explain why the continent's agricultural output lags behind global production. Therefore, attention is also drawn to foreign donors of agricultural inputs in Africa including the Bill and Melinda Gates Foundation, the International Fund for Agricultural Development, the World Health Organisation and the United Nations Population Fund (Shaw and Wilson,

2020). These entities can reconsider their benevolence by sourcing and gifting traditional seeds to rural households, at the same time as promoting awareness on the hazards of xenobiotics and the potential benefits of broadcast planting. In addition to surface water and aquatic life, the Ethiopian Pesticide Risk Reduction Programme should also consider rural households' expected harvest to be an area of concern.

7.3 Limitations and Future Research

There are a few shortcomings in this study that prompt follow-up research. First, the parallels drawn with contingency and sustainability theory are only reflective and conceptual. Nevertheless, this paves way for new studies to empirically measure and report contingent and sustainable behaviour. Second, the structural model did not check the type of xenobiotics used and how their toxological properties affect expected crop harvest not just in terms of quantity but also in quality. In the interest of empirical specificity, future studies could be predicated on this thesis as such results will deepen insights on the exact effects of xenobiotic use. Third, in new studies, researchers may also examine actual harvest rather than expected harvest as the dependent variable. Fourth, although the findings may also be useful in countries neighbouring Ethiopia in East Africa, other studies are invited to adopt the path diagram and possibly validate the current results for generalisability. Finally, in the grand scheme, the current findings hint at the benefits of organic farming without the need for improved seeds and xenobiotics. Future research can conceptualise organic farming practices in the farming innovation and productivity nexus to advance knowledge in this area.

ACKNOWLEDGMENT

We thank Mohamed Yacine Haddoud for his useful comments in the conceptualisation of this study. We also thank the anonymous reviewers for highlighting areas to improve the robustness and presentation.

DISCLAIMER

The authors acknowledge that the original collector of the data, the authorised distributor of the data, and the relevant funding agency bear no responsibility for use of the data or for interpretations or inferences based upon such uses.

References

- Abate, G., Dereje, M., Hirvonen, K. and Minten, B. (2020), 'Geography of public service delivery in rural Ethiopia', *World Development*, Vol. 136, p.105133.
<https://doi.org/10.1016/j.worlddev.2020.105133>.
- Abay, K., Blalock, G. and Berhane, G. (2017), 'Locus of control and technology adoption in developing country agriculture: Evidence from Ethiopia', *Journal of Economic Behavior and Organization*, Vol. 143 No. C, pp. 98-115.
- Abebaw, D. and Haile, M. (2013), 'The impact of cooperatives on agricultural technology adoption: Empirical evidence from Ethiopia', *Food Policy*, Vol. 38, pp. 82-91.
<https://doi.org/10.1016/j.foodpol.2012.10.003>.

- Abebe, G. and Alemu, A. (2017), 'Role of Improved Seeds Towards Improving Livelihood and Food Security at Ethiopia', *International Journal of Research – Granthaalayah*, Vol. 5 No. 2, pp. 338-356.
- Agrow (2006) *World Agchem Market Steady*, Agrow, Ohio.
- Aguilar, P. and Jacobsen, S. (2003), 'Cultivation of Quinoa on the Peruvian Altiplano', *Food Reviews International*, Vol. 19 No. 1-2, pp. 31-41.
- Ahmed, M., Geleta, K., Tazeze, A., Mesfin, H. and Tilahun, E. (2017), 'Cropping systems diversification, improved seed, manure and inorganic fertilizer adoption by maize producers of eastern Ethiopia', *Journal of Economic Structures*, Vol. 6 No. 1, pp. 1-16.
- Ajayi, O., Akinnifesi, F., Sileshi, G. and Ajayi, A. (2009), 'Agricultural Policies and the Emergence of Cotton as the Dominant Crop in Northern Côte d'Ivoire: Historical Overview and Current Outlook', *Natural Resources Forum*, Vol. 33 No. 2, pp. 111-122.
- Allen-Sader, C., Thurston, W., Meyer, M., Nure, E., Bacha, N., Alemayehu, Y., Stutt, R., Safka, D., Craig, A., Derso, E. and Burgin, L. (2019), 'An early warning system to predict and mitigate wheat rust diseases in Ethiopia', *Environmental Research Letters*, Vol. 14 No. 11, pp. 1-11.
- Aregbeshola, B. and Khan, S. (2018), 'Out-of-pocket payments, catastrophic health expenditure and poverty among households in Nigeria 2010', *International Journal of Health Policy and Management*, Vol. 7 No. 9, pp. 798-806.
- Arias-Estévez, M., López-Periágo, E., Martínez-Carballo, E., Simal-Gándara, J., Mejuto, J.C. and García-Río, L., (2008), 'The mobility and degradation of pesticides in soils and the pollution of groundwater resources', *Agriculture, Ecosystems & Environment*, Vol. 123 No. 4, pp. 247-260.
- Armanda, D., Guinée, J. and Tukker, A. (2019), 'The second green revolution: Innovative urban agriculture's contribution to food security and sustainability - A review', *Global Food Security*, Vol. 22, pp. 13-24. <https://doi.org/10.1016/j.gfs.2019.08.002>.
- Arthur, W. (1994), 'Bounded rationality and inductive behavior', *American Economic Review*, Vol. 84 No. 2, pp. 406-411.
- Atkinson, J. (1957), 'Motivational Determinants of Risk-taking Behavior', *Psychological Review*, Vol. 64 No. 6, pp. 359-372.
- Ayalew, M., Dakhli, M. and Lafhaj, Z. (2016), 'The future of lean construction in Ethiopian construction industry', *International Journal of Engineering Research & Technology*, Vol. 5 No. 2, pp. 107-113.
- Bachewe, F., Berhane, G., Minten, B. and Taffesse, A. (2018), 'Agricultural Transformation in Africa? Assessing the Evidence in Ethiopia', *World Development*, Vol. 105 No. C, pp. 286-298.
- Baiphethi, M. and Jacobs, P. (2009), 'The contribution of subsistence farming to food security in South Africa'. *Agrekon*, Vol. 48 No. 4, pp. 459-482.
- Banjo, A., Aina, S. and Rije, O. (2010), 'Farmers' knowledge and perception towards herbicides and pesticides usage in Fadama area of Okun-Owa, Ogun State of Nigeria', *African Journal of Basic and Applied Sciences*, Vol. 2 No. 5-6, pp. 188-194.
- Berhane, G., Ragasa, C., Abate, G. and Assefa, T. (2018), *The State of Agricultural Extension Services in Ethiopia and their Contribution to Agricultural Productivity*, International Food Policy Research Institute, Addis Ababa.
- Berhanu, K. and Poulton, C. (2014), 'The political economy of agricultural extension policy in Ethiopia: economic growth and political control', *Development Policy Review*, Vol. 32 No. 2, pp. 197-213.
- Bjornlund, V., Bjornlund, H. and van Rooyen, A. (2020), 'Why agricultural production in sub-Saharan Africa remains low compared to the rest of the world – a historical perspective', *International Journal of Water Resources Development*, Vol. 36 No. 1, pp. 20-53.
- Bonner, M. and Alavanja, M. (2017), 'Pesticides, human health, and food security', *Food and Energy Security*, Vol. 6 No. 3, pp. 89-93.
- Bouwman, T., Andersson, J. and Giller, K. (2021), 'Herbicide induced hunger? Conservation agriculture, Ganyu labour and rural poverty in Central Malawi', *The Journal of Development Studies*, Vol. 57 No. 2, pp. 244-263.

- Bracken, B. (1996), 'Clinical applications of a context-dependent, multidimensional model of self-concept', in Bracken, E. (Ed.) *Handbook of Self-concept*, John Wiley, New York, pp. 463-503.
- Central Statistics Agency (2018), *Agricultural Sample Survey 2016/2017: Report on Area and Production of Major Crops*, Central Statistics Agency, Addis Ababa.
- Christiaensen, L., Demery, L. and Paternostro, S. (2003), 'Macro and micro perspectives of growth and poverty in Africa', *The World Bank Economic Review*, Vol. 17 No. 3, pp. 317-347.
- Cieslik, K. and D'Aoust, O. (2018), 'Risky business? Rural entrepreneurship in subsistence markets: evidence from Burundi', *The European Journal of Development Research*, Vol. 30 No. 4, pp. 693-717.
- Çıldır, D. and Liman, R. (2020), 'Cytogenetic and genotoxic assessment in *Allium cepa* exposed to imazalil fungicide', *Environmental Science and Pollution Research*, Vol. 27 No. 16, pp. 20335-20343.
- Clapham, C. (2018), 'The Ethiopian Developmental State', *Third World Quarterly*, Vol. 39 No. 6, pp. 1151-1165.
- Cook, R. and Weller, D. (2004), 'In defense of crop monoculture: new directions for a diverse planet', Proceedings of the 4th International Crop Science Congress, 26 Sep–1 Oct 2004, Brisbane, Australia. Published on CDROM. Web site [www.cropscience.org.au](http://www.cropsscience.org.au) [Accessed 21st March 2021].
- Cooper, J. and Dobson, H. (2007), 'The benefits of pesticides to mankind and the environment', *Crop Protection*, Vol. 26 No. 9, pp. 1337-1348.
- Craven, R., Marsh, H. and Burnett, P. (2003), 'Cracking the self-concept enhancement conundrum: A call and blueprint for the next generation of self-concept enhancement research', in Marsh, H., Craven, R. and McInerney, D. (Eds.) *International Advances in Self Research*, Information Age, Connecticut, pp. 67-90.
- Daellenbach, G., Kerridge, P., Wolfe, M., Frossard, E. and Finckh, M. (2005), 'Plant productivity in cassava-based mixed cropping systems in Colombian hillside farms', *Agriculture, Ecosystems & Environment*, Vol. 105 No. 4, pp. 595-614.
- Dalla Lana, F., Ziegelmann, P., Maia, A., Godoy, C. and Del Ponte, E. (2015), 'Meta-analysis of the relationship between crop yield and soybean rust severity', *Phytopathology*, Vol. 105 No. 3, pp. 307-315.
- Damania, R., Berg, C., Russ, J., Federico Barra, A., Nash, J. and Ali, R. (2017), 'Agricultural technology choice and transport', *American Journal of Agricultural Economics*, Vol. 99 No. 1, pp. 265-284.
- Danso-Abbeam, G. and Baiyegunhi, L. (2017), 'Adoption of agrochemical management practices among smallholder cocoa farmers in Ghana', *African Journal of Science, Technology, Innovation and Development*, Vol. 9 No. 6, pp. 717-728.
- Davies, J. and Caseley, J. (1999), 'Herbicide Safeners: A Review', *Pesticide Science*, Vol. 55 No. 11, pp. 1043-1058.
- Dawud, S., Dugda, D. and Giza, B. (2019), 'Assessment of pesticides use and its effects on honeybee colonies in West and East Hararghe Zones of Oromia region', *Regional Review Workshop on Completed Research Activities of Livestock Research Directorate*, Adami Tulu Agricultural Research Center, Adami Tulu 07-12 October.
- Dayan, F. (2019), 'Current status and future prospects in herbicide discovery', *Plants*, Vol. 8 No. 9, pp. 341-358.
- De la Cruz, E., Bravo-Durán, V., Ramírez, F. and Castillo, L. (2014), 'Environmental hazards associated with pesticide import into Costa Rica, 1977-2009', *Journal of Environmental Biology*, Vol. 35 No. 1, pp. 43-55.
- Deichmann, U., Shilpi, F. and Vakis, R. (2009), 'Urban proximity, agricultural potential and rural non-farm employment: Evidence from Bangladesh', *World Development*, pp. 37 Vol. 3, pp. 645-660.
- Demiessie, H. (2021), *Ethiopia: From Revolutionary Democracy to Prosperity: The Moral and Philosophical Foundations of a Prosperous Society*, Independent Publisher, Jijjiga.
- Demiessie, B. and Kasie, T. (2017), 'Rural households' vulnerability to poverty in Ethiopia', *Journal of Poverty*, Vol. 21 No. 6, pp. 528-542.

- Demlew, M., Alemu, B. and Awuk, A. (2019), 'Nutritive value Evaluation of Buffel grass and Silver leaf Desmodium Grown in Pure Stands and in Mixture at Different Harvesting Times in Gozamen District, East Gojjam Zone, Ethiopia', *Greener Journal of Agricultural Sciences*, Vol. 9 No. 3, pp. 315-321.
- Dennis, D. (1989), 'An economic analysis of harvest behavior: integrating forest and ownership characteristics', *Forest Science* Vol. 35 No. 4, pp. 1088-1104.
- Devi, S., Hooda, V., Kamboj, N. and Singh, J. (2018), 'Bioefficacy of herbicides in relation to planting techniques in wheat', *Chemical Science Review and Letters*, Vol. 7 No. 25, pp. 77-82.
- Dorfman, J. (1996), 'Modeling multiple adoption decisions in a joint framework', *American Journal of Agricultural Economics*, Vol. 78 No. 3, pp. 547-557.
- Dubey, K. and Fulekar, M. (2011), 'Effect of pesticides on the seed germination of *Cenchrus setigerus* and *Pennisetum pedicellatum* as monocropping and co-cropping system: implications for rhizospheric bioremediation', *Romanian Biotechnological Letters*, Vol. 16 No. 1, pp. 5908-5918.
- Duke, S. and Powles, S. (2008), 'Glyphosate: a once-in-a-century herbicide', *Pest Management Science*, Vol. 64 No. 4, pp. 319-325.
- Eccles, J. and Wigfield, A. (2020), 'From expectancy-value theory to situated expectancy-value theory: A developmental, social cognitive, and sociocultural perspective on motivation', *Contemporary Educational Psychology*, Vol. 61, pp.101859. <https://doi.org/10.1016/j.cedpsych.2020.101859>
- Elkington, J. (1994), 'Towards the sustainable corporation: Win-win-win business strategies for sustainable development', *California Management Review*, Vol. 36 No. 2, pp. 90-100.
- Emerton, L. and Snyder, K. (2018), 'Rethinking sustainable land management planning: Understanding the social and economic drivers of farmer decision-making in Africa', *Land Use Policy*, Vol. 79 No. C, pp. 684-694.
- Endalew, Z., Terefe, H. and Dejene, M. (2020), 'Management of Rust (*Puccinia allii* R.) through Host Resistance and Alternate Application of Fungicides on Garlic (*Allium sativum* L.) in North Wollo, Ethiopia', *East African Journal of Sciences*, Vol. 14 No. 1, pp. 65-82.
- Ezulike, T. and Igwatu, R. (1993), 'Effects of intercropping cassava and pigeon pea on green spider mite *Mononychellus tanajoa* (Bondar) infestation and on yields of the associated crops', *Discovery and Innovation*, Vol. 5 No. 4, pp. 355-359.
- Fahong, W., Xuqing, W. and Sayre, K. (2004), 'Comparison of conventional, flood irrigated, flat planting with furrow irrigated, raised bed planting for winter wheat in China', *Field Crops Research*, Vol. 87 No. 1, pp. 35-42.
- FAO (2009a), *Technical papers from the Expert Meeting on How to Feed the World in 2050*, Food and Agriculture Organisation of the United Nations, Rome.
- FAO (2009b), *Responding to the challenges of a changing world: The role of new plant varieties and high quality seeds in agriculture*, Food and Agriculture Organisation of the United Nations, Rome.
- FAO (2015), *Socio-economic context and role of Agriculture*, Food and Agriculture Organisation of the United Nations, Rome.
- FAO (2021a), *Family Farming in Africa*, Available online: <http://www.fao.org/family-farming/regions/africa/en/> [Accessed 7th March 2021].
- FAO (2021b), *Plan Pest and Diseases*, Available online: [http://www.fao.org/emergencies/emergency-types/plant-pests-and-diseases/en/?page=4&ipp=10&tx_dynalist_pi1\[par\]=YToxOntzOjE6lkwiO3M6MToiMCI7fQ](http://www.fao.org/emergencies/emergency-types/plant-pests-and-diseases/en/?page=4&ipp=10&tx_dynalist_pi1[par]=YToxOntzOjE6lkwiO3M6MToiMCI7fQ) [Accessed 7th March 2021].
- Feder, G. (1982), 'Adoption of interrelated agricultural innovations: Complementarity and the impacts of risk, scale, and credit', *American Journal of Agricultural Economics*, Vol. 64 No.1, pp. 94-101.
- Fikadu, Z. (2020), 'Pesticides use, practice and its effects on honeybee in Ethiopia: A review. *International Journal of Tropical Insect Science*', Vol. 40, pp. 473-481. <http://creativecommons.org/licenses/by/4.0/>.

- Fondong, V., Thresh, J. and Zok, S. (2002), 'Spatial and temporal spread of cassava mosaic virus disease in cassava grown alone and when intercropped with maize and/or cowpea', *Journal of Phytopathology*, Vol. 150 No. 7, pp. 365-374.
- Gebissa, B., Shumeta, Z. and Regassa, G. (2019), 'Gender Difference and Its Effect on Agricultural Productivity: The Case of Yubdo District in Ethiopia', *World Journal of Agricultural Sciences*, Vol. 15 No. 2, pp. 54-59.
- Gebre, G., Isoda, H., Rahut, D., Amekawa, Y. and Nomura, H. (2019), 'Gender differences in the adoption of agricultural technology: The case of improved maize varieties in southern Ethiopia', *In Women's Studies International Forum*, Vol. 76, pp. 102264. <https://doi.org/10.1016/j.wsif.2019.102264>.
- Gebremariam, G. and Tesfaye, W. (2018), 'The heterogeneous effect of shocks on agricultural innovations adoption: Microeconomic evidence from rural Ethiopia', *Food Policy*, Vol. 74 No. C, pp. 154-61.
- Genet, Y., Nigussie, A., Fikre, T., Assefa, K. and Musema, R. (2020), 'Onfarm Validation of Agricultural Technologies for Supporting Tef Extension Package Formulation in Ethiopia', *International Journal of Applied Agricultural Sciences*, Vol. 6 No. 6, pp. 172-178.
- Gessesse, E. (2020), 'Teff (*Eragrostis*) Row Planting Practice and its Contribution to Household Income in Ebinat District, Ethiopia', University of Gondar, Gondar.
- Getu, M. (2009), 'Ethiopian floriculture and its impact on the environment', *Mizan Law Review*, Vol. 3 No. 2, pp. 240-270.
- Gianessi, L. (2013), 'The increasing importance of herbicides in worldwide crop production', *Pest Management Science*, Vol. 69 No. 10, pp. 1099-1105.
- Gianessi, L. and Williams, A. (2011), 'Overlooking the obvious: The opportunity for herbicides in Africa', *Outlooks on Pest Management*, Vol. 22 No. 5, pp. 211-215.
- Gockowski, J. and Ndoumbe, M. (2004), 'The Adoption of Intensive Monocrop Horticulture in Southern Cameroon', *Agricultural Economics*, Vol. 30 No. 3, pp. 195-202.
- Gold, C. (1993), 'Effects of cassava intercropping and varietal mixtures on herbivore load, plant growth, and yields: applications for small farmers in Latin America', in Altieri, M. (Ed.) *Crop Protection Strategies for Small Farmers*, Westview Press, London, pp. 117-141.
- Graziosi, I., Tembo, M., Kuate, J. and Muchugi, A. (2020). 'Pests and diseases of trees in Africa: A growing continental emergency', *Plants, People, Planet*, Vol. 2 No. 1, pp. 14-28.
- Green, J. (2012), 'The benefits of herbicide-resistant crops', *Pest Management Science*, Vol. 68 No. 10, pp. 1323-1331.
- Gross, E., Lahkar, B., Subedi, N., Nyirenda, V., Lichtenfeld, L. and Jakoby, O. (2018), 'Seasonality, crop type and crop phenology influence crop damage by wildlife herbivores in Africa and Asia', *Biodiversity and Conservation*, Vol. 27 No. 8, pp. 2029-2050.
- Haggblade, S., Diarra, A., Jiang, W., Assima, A., Keita, N., Traore, A. and Traore, M. (2021), 'Fraudulent pesticides in West Africa: A Quality Assessment of Glyphosate Products in Mali', *International Journal of Pest Management*, Vol. 67 No. 1, pp. 32-45.
- Hailu, G., Weersink, A. and Minten, B. (2017), 'Determinants of the productivity of teff in Ethiopia', *The European Journal of Development Research*, Vol. 29 No. 4, pp. 866-892.
- Hair J., Matthews, L., Matthews, R. and Sarstedt, M. (2017), 'PLS-SEM or CB-SEM: updated guidelines on which method to use', *International Journal of Multivariate Data Analysis*, Vol. 1 No. 2, pp. 107-123.
- Hair, J., Ringle, C. and Sarstedt, M. (2011), 'PLS-SEM: Indeed a Silver Bullet', *Journal of Marketing Theory and Practice*, Vol. 19 No. 2, pp. 139-152.
- Haverkort, A. and Bicamumpaka, M. (1986), 'Correlation between intercepted radiation and yield of potato crops infested by *Phytophthora infestans* in central Africa', *Netherlands Journal of Plant Pathology*, Vol. 92 No. 5, pp. 239-247.
- Hempel, L. (1999), 'Conceptual and analytical challenges in building sustainable communities', in Mazmanian, D. and Kraft, M. (Eds.) *Toward Sustainable Communities:*

- Transition and Transformation in Environmental Policy*, MIT Press, Cambridge MA, pp. 43–74.
- Hossard, L., Philibert, A., Bertrand, M., Colnenne-David, C., Debaeke, P., Munier-Jolain, N., Jeuffroy, M., Richard, G. and Makowski, D. (2014), 'Effects of halving pesticide use on wheat production', *Scientific Reports*, Vol. 4 No. 1, pp. 1-7.
- Howeler, R. and Cadavid, L. (1990), 'Short-and long-term fertility trials in Colombia to determine the nutrient requirements of cassava', *Fertilizer Research*, Vol. 26 No. 1, pp. 61-80.
- Imfeld, G. and Vuilleumier, S. (2012), 'Measuring the effects of pesticides on bacterial communities in soil: a critical review', *European Journal of Soil Biology*, Vol. 49 No. 22-30. <http://dx.doi.org/10.1016/j.ejsobi.2011.11.010>.
- Iqbal, S., Khalid, U., Awan, T., Iqbal, N., Saleem, M., Iram, A., Raza, M., Riaz, A., Latif, T. and Ahmad, N. (2019), 'Qualitative response of super basmati rice to different nitrogen levels under varying rice ecosystem', *African Journal of Agricultural Research*, Vol. 14 No. 9, pp. 548-558.
- Jacoby, H. (2000), 'Access to markets and the benefits of rural roads', *The Economic Journal*, Vol. 110 No. 465, pp. 713-737.
- Jacques, P. and Jacques, J. (2012), 'Monocropping cultures into ruin: the loss of food varieties and cultural diversity', *Sustainability*, Vol. 4 No. 11, pp. 2970-2997.
- Jayne, T., Muyanga, M., Wineman, A., Ghebru, H., Stevens, C., Stickler, M., Chapoto, A., Anseeuw, W., van der Westhuizen, D. and Nyange, D. (2019), 'Are medium-scale farms driving agricultural transformation in sub-Saharan Africa?', *Agricultural Economics*, Vol. 50 No. 1, pp. 75-95.
- Kang, M. and Jung, C. (2017), 'Avoidance behavior of honey bee, *Apis mellifera* from commonly used fungicides, acaricides and insecticides in apple orchards', *Journal of Apiculture*, Vol. 32 No. 4, pp. 295-302.
- Kapsi, M., Tsoutsis, C., Paschalidou, A. and Albanis, T. (2019), 'Environmental monitoring and risk assessment of pesticide residues in surface waters of the Louros River', *Science of The Total Environment*, Vol. 650 No. 2, pp. 2188-2198.
- Kassaw, A., Abera, M. and Belete, E. (2021), 'The response of potato late blight to potato varieties and fungicide spraying frequencies at meket, Ethiopia', *Cogent Food & Agriculture*, Vol. 7 No. 1, p.1870309.
- Kebede, D., Ketema, M., Dechassa, N. and Hundessa, F. (2017), 'Determinants of adoption of wheat production technology package by smallholder farmers: Evidences from eastern Ethiopia', *Turkish Journal of Agriculture-Food Science and Technology*, Vol. 5 No. 3, pp. 267-274.
- Kebede, M. and Anbasa, F. (2017), 'Efficacy of pre-emergence herbicides for the control of major weeds in maize (*Zea mays* L.) at Bako, Western Oromia, Ethiopia', *American Journal of Agriculture and Forestry*, Vol. 5 No. 5, pp. 173-180.
- Kecha, M. (2020), 'Evaluation of some fungicides for Management of Sugar cane rust (*Puccinia melanocephala*) disease in Finchaa Sugar Estate, Horo-Guduru Wollega, Oromia Regional State, Ethiopia', *International Journal of Agricultural Science and Food Technology*, Vol. 6 No. 2, pp. 204-210.
- Kock, N. (2019), *WarpPLS User Manual: Version 7.0.*, ScriptWarp Systems, Texas.
- Lakshman, Y. (1993), 'What Are Improved Seeds? An Epistemology of the Green Revolution', *Economic Geography*, Vol. 69 No. 3, pp. 254-273.
- Lalitha, M. and Chhabra, V. (2020), 'Response of Oilseed Crops to Planting Methods and Spacing', *IAR Journal of Agriculture Research and Life Sciences*, Vol. 1 No. 6, pp. 172-175.
- Li, M., Li, R., Zhang, J., Liu, S., Hei, Z. and Qiu, S. (2019), 'A combination of rice cultivar mixed-cropping and duck co-culture suppressed weeds and pests in paddy fields', *Basic and Applied Ecology*, Vol. 40, pp. 67-77. <https://doi.org/10.1016/j.baae.2019.09.003>.
- Loha, K., Lamoree, M., Weiss, J. and de Boer, J. (2018), 'Import, disposal, and health impacts of pesticides in the East Africa Rift (EAR) zone: A review on management and policy analysis', *Crop Protection*, Vol. 112, pp. 322-331, <https://doi.org/10.1016/j.cropro.2018.06.014>.

- Lusembo, P., Ebong, C. and Sabiiti, E. (1998), 'Integration of cassava tuber and forage legume seed production for sustained soil fertility', *Tropical Agriculture*, Vol. 75 No 1-2, pp. 18-20.
- Luthans, F. and Stewart, T. (1977), 'A general contingency theory of management', *Academy of Management Review*, Vol. 2 No. 2, pp. 181-195.
- Mahajan, G., Rachaputi, R. and Chauhan, B. (2019), 'Integrated weed management using row arrangements and herbicides in pigeonpea (*Cajanus cajan*) in Australia', *Crop and Pasture Science*, Vol. 70 No. 8, pp. 676-683.
- Mariyono, J., Kuntariningsih, A. and Kompas, T. (2018), 'Pesticide use in Indonesian vegetable farming and its determinants', *Management of Environmental Quality*, Vol. 29 No. 2, pp. 305-323.
- Matthews, G., Wiles, T. and Baleguel, P. (2003), 'A survey of pesticide application in Cameroon', *Crop Protection*, Vol. 22 No. 5, pp. 707-714.
- Maxwell, S. and Fernando, A. (1989), 'Cash Crops in Developing Countries: The Issues, The Facts, The Policies', *World Development*, Vol. 17 No. 11, pp. 1677-1708.
- Mbah, T., Obike, A., Ejiofor, E., Ikechukwu, G., Mbah, G. and Omeh, Y. (2018), 'Effect of fungicide application on some nutritional and anti-nutritional composition of *Colocassia esculenta* (Cocoyam)', *Journal of Agronomy*, Vol. 17 No. 3, pp. 147-153.
- McGuire, S. and Sperling, L. (2011), 'The links between food security and seed security: facts and fiction that guide response', *Development in Practice*, Vol. 21 No. 4-5, pp. 493-508.
- McMaster, G., Aiken, R. and Nielsen, D. (2000), 'Optimizing wheat harvest cutting height for harvest efficiency and soil and water conservation', *Agronomy Journal*, Vol. 92 No. 6, pp. 1104-1108.
- Mekonen, S., Ambelu, A. and Spanoghe, P. (2014), 'Pesticide residue evaluation in major staple food items of Ethiopia using the Quechers method: A case study from the Jimma zone', *Environmental Toxicology and Chemistry*, Vol. 33 No. 6, pp. 1294-1302.
- Mengistie, B., Mol, A. and Oosterveer, P. (2017), 'Pesticide use practices among smallholder vegetable farmers in Ethiopian Central Rift Valley', *Environment, Development and Sustainability*, Vol. 19 No. 1, pp. 301-324.
- Mitiku, A. and Dalga, D. (2014), 'Effect of herbicides on weed dynamics and yield and yield attribute of bread wheat (*Triticum aestivum* L.) in south eastern part of Ethiopia', *International Journal of Technology Enhancements and Emerging Engineering Research*, Vol. 2 No. 4, pp. 130-133.
- MoARD (2007), 'Livestock Development Master Plan Study. Phase I Report - Data Collection and Analysis', Ministry of Agriculture and Rural Development, Addis Ababa.
- Morton, V. and Staub, T. (2008), *A short history of fungicides*, The American Phytopathological Society, Minnesota.
- Muzhinji, N., Woodhall, J., Truter, M. and Van der Waals, J. (2018), 'Variation in fungicide sensitivity among *Rhizoctonia* isolates recovered from potatoes in South Africa', *Plant Disease*, Vol. 102 No. 8, pp. 1520-1526.
- Namugga, P., Melis, R., Sibiya, J. and Barekye, A. (2017), 'Participatory assessment of potato farming systems, production constraints and cultivar preferences in Uganda', *Australian Journal of Crop Science*, Vol. 11 No. 8, pp. 932-940.
- Negatu, B., Kromhout, H., Mekonnen, Y. and Vermeulen, R. (2017), 'Occupational pesticide exposure and respiratory health: a large-scale cross-sectional study in three commercial farming systems in Ethiopia', *Thorax*, Vol. 72 No. 6, pp. 498-499.
- Nuyttens, D., Devarrewaere, W., Verboven, P. and Foqué, D. (2013), 'Pesticide-laden dust emission and drift from treated seeds during seed drilling: a review', *Pest Management Science*, Vol. 69 No. 5, pp. 564-575.
- O'Mara, A., Marsh, H., Craven, R. and Debus, R. (2006), 'Do self-concept interventions make a difference? A synergistic blend of construct validation and meta-analysis', *Educational Psychologist*, Vol. 41 No. 3, pp. 181-206.
- Ochieng, J., Schreinemachers, P., Ogada, M., Dinssa, F., Barnos, W. and Mndiga, H. (2019), 'Adoption of improved amaranth varieties and good agricultural practices in East

- Africa', *Land Use Policy*, Vol. 83, pp. 187-194.
<https://doi.org/10.1016/j.landusepol.2019.02.002>
- Odurukwe, S. and Ikeorgu, J. (1994), 'Effects of fertilizer and time of introduction of cassava in yam/maize/cassava intercrop on component yield', *Acta Horticulturae*, Vol. 380, pp. 72-77. [10.17660/ActaHortic.1994.380.11](https://doi.org/10.17660/ActaHortic.1994.380.11).
- Olasantan, F., Ezumah, H. and Lucas, E. (1996), 'Response of cassava and maize to fertilizer application, and a comparison of the factors affecting their growth during intercropping' *Nutrient Cycling in Agroecosystems*, Vol. 46 No.3, pp. 215-223.
- Oliver, R. and Hewitt, H. (2014), *Fungicides in Crop Protection*, CAB International, Oxfordshire.
- Olivera F., Newcomb, M., Flath, K., Sommerfeldt-Impe, N., Szabo, L., Carter, M., Luster, D. and Jin, Y. (2017), 'Characterization of Puccinia graminis f. sp. tritici isolates derived from an unusual wheat stem rust outbreak in Germany in 2013', *Plant Pathology*, Vol. 66 No. 8, pp. 1258-1266.
- Omamo, S. (1998), 'Farm- to- market transaction costs and specialisation in small- scale agriculture: Explorations with a non- separable household model', *The Journal of Development Studies*, Vol. 35 No. 2, pp. 152-163.
- Orchard, S., Stringer, L. and Manyatsi, A. (2017), 'Farmer perceptions and responses to soil degradation in Swaziland', *Land Degradation & Development*, Vol. 28 No. 1, pp. 46-56.
- Orth, R., Marion, S., Granger, S. and Traber, M. (2009), 'Evaluation of a mechanical seed planter for transplanting Zostera marina (eelgrass) seeds', *Aquatic Botany*, Vol. 90 No. 2, pp. 204-208.
- Peet, E., Fink, G. and Fawzi, W. (2015), 'Returns to education in developing countries: Evidence from the living standards and measurement study surveys', *Economics of Education Review*, Vol. 49 No. C, pp. 69-90.
- Pratt, C., Constantine, K. and Murphy, S. (2017), 'Economic impacts of invasive alien species on African smallholder livelihoods', *Global Food Security*, Vol. 14, No. 31-37. <https://doi.org/10.1016/j.gfs.2017.01.011>.
- Reddy, K. (2003), 'Impact of Rye Cover Crop and Herbicides on Weeds, Yield, and Net Return in Narrow-Row Transgenic and Conventional Soybean (Glycine max) 1', *Weed Technology*, Vol. 17 No. (1), pp. 28-35.
- Rhoades, R. and Bebbington, A. (1990), 'Mixing it up: Variations in Andean farmers' rationales for intercropping of potatoes', *Field Crops Research*, Vol. 25 No. 1-2, pp. 145-156.
- Rock, J. (2019) 'We are not starving': challenging genetically modified seeds and development in Ghana', *Culture, Agriculture, Food and Environment*, Vol. 41 No. 1, pp. 15-23.
- Rodenburg, J., Johnson, J., Dieng, I., Senthilkumar, K., Vandamme, E., Akakpo, C., Allarangaye, M., Baggie, I., Bakare, S., Bam, R. and Bassoro, I. (2019), 'Status quo of chemical weed control in rice in sub-Saharan Africa', *Food Security*, Vol. 11 No. 1, pp. 69-92.
- Rüegg, W., Quadranti, M. and Zoschke, A. (2007), 'Herbicide research and development: challenges and opportunities', *Weed Research*, Vol. 47 No. 4, pp. 271-275.
- Runo, S., Alakonya, A., Machuka, J. and Sinha, N. (2011), 'RNA interference as a resistance mechanism against crop parasites in Africa: a 'Trojan horse' approach', *Pest Management Science*, Vol. 67 No. 2, pp. 129-136.
- Sajid, M., Muhammad, N., Zakaria, R., Shahbaz, A. and Nauman, A. (2020) 'Associated Factors of Cardiovascular Diseases in Pakistan: Assessment of Path Analyses Using Warp Partial Least Squares Estimation', *Pakistan Journal of Statistics and Operation Research*, Vol. 16 No. 2, pp. 265-277.
- Sargent, T. (1993), *Bounded rationality in macroeconomics: The Arne Ryde Memorial Lectures*, Oxford University Press, Oxford.
- Schmidt, E., Dorosh, P., Jemal, M. and Smart, J. (2018), *Ethiopia's Spatial and Structural Transformation: Public Policy and Drivers of Change*, International Food Policy Research Institute, Addis Ababa.

- Schoar, A. (2010), 'The divide between subsistence and transformational entrepreneurship', *Innovation Policy and the Economy*, Vol. 10 No. 1, pp. 57-81.
- Shaw, A. and Wilson, K. (2020), 'The Bill and Melinda Gates Foundation and the necro-populationism of 'climate-smart'agriculture', *Gender, Place & Culture*, Vol. 27 No. 3, pp. 370-393.
- Shita, A., Kumar, N. and Singh, S. (2018), 'Agricultural technology adoption and its determinants in Ethiopia: a reviewed paper', *Asia Pacific Journal of Research*, Vol. 1 No. 55, pp. 99-104.
- Sisay, B., Simiyu, J., Malusi, P., Likhayo, P., Mendesil, E., Elibariki, N., Wakgari, M., Ayalew, G. and Tefera, T. (2018), 'First report of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), natural enemies from Africa', *Journal of Applied Entomology*, Vol. 142 No. 8, pp. 800-804.
- Sissons, P., Green, A. and Broughton, K. (2019), 'Inclusive growth in English cities: Mainstreamed or Sidelined?', *Regional Studies*, Vol. 53 No. 3, pp. 435-446
- Slaper, T. and Hall, T. (2011), 'The triple bottom line: What is it and how does it work', *Indiana Business Review*, Vol. 86 No. 1, pp. 4-8.
- Steinberg, G. and Gurr, S. (2020), 'Fungi, fungicide discovery and global food security', *Fungal Genetics and Biology*, Vol. 144, p.103476. <https://doi.org/10.1016/j.fgb.2020.103476>.
- Stifel, D. and Minten, B. (2017), 'Market access, well-being, and nutrition: evidence from Ethiopia', *World Development*, Vol. 100 No. 90, pp. 229-241.
- Stokstad, E. (2017), 'New crop pest takes Africa at lightning speed'. *Science*, Vol. 356 No. 6337, pp. 473-474.
- Tamru, S., Minten, B., Alemu, D. and Bachewe, F. (2017), 'The rapid expansion of herbicide use in smallholder agriculture in Ethiopia: Patterns, drivers, and implications', *The European Journal of Development Research*, Vol. 29 No. 3, pp. 628-647.
- TechSci (2021), *Global Pesticides Market*, TechSci Research LLC, New York.
- Teklu, B., Adriaanse, P., Ter Horst, M., Deneer, J. and Van den Brink, P. (2015), 'Surface water risk assessment of pesticides in Ethiopia', *Science of the Total Environment*, Vol. 508, pp. 566-574. <https://doi.org/10.1016/j.scitotenv.2014.11.049>.
- Tessema, Y., Asafu- Adjaye, J. and Shiferaw, B. (2018), 'The impact of conservation tillage on maize yield and input demand: the case of smallholder farmers in north-west Ethiopia', *Australian Journal of Agricultural and Resource Economics*, Vol. 62 No. 4, pp. 636-653.
- Tolera, S. (2020), 'Systematic Review on Adverse Effect of Pesticide on Top Ten Importers of African Countries', *Journal of Applied Sciences and Environmental Management*, Vol. 24 No. 9, pp. 1607-1616.
- van der Meulen, H., Dolman, M., Jager, J. and Venema, G. (2014), 'The impact of farm size on sustainability of dutch dairy farms', *International Journal of Agricultural Management*, Vol. 3 No. 2, pp. 119-123.
- Vandecasteele, J., Dereje, M., Minten, B. and Taffesse, A. (2018), 'Labour, profitability and gender impacts of adopting row planting in Ethiopia', *European Review of Agricultural Economics*, Vol. 45 No. 4, pp. 471-503.
- Venugopal, S., Viswanathan, M. and Jung, K. (2015), 'Consumption constraints and entrepreneurial intentions in subsistence marketplaces', *Journal of Public Policy & Marketing*, Vol. 34 No. 2, pp. 235-251.
- Verkaart, S., Mausch, K., Claessens, L. and Giller, K. (2019), 'A recipe for success? Learning from the rapid adoption of improved chickpea varieties in Ethiopia', *International Journal of Agricultural Sustainability*, Vol. 17 No. 1, pp. 34-48.
- Wandiga, S. (2001), 'Use and Distribution of Organochlorine Pesticides. The Future in Africa', *Pure and Applied Chemistry*, Vol. 73 No. 7, pp. 1147-1155.
- Wigfield, A. (1994), 'Expectancy-value theory of achievement motivation: A developmental perspective', *Educational Psychology Review*, Vol. 6 (1), pp. 49-78.
- Williamson, S. (2003), *Pesticide Provision in Liberalized Africa: Out of Control? Agricultural Research and Extension Network Paper No. 126*, Overseas Development Institute, London.

- Williamson, S., Ball, A. and Pretty, J. (2008), 'Trends in pesticide use and drivers for safer pest management in four African countries', *Crop Protection*, Vol. 27 No. 10, 1327-1334.
- World Bank (2021a), Data: Ethiopia, Available online: <https://data.worldbank.org/country/ethiopia> [Accessed 10th March 2021].
- World Bank (2021b), Data: Ethiopia Agriculture & Rural Development, Available online: <https://data.worldbank.org/topic/agriculture-and-rural-development?locations=ET> [Accessed 10th March 2021].
- World Bank (2021c) *Microdata Library*, The World Bank Group, Washington DC
- Yigezu, Y., El-Shater, T., Boughlala, M., Bishaw, Z., Niane, A., Maalouf, F., Degu, W., Wery, J., Boutfiras, M. and Aw-Hassan, A. (2019). Legume-based rotations have clear economic advantages over cereal monocropping in dry areas, *Agronomy for Sustainable Development*, Vol. 39 No. 6, pp. 1-14.
- Zhai, Y., Wu, Q., Chen, G., Zhang, H., Yin, X. and Chen, F. (2018), 'Broadcasting Winter Wheat Can Increase Grain Yield without Reducing the Kernels per Spike and the Kernel Weight', *Sustainability*, Vol. 10 No. 12, pp. 1-16.
- Zhang, D., Li, H., Fu, Z., Cai, S., Xu, S., Zhu, H. and Shen, J. (2019), 'Increased planting density of Chinese milk vetch (*Astragalus sinicus*) weakens phosphorus uptake advantage by rapeseed (*Brassica napus*) in a mixed cropping system', *AoB Plants*, Vol. 11 No. 4, pp. 1-9.
- Zhang, R., Pan, Z., Wang, X., Shen, M., Zhou, J., Fu, Z. and Jin, Y. (2018), 'Short-term propamocarb exposure induces hepatic metabolism disorder associated with gut microbiota dysbiosis in adult male zebrafish', *Acta Biochimica Biophysica Sinica*, Vol. 51 No. 1, pp. 88-96.
- Zhang, W. (2018), 'Global pesticide use: Profile, trend, cost/benefit and more'. *International Academy of Ecology and Environmental Sciences*, Vol. 8 No. 1, pp. 1-27.