

## Psychosocial determinants of depression in the community of the elderly with cardiovascular disease

Item Type	Journal article
Authors	Xu, Man;Chen, Ruoling;Liu, Bing;Chai, Yun;Boer, Dorothy D;Hu, Ping;Hu, Zhi
Citation	Xu, M., Chen, R. et al. (2018) Psychosocial determinants of depression in the community of the elderly with cardiovascular disease, Psychiatry Research, 268 (October 2018), pp. 123-130.
DOI	<a href="https://doi.org/10.1016/j.psychres.2018.03.008">10.1016/j.psychres.2018.03.008</a>
Publisher	Elsevier
Journal	Psychiatry Research
Download date	2026-03-12 20:48:37
License	<a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Link to Item	<a href="http://hdl.handle.net/2436/621594">http://hdl.handle.net/2436/621594</a>

**Psychosocial determinants of depression in the community of the elderly with cardiovascular disease.**

Man Xu <sup>a,1</sup>, Ruoling Chen <sup>b,1</sup>, Bing Liu <sup>c</sup>, Yun Chai <sup>d</sup>, Dorothy D. Boer <sup>e</sup>, Ping Hu <sup>a</sup>, Zhi Hu <sup>f</sup>

<sup>a</sup> Key Laboratory of Environment and Health (HUST), Ministry of Education & Ministry of Environmental Protection, and State Key Laboratory of Environmental Health (Incubation), School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

<sup>b</sup> Center for Health and Social Care Improvement (CHSCI), University of Wolverhampton, Wolverhampton, United Kingdom

<sup>c</sup> Center of Health Administration and Development Studies, Hubei University of Medicine, Shiyan, Hubei, China

<sup>d</sup> School of Public Health and Health Management, Hubei University of Medicine, Shiyan, Hubei, China

<sup>e</sup> English Language Teaching Department, Hubei University of Medicine, Shiyan, Hubei, China

<sup>f</sup> School of Health Administrations, Anhui Medical University, Hefei, China

## **Abstract**

The co-morbidity of cardiovascular disease (CVD) and depression is quite frequent in old people, and some potential biological and behavioural mechanisms linking them have been reported. Yet the impact of psychosocial factors on depression in the elderly with CVD remains unclear. This study aimed to analyze the psychosocial determinants of depression in the elderly with CVD. Using the Geriatric Mental Status–Automated Geriatric Examination for Computer Assisted Taxonomy, a community-based household survey was performed in 2,199 elderly people from the Anhui cohort third-wave survey from 2007 to 2009 and an extended study in Hubei from 2010 to 2011. Multiple logistic regression analyses were employed to assess the influence of psychosocial factors on depression. Among them, the prevalence of depression was 4.77%. Three factors were associated with depression in elderly in the community: self-assessed physical health status, anything else severely upsetting and unpleasantness with relatives, friends, or neighbors. In particular, associations of psychosocial factors with depression were more evident in individuals with CVD. This study confirms several psychosocial determinants of depression and the impact of CVD on the associations among the elderly, which provides some clues for interventional strategies of late-life depression.

## **Keywords:**

Geriatric Mental Status–Automated Geriatric Examination for Computer Assisted Taxonomy; Psychological status; Negative life events; Social network; Activities of daily living; Depression; Older people; Cardiovascular disease

## **1.Introduction**

Depression is a complex public health problem and leads to global disease and economic burden (WHO,2014). Despite decades of widespread intervention and efforts to raise public awareness of depression, its prevalence rates were still stable (Ferrari et al., 2013).Cardiovascular disease (CVD) is often regarded as a frequent co-morbidity of depression (Chauvet-Gelinier et al., 2013; Gan et al., 2014; Ghaemmohamadi et al., 2017), and there is a bidirectional relationship between them (Chauvet-Gelinier et al., 2013; Hare et al., 2014). Coronary heart disease is present in 30% of patients with mood disorders (Chauvet-Gelinier et al., 2013), while depression can increase 30% risk of both coronary heart disease and myocardial infarction (Gan et al., 2014). Moreover, potential mechanisms linking depression and adverse cardiac outcomes include some plausible biological pathways, such as elevated inflammation and altered function in autonomic nervous system (Chauvet-Gelinier et al., 2013). Behavioral pathways are also involved, such as poor adherence to diet and medication treatments and a sedentary lifestyle (Chauvet-Gelinier et al., 2013).

Accumulated evidence among general older people has demonstrated associations of depression with some psychosocial elements, including unsatisfied one's quality of life, income and physical health (Davison et al., 2012; Feng et al., 2014), burden of more chronic diseases (Feng et al., 2014),greater levels of disability (Davison et al., 2012), negative life events (Chen et al., 2005; Feng et al., 2014), and social support (Glaesmer et al., 2011; Morikawa et al., 2013; Wang and Zhao, 2012). However, few psychosocial factors were reported to be associated with depression among older people with CVD. For example, depression was previously found to correlate with total scores of social support as well as two domains of subjective support and utilization among 81 Chinese elderly post-stroke patients (Zhang et al., 2011). Another cross-sectional study conducted among 1064 elderly subjects with hypertension living at home in China showed that individuals under conditions of illiteracy, experiencing a pronounced stressful life event, or having poor functional status were more likely to develop depression symptoms (Ma et al., 2015).

Therefore, further investigations on effects of self-rated psychological status, negative life events, social network, and activities of daily life among aged residents with CVD are warranted, particularly given that the presence of CVD is usually associated with loss of activities of daily living (ADL) and increased demands for more various support. Besides, to date, the majority of previous studies assessing depression were based on self-rated questionnaires, which could be easily influenced by participants' subjective consciousness. To fill in these gaps, our study illustrated the psychosocial determinants of depression among old adults with CVD. Among those patients, psychosocial factors were hypothesized to be associated with depression, although their impacts on depression would be partially overlapped or offset. Our findings would add clues for more feasible psychological intervention strategies for preventing and controlling late-life depression.

## **2. Methods**

### *2.1. Study population*

The study population consisted of elderly patients from two community-based household survey studies: the Anhui cohort 3<sup>rd</sup> wave survey and an extended study in Hubei province with a common research protocol (Chen et al., 2013).

#### *2.1.1 The Anhui cohort 3<sup>rd</sup> wave survey*

The methods of the Anhui cohort study have been fully described previously (Chen et al., 2008; Chen et al., 2014; Chen et al., 2013; Chen et al., 2004; Chen et al., 2005). In brief, to collect baseline data based on the residency list from the committees of the village and the district (wave 1), we randomly selected a sample of 3,336 rural and urban residents (aged $\geq$ 60 years) who had lived for 5 years or longer in the survey districts of Anhui province during 2001 to 2003. As the survey was anonymous and the data was only used for scientific research, after some explanations about the purposes and contents of the survey with each participant or the closest adult care-giver, we obtained the permission for interview and oral informed consent. Ethical approval for the study was obtained from Anhui Medical University and the

Research Ethics Committee, University College London (Chen et al., 2005). One year after the baseline investigation, we re-examined 2,608 cohort members (wave 2). During 2007 to 2009, we conducted the 3<sup>rd</sup> wave survey and successfully followed up 1,757 surviving cohort members with a response rate of 82.4%.

### *2.1.2 An extended study in Hubei*

During 2010 to 2011, we extended the project to include the Hubei province. Two researchers in this survey team were trained at Anhui Medical University and gained the experience of having undertaken several surveys in the old, and transferred these skills to local research team members (Chen et al., 2004; Chen et al., 2005). The survey was conducted in Maojian subdistrict in Shiyan, Hubei, aiming to recruit more than 500 participants in each community. Ethical approval for the study was obtained from the Research Ethics Committee of Hubei University of Medicine. In total, we successfully obtained 501 complete questionnaires among 560 participants with a response rate of 89.5%.

### *2.2. Interview and data collection*

The main interview material contained a general health and risk factors record and the GMS questionnaire (Copeland et al., 1986). The general health and risk factors record, derived partially from the MRC-ALPHA study (Copeland et al., 1999) and the Scottish MONICA surveys (Chen and Tunstall-Pedoe, 2005), was used for collecting data on five aspects of participants in this study. Data obtained included: (1) demographic information, (2) self-assessed psychological aspects, (3) social network, (4) negative life events occurring in the past 2 years, (5) ADL, (6) CVD. Participants reported their level of difficulty in questions of ADL scale. The valid response was 'no difficulty alone' (score 0), 'manages alone with difficulty' (score 1), and 'cannot do alone' (score 2). The scale includes the following 14 items: taking a bath or all-over wash, washing hands and face, putting on shoes and stockings/socks, doing up buttons and zips, dressing yourself other than the above, getting to and using the washroom, getting in and out of bed, feeding yourself, shaving (men) or doing up your hair (women), cutting your own toenails, getting upstairs and downstairs, getting around the house, going out alone and taking medicine. We asked each participant whether he

or she had a doctor diagnosis of hypertension, heart disease, stroke, angina and recorded as yes/no. Those who had any type of these diseases were defined as participants with CVD. Depression was defined using the Geriatric Mental State-Automated Geriatric Examination for Computer-Assisted Taxonomy (GMS-AGECAT) (Copeland et al., 1986), which was fully described previously for depression diagnosis (Chen et al., 2004; Copeland et al., 1986) and validated in China (Chen et al., 2004).

### 2.3. *Statistical analysis*

Pearson's  $\chi^2$  tests were conducted to determine the distribution of depression according to demographic and psychological characteristics. Multiple logistic regression analysis was conducted in a full model to evaluate risk factors (psychosocial variables as independent variables and depression as the dependent variable) adjusting for demographic variables and CVD with a significant level of 0.05 to enter and to stay in the model. Subsequently, a stratification analysis was conducted in the multiple logistic regression analysis in the same model to separately assess associations of psychosocial variables with depression among participants classified by the presence of CVD.

Data analyses were performed using SPSS statistical package, version 20.0 (SPSS Inc., Chicago, IL, USA). Our missing data analysis procedures mainly involved data missing completely at random, and the method of missing data processing was conducted by replacing the missing values with the column mean / median (Engels and Diehr, 2003).

## **3. Results**

### 3.1. *Baseline characteristics of participants*

Figure 1 presented the flow chart of the source of participants from the two communities for analysis. Of the 2199 participants, 105 (4.77%) was diagnosed with depression. Among all the participants, 1063 (48.34%) had one or more types of doctor-diagnosed CVD, including hypertension (37.79%), heart disease (19.33%),

angina (6.41%) and stroke (4.50%). Among participants aged 60-100 years (M=74.12; SD=7.09) in our study, 1174(53.39%) were female, 1124(51.11%) lived in urban districts, and 1127(51.25%) were illiterate. And 1066(48.48%) participants were peasants and 450 (20.46%) were manual laborers.

Significant differences in distributions of depression were found according to all the demographic and psychosocial variables, except for variables of good relationships with parents ( $\chi^2=181.63$ ,  $P=0.178$ ). The prevalence of CVD was significantly different between depression and non-depression group ( $\chi^2 = 5.06$ ,  $P =0.024$ ) (Table 1).

### *3.2. Associations of psychosocial factors with depression*

Associations of depression with self-assessed physical health status and life events of anything else severely upsetting and unpleasantness with relatives, friends, or neighbors were significant among all the participants, after adjusting for gender, age, urban-rurality, education level, main occupation and CVD (Table 2).

### *3.3. Stratified analyses for association of psychosocial factors with depression*

We subsequently conducted a stratification analysis in the participants by the presence of CVD. After adjusting for the same covariates as in the Table 2, associations of psychosocial factors with depression were separately found among participants with and without CVD. In participants with CVD, self-assessed physical health status, death of closely related person, and unpleasantness with relatives, friends, or neighbors were significantly associated with depression. But only the life event of anything else severely upsetting was associated with depression among participants without CVD (Table 3).

## **4. Discussion**

In our study, the prevalence of depression was 4.77%, which was similar to that found in 7,072 community elderly from six provinces of China based on the same diagnosis procedure (Chen et al., 2013). As expected, the associations of depression with self-assessed psychological status and negative life events were confirmed among older people from the community. But more associations were only seen in old

patients with CVD than those without CVD, when participants were stratified by the presence of CVD. Those findings provide evidence on implications of several psychosocial determinants of depression and the impact of CVD on these associations in the older population, which offers some potential ways to reduce later-life depression.

#### *4.1. Impact of psychosocial factors on depression in old people*

Later lifetime is often featured by a series of loss experiences, such as bereavement, loss of functional role in society, and isolation due to chronic diseases (Stephens MA, 1990). In our study, the aged who experienced negative life events, such as unpleasant things with intimate friends, neighbors or relatives, or anything else severely upsetting, were more likely to suffer from depression. Devanand et al. (Devanand et al., 2002) conducted a survey of 50 American elderly outpatients and 40 health controls in an American Late-Life Depression clinic, and found that individuals with major depression reported higher prevalence of adverse life events about interpersonal conflicts than controls. Interpersonal conflicts included major problems with friends/neighbors and major family problems/conflicts other than with spouse. This study was in line with our results. But in a community survey of 654 aged residents in an district in London, end of relationship and problem with close friend or relative were not associated with pervasive depression (Prince et al., 1997). Some other similar studies just showed negative effect of stressful life events on later-life depression but life events were not assessed in specific types (Feng et al., 2014; Luppá et al., 2012). The inconsistent conclusions may be caused by differences in sample size, evaluation methods of life events and culture backgrounds of various countries. In addition, according to socioemotional selectivity theory (Carstensen, 1995), aging is associated with motivational shifts that greater importance is paid to current emotional goals rather than instrumental goals. To maximize their positive emotional experience and minimize their emotional risk in the limited life, old people would selectively narrow their social communication and pay more attention to people who are close to them. Unsurprisingly, old people who experienced the two emotional stressful events were more likely to develop depression in our study.

Also, our study found that community elderly people with poor self-assessed health status were more likely to have depression than those with satisfactory self-assessed health. Meanwhile, those with the presence of CVD or those with higher scores on dependency of ADL had a higher prevalence of depression. The results were in line with findings from two previous cross-sectional studies (Davison et al., 2012; Nicolosi et al., 2011). In a case-control study of 50 aged care residents with a diagnosis of major depressive disorder and 50 matched residents without depression, higher level of depressive symptoms was associated with poorer self-rated health, more chronic health problems and greater levels of disability (Davison et al., 2012). In another study of 303 Brazilian elderly in a poor subdistrict of the city, individuals with more depressive symptoms were those with poorer self-perceived health and more health problems (Nicolosi et al., 2011). However, compared with the above two studies, our study had available formal diagnosis of depression using GMS-AGECAT, while they just assessed the severity of depression by geriatric depression scales. In addition, our study was based on the general community dwelling elderly with much larger sample size, thus have more reliable conclusions.

Stratification analysis showed that associations of self-assessed physical health, death of closely related person, unpleasantness with relatives, friends, or neighbor with depression were more pronounced for aged adults with CVD. In fact, CVD often has a long disease course and requires daily care and long-term medications, during which negative consequences of the persistent complications, decline of physical functions, heavy financial burden, and increased dependence on people around may affect older CVD patients' self-rated health and cognitive functions (Ma et al., 2015; Zhang et al., 2017). Then, subsequent somatic dysfunction and emotional distress may trigger and exacerbate the onset and development of depression (Ma et al., 2015; Zhang et al., 2017). Meanwhile, such emotional stressful events could make them nervous, sad, and even contribute to a disturbing sense of helplessness or insecurity, thus may also result in depression when accumulated for a long time. Moreover, chronic mental stress is a probable common agitator underlying the co-morbidity between depression and cardiovascular pathology by shifting the homeostatic balance

in the autonomic nervous system (Halaris, 2013). Besides, for the elderly without CVD, only the life event of anything else severely upsetting was associated with depression. We may speculate that for the old without somatic and psychological consequences of CVD, only the life events that severely make one upsetting may put them into depression. These findings suggested that, compared with older adults without CVD, stressful life events and poor self-reported physical health were more likely to be risk factors of depression among old people with CVD. Although the underlying pathophysiological or psychosocial mechanisms are unclear, it is possible that suffering from CVD may make the old more susceptible to those emotional stressful events and poor self-rated health status, which ,to a great extent, facilitates the process of geriatric depression onset.

#### *4.2. Implications for clinical practice*

Depression and CVD are at present the two most common disease that cause disability in high-income countries and this trend is expected to expand for countries of all income levels by 2030 (WHO,2008). Strategies to alleviate depression in old population should be a combined effect of the whole society, the community and family members. On one hand, public health planners and practitioners need give more consideration to developing comprehensive disease management for the old, including timely screening of CVD and depression, cardiac rehabilitation and exercise, cognitive behavioral therapy and antidepressant medication (Hare et al., 2014). On the other hand, previous studies showed that worse family functioning and lower social support were negatively associated with depression in Chinese elderly (Wang and Zhao, 2012);family emotional involvement had a buffering effect on mental symptoms of Chinese community elderly with chronic medical diseases (Leung et al., 2007).Therefore, good family function and various social support (i.e., emotional or material support) from people around them should be necessarily needed for the elderly to maintain their emotional fulfillment. Especially, specific measures should be taken to reduce the severity of grief and depression for older patients who experienced death of closely related person. Physicians in primary care should pay more attention to recognizing and appropriately referring for their emotional distress

in the context of death of the close, as standard depression care management may not be effective (Ghesquiere et al., 2013; Ghesquiere et al., 2014). Only in this context could targeted psychosocial interventions be taken to reduce depression risk of the elderly.

#### *4.3. Strengths and limitations*

The main contribution of this study lies in demonstrating the impact of psychosocial variables on depression among older adults and explaining the role of CVD in these associations. Also, whereas other studies used questionnaires as a substitute for depression diagnosis, we conducted a standardized face-to-face interview and depression diagnostic system using the GMS-AGECAT package. However, some limitations should be mentioned. First, we defined medically CVD based on information obtained from participants or their closest caregivers instead of accessing medical records. This approach might underestimate the prevalence of CVD diagnosed by doctors, because participants may not recall whether they have been given a diagnosis or may be unwilling to disclose this information. We acknowledge that more robust information should have been obtained through checking relevant medical records. Secondly, most assumed risk factors were collected through self-assessment, such as self-assessed psychological status, social network, and adverse life events (as in many European studies). Hence, any of those could cause bias if adverse data was reported by depressed participants. In addition, the sample size was relatively small due to the low prevalence of depression in communities and the study was cross-sectional. Nevertheless, we were able to identify certain high-risk groups which might benefit from psychological interventions, but the causal relationship between depression, CVD and correlative psychosocial factors requires longitudinal studies with a larger sample size for confirmation.

#### *4.4. Final conclusions*

Our studies confirmed associations of depression with self-assessed psychological status and negative life events among older people from the community and found the impact of CVD on the associations. From this study, we can propose that what probably affects patients is not only those psychosocial determinants, but also the way

in which physical limitations alter their lives due to CVD. It is possible that psychosocial factors may act as a stress-physiology mediator, a node through which CVD exert influence on both physiological functions and mental health of the old. However, our cross-sectional study lacks evidence for causal relationship. Thus, more longitudinal researches are needed to establish the pathways.

## **Conflict of interest**

None

## **Acknowledgements**

We wish to acknowledge the two research consortia from Medical University of Anhui and Hubei University of Medicine as well as all the participants and interviewers in each research center.

## **Funding**

This work was supported by the Health Department of Hubei Province[grant number ZX2012-11]; the Science and Technology Department of Hubei Province [grant number 2014CFB646].


## References

- Carstensen, L.L., 1995. Evidence for a life-span theory of socioemotional selectivity. *Curr. Dir. Psychol. Sci.* 4, 151-156.
- Chauvet-Gelinier, J.C., Trojak, B., Verges-Patois, B., Cottin, Y., Bonin, B., 2013. Review on depression and coronary heart disease. *Arch. Cardiovasc. Dis.* 106, 103-110.
- Chen, R., Hu, Z., Wei, L., Qin, X., McCracken, C., Copeland, J.R., 2008. Severity of depression and risk for subsequent dementia: cohort studies in China and the UK. *Br. J. Psychiatry.* 193, 373-377.
- Chen, R., Hu, Z., Wei, L., Wilson, K., 2014. Socioeconomic status and survival among older adults with dementia and depression. *Br. J. Psychiatry.* 204, 436-440.
- Chen, R., Tunstall-Pedoe, H., 2005. Socioeconomic deprivation and waist circumference in men and women: The Scottish MONICA surveys 1989--1995. *Eur. J. Epidemiol.* 20, 141-147.
- Chen, R.L., Hu, Z., Chen, R.L., Ma, Y., Zhang, D., Wilson, K., 2013. Determinants for undetected dementia and late-life depression. *Br. J. Psychiatry.* 203, 203-208.
- Chen, R.L., Hu, Z., Qin, X., Xu, X.C., Copeland, J.R.M., 2004. A community-based study of depression in older people in Hefei, China - the GMS-AGECAT prevalence, case validation and socio-economic correlates. *Int. J. Geriatr. Psychiatry.* 19, 407-413.
- Chen, R.L., Wei, L., Hu, Z., Qin, X., Copeland, J.R.M., Hemingway, H., 2005. Depression in older people in rural China. *Arch. Intern. Med.* 165, 2019-2025.
- Copeland, J.R., Chen, R., Dewey, M.E., McCracken, C.F., Gilmore, C., Larkin, B., et al, 1999. Community-based case-control study of depression in older people. Cases and sub-cases from the MRC-ALPHA Study. *Br. J. Psychiatry.* 175, 340-347.
- Copeland, J.R., Dewey, M.E., Griffiths-Jones, H.M., 1986. A computerized psychiatric diagnostic system and case nomenclature for elderly subjects: GMS and AGECAT. *Psychol. Med.* 16, 89-99.
- Davison, T.E., McCabe, M.P., Knight, T., Mellor, D., 2012. Biopsychosocial factors related to depression in aged care residents. *J. Affect. Disord.* 142, 290-296.
- Devanand, D.P., Kim, M.K., Paykina, N., Sackeim, H.A., 2002. Adverse life events in elderly patients with major depression or dysthymic disorder and in healthy-control subjects. *Am. J. Geriatr. Psychiatry.* 10, 265-274.
- Engels, J.M., Diehr, P., 2003. Imputation of missing longitudinal data: a comparison of methods. *J. Clin. Epidemiol.* 56, 968-976.
- Feng, L., Li, P., Lu, C., Tang, W., Mahapatra, T., Wang, Y., et al, 2014. Burden and correlates of geriatric depression in the Uyghur elderly population, observation from Xinjiang, China. *PloS one.* 9, e114139. doi: 10.1371/journal.pone.0114139.
- Ferrari, A.J., Charlson, F.J., Norman, R.E., Patten, S.B., Freedman, G., Murray, C.J.,

- et al, 2013. Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Med.*10, e1001547. doi: 10.1371/journal.pmed.1001547.
- Gan, Y., Gong, Y., Tong, X., Sun, H., Cong, Y., Dong, X., et al, 2014. Depression and the risk of coronary heart disease: a meta-analysis of prospective cohort studies. *BMC psychiatry.*14. Available from: <<https://bmcp psychiatry.biomedcentral.com/articles/10.1186/1471-244X-14-7>>
- Ghaemmohamadi, M.S., Behzadifar, M., Ghashghaee, A., Mousavinejad, N., Ebadi, F., Saeedi Shahri, S.S., et al, 2017. Prevalence of depression in cardiovascular patients in Iran: A systematic review and meta-analysis from 2000 to 2017. *J. Affect. Disord.*227, 149-155.
- Ghesquiere, A., Shear, M.K., Duan, N., 2013. Outcomes of bereavement care among widowed older adults with complicated grief and depression. *J. Prim. Care Community Health.* 4, 256-264.
- Ghesquiere, A.R., Park, M., Bogner, H.R., Greenberg, R.L., Bruce, M.L., 2014. The effect of recent bereavement on outcomes in a primary care depression intervention study. *Am. J. Geriatr. Psychiatry.* 22, 1555-1564.
- Glaesmer, H., Riedel-Heller, S., Braehler, E., Spangenberg, L., Lupp, M., 2011. Age- and gender-specific prevalence and risk factors for depressive symptoms in the elderly: a population-based study. *Int. Psychogeriatr.*23, 1294-1300.
- Halaris, A., 2013. Co-morbidity between cardiovascular pathology and depression: role of inflammation. *Mod. Trends Pharmacopsychiatry.* 28, 144-161.
- Hare, D.L., Toukhsati, S.R., Johansson, P., Jaarsma, T., 2014. Depression and cardiovascular disease: a clinical review. *Eur. Heart. J.* 35, 1365-1372.
- Leung, K.K., Chen, C.Y., Lue, B.H., Hsu, S.T., 2007. Social support and family functioning on psychological symptoms in elderly Chinese. *Arch. Gerontol. Geriatr.*44, 203-213.
- Lupp, M., Sikorski, C., Luck, T., Weyerer, S., Villringer, A., Konig, H.H., et al, 2012. Prevalence and risk factors of depressive symptoms in latest life--results of the Leipzig Longitudinal Study of the Aged (LEILA 75+). *Int. J. Geriatr. Psychiatry.*27, 286-295.
- Ma, L., Tang, Z., Sun, F., Diao, L., Li, Y., Wang, J., et al, Risk factors for depression among elderly subjects with hypertension living at home in China, 2015. *Int. J. Clin. Exp. Med.* 8, 2923-2928.
- Morikawa, M., Okamoto, N., Kiuchi, K., Tomioka, K., Iwamoto, J., Harano, A., et al , 2013. Association between depressive symptoms and metabolic syndrome in Japanese community-dwelling older people: a cross-sectional analysis from the baseline results of the Fujiwara-kyo prospective cohort study. *Int. J. Geriatr. Psychiatry.* 28, 1251-1259.
- Nicolosi, G.T., da Silva Falcao, D.V., Tavares Batistoni, S.S., Lopes, A., Cachioni, M., Neri, A.L., et al, 2011. Depressive symptoms in old age: relations among sociodemographic and self-reported health variables. *Int. Psychogeriatr.* 23, 941-949.
- Prince, M.J., Harwood, R.H., Blizard, R.A., Thomas, A., Mann, A.H., 1997. Social

- support deficits, loneliness and life events as risk factors for depression in old age. *The Gospel Oak Project VI. Psychol. Med.*27, 323-332.
- Stephens, M.A., Crowther, J.G., Hobfoll, S.E., Tennenbaum D.I. (Eds.), 1990. *Stress and Coping in Later-Life Families*. Hemisphere Publishing Company, New York, pp. 131-152.
- Wang, J., Zhao, X., 2012. Family functioning and social support for older patients with depression in an urban area of Shanghai, China. *Arch. Gerontol. Geriatr.* 55, 574-579.
- WHO, 2014. Global health estimates 2014 summary tables. Available from: <[http://www.who.int/healthinfo/global\\_burden\\_disease/estimates/en/index2.html](http://www.who.int/healthinfo/global_burden_disease/estimates/en/index2.html)>.
- WHO, 2008. The global burden of disease: 2004 update. Available from: <[http://www.who.int/healthinfo/global\\_burden\\_disease/2004\\_report\\_update/en/](http://www.who.int/healthinfo/global_burden_disease/2004_report_update/en/)>.
- Zhang, H.Y., Zhou, T., Zhang, Y.Q., Zhang, Y., Xu, Y., 2011. Correlation between social support and depression in elderly stroke patients in the sequelae stage from five communities of Shanghai, China. *Neural. Regen. Res.*6, 1493-1497.
- Zhang, Y., Chen, Y., Ma, L., 2018. Depression and cardiovascular disease in elderly: Current understanding. *J. Clin. Neurosci.*47,1-5.





**Figure 1. Flow chart of the elderly for analysis from two community-based surveys (N=2199)**

**Table 1. Distribution of depression according to demographic and psychosocial variables.**

Variable	Non-depressed		Depressed		$\chi^2_{(df)}$	P
	n	%	n	%		
<b><u>Demographic factors</u></b>						
<b>Urban-rurality</b>						
Urban	1085	51.81	39	37.14	8.61 <sub>(1)</sub>	0.003
Rural	1009	48.19	66	62.86		
<b>Gender</b>						
Female	1102	52.63	72	68.57	10.22 <sub>(1)</sub>	0.001
Male	992	47.37	33	31.43		
<b>Age(years)</b>						
85-100	196	9.36	15	14.29	3.8 <sub>(2)</sub>	0.015
75-84	885	42.26	37	35.24		
60-74	1013	48.38	53	50.48		
<b>Education level</b>						
Illiterate	1061	50.67	66	62.86	10.73 <sub>(4)</sub>	0.030
Primary school	369	17.62	21	20		
Secondary school	301	14.37	7	6.67		
High secondary school	215	10.27	8	7.62		
College	148	7.07	3	2.86		
<b>Main occupation</b>						
Peasant	1004	47.95	62	59.05	11.91 <sub>(3)</sub>	0.008
Manual laborer	432	20.63	18	17.14		
Business/other (housewife)	252	12.03	17	16.19		
Officer/teacher	406	19.39	8	7.62		
<b><u>Self-assessed psychological status</u></b>						
<b>Satisfaction with current life</b>						
Poor	55	2.63	16	15.24	65.37 <sub>(2)</sub>	<0.0001
Fair	757	36.15	52	49.52		
Satisfactory	1282	61.22	37	35.24		
<b>Self-assessed physical health status</b>						
Poor	216	10.32	36	34.29	64.43 <sub>(2)</sub>	<0.0001
Fair	1019	48.66	51	48.57		
Good	859	41.02	18	17.14		
<b>Satisfaction with current income</b>						
Poor	135	6.45	26	24.76	56.82 <sub>(2)</sub>	<0.0001
Fair	952	45.46	52	49.52		
Satisfactory	1007	48.09	27	25.71		

(Continued)

Variable	Non-depressed		Depressed		$\chi^2_{(df)}$	P
	n	%	n	%		
<b><u>Social network</u></b>						
<b>Living with</b>						
Nobody	318	15.19	27	25.71	8.38 <sub>(1)</sub>	0.004
Family members /others	1776	84.81	78	74.29		
<b>Marital status</b>						
Widowed	84	4.01	5	4.76	12.14 <sub>(2)</sub>	0.002
Never married/divorced	493	23.54	40	38.1		
Married	1517	72.45	60	57.14		
<b>Available help when needed</b>						
No	93	4.44	14	13.33	17.08 <sub>(1)</sub>	<0.0001
Yes	2001	95.56	91	86.67		
<b>Satisfaction with the help and support from others</b>						
No	61	2.91	13	12.38	2472.70 <sub>(1)</sub>	<0.0001
Yes	2033	97.09	92	87.62		
<b>Good relationships with others and ease in acquiring friends</b>						
No	272	12.99	25	23.81	10.02 <sub>(1)</sub>	0.002
Yes	1822	87.01	80	76.19		
<b>Good relationships with parents</b>						
No	33	1.58	4	3.81	181.63 <sub>(1)</sub>	0.178
Yes	2061	98.42	101	96.19		
<b>Good relationships with neighbors</b>						
No	729	34.81	52	49.52	9.45 <sub>(1)</sub>	0.002
Yes	1365	65.19	53	50.48		
<b><u>Negative life events</u></b>						
<b>A sudden decline of health status</b>						
No	1604	76.6	60	57.14	20.56 <sub>(1)</sub>	<0.0001
Yes	490	23.4	45	42.86		
<b>Serious financial problems</b>						
No	1926	91.98	82	78.1	24.29 <sub>(1)</sub>	<0.0001
Yes	168	8.02	23	21.9		
<b>Death of closely related person</b>						
No	1790	85.48	77	73.33	11.51 <sub>(1)</sub>	0.001
Yes	304	14.52	28	26.67		

(Continued)

Variable	Non-depressed		Depressed		$\chi^2_{(df)}$	P
	n	%	n	%		
<b>Something important lost or stolen</b>						
No	2038	97.33	98	93.33	438.20 <sub>(1)</sub>	0.036
Yes	56	2.67	7	6.67		
<b>Anything else severely upsetting</b>						
No	1977	94.41	74	70.48	91.26 <sub>(1)</sub>	<0.0001
Yes	117	5.59	31	29.52		
<b>Unpleasantness with relatives, friends, or neighbors</b>						
No	2070	98.85	93	88.57	5941.83 <sub>(1)</sub>	<0.0001
Yes	24	1.15	12	11.43		
<b><u>ADL</u></b>						
<b>ADL (scores)</b>						
0	1696	80.99	74	70.48	20.13 <sub>(2)</sub>	<0.0001
1-4	282	13.47	14	13.33		
5-28	116	5.54	17	16.19		
<b>CVD</b>						
No	1093	52.2	43	40.95	5.06 <sub>(1)</sub>	0.024
Yes	1001	47.8	62	59.05		

**Table 2. Associations between psychosocial factors and depression among the community elderly ( $n=2199$ ).<sup>‡</sup>**

Variable	OR(95%CI)
<b><u>Self-assessed psychological status</u></b>	
<b>Satisfaction with current life</b>	
Poor	2.02(0.75,5.45)
Fair	1.45(0.78,2.71)
Satisfactory	Reference
<b>Self-assessed physical health status</b>	
Poor	3.27(1.59,6.73)*
Fair	1.68(0.91,3.12)
Good	Reference
<b>Satisfaction with current income</b>	
Poor	2.18(0.94,5.06)
Fair	1.1(0.56,2.19)
Satisfactory	Reference
<b><u>Social network</u></b>	
<b>Living with</b>	
Nobody	1.52(0.85,2.72)
Family members /others	Reference
<b>Marital status</b>	
Never married	1.21(0.41,3.62)
Widowed (including several divorces)	1.39(0.81,2.37)
Married	Reference
<b>Available help when needed</b>	
No	0.83(0.29,2.39)
Yes	Reference
<b>Satisfaction with the help and support from others</b>	
No	2.41(0.79,7.36)
Yes	Reference
<b>Good relationships with others and ease in acquiring friends</b>	
No	0.9(0.48,1.67)
Yes	Reference
<b>Good relationships with parents</b>	
No	2.2(0.66,7.4)
Yes	Reference
<b>Good relationships with neighbors</b>	
No	1.16(0.71,1.9)
Yes	Reference

(continued)

Variable	OR(95% CI)
<b><u>Negative life events</u></b>	
<b>A sudden decline in health status</b>	
Yes	1.23(0.75,2)
No	Reference
<b>Serious financial problems</b>	
Yes	0.83(0.43,1.57)
No	Reference
<b>Death of closely related person</b>	
Yes	1.6(0.93,2.76)
No	Reference
<b>Something important lost or stolen</b>	
Yes	0.97(0.37,2.55)
No	Reference
<b>Anything else severely upsetting</b>	
Yes	3.04(1.64,5.62)**
No	Reference
<b>Unpleasantness with relatives, friends, or neighbors</b>	
Yes	4(1.55,10.31)*
No	Reference
<b><u>ADL</u></b>	
<b>ADL (scores)</b>	
1-4	0.84(0.44,1.6)
5-28	1.06(0.51,2.21)
0	Reference
<b>CVD</b>	
Yes	1.67(1.05,2.65)*
No	Reference

†Adjusted for the gender, age, urban-rurality, education level and main occupation.

\*\*  $P < 0.001$

\*  $P < 0.05$

**Table 3. Associations of psychosocial factors with depression stratified by CVD\*.**

Variable	Participants without CVD		Participants with CVD	
	<i>N(% of total 1136)</i>	<i>OR(95%CI) for depression</i>	<i>N (% of total 1063)</i>	<i>OR(95%CI) for depression</i>
<b><u>Self-assessed psychological status</u></b>				
<b>Satisfaction with current life</b>				
Poor	40(3.66)	2.75(0.64,11.8)	15(1.50)	1.21(0.22,6.68)
Fair	427(39.07)	1.2(0.45,3.18)	330(32.97)	2.17(0.88,5.32)
Satisfactory	626(57.27)	Reference	656(65.53)	Reference
<b>Self-assessed physical health status</b>				
Poor	91(8.33)	1.96(0.61,6.38)	125(12.49)	4.62(1.73,12.37) *
Fair	524(47.94)	1.31(0.53,3.25)	495(49.45)	2.25(0.92,5.52)
Good	478(43.73)	Reference	381(38.06)	Reference
<b>Satisfaction with current income</b>				
Poor	83(7.59)	3.25(0.77,13.73)	52(5.19)	1.55(0.49,4.9)
Fair	539(49.31)	2.01(0.64,6.26)	413(41.26)	0.58(0.22,1.53)
Satisfactory	471(43.09)	Reference	536(53.55)	Reference
<b>Living with</b>				
Nobody	174(15.92)	1.59(0.62,4.06)	144(14.39)	1.35(0.6,3.03)
Family members/others	919(84.08)	Reference	857(85.61)	Reference
<b><u>Social network</u></b>				
<b>Marital status</b>				
Never married	43(3.93)	2.21(0.46,10.67)	41(4.10)	0.65(0.11,3.78)
Widowed(including several divorces)	261(23.88)	0.93(0.38,2.28)	232(23.18)	1.78(0.87,3.64)
Married	789(72.19)	Reference	728(72.73)	Reference
<b>Available help when needed</b>				
No	46(4.21)	0.53(0.09,3.01)	47(4.70)	1.14(0.25,5.14)
Yes	1047(95.79)	Reference	954(95.30)	Reference
<b>Satisfaction with the help and support from others</b>				
No	31(2.84)	3.61(0.57,22.72)	30(3.00)	1.73(0.32,9.23)
Yes	1062(97.16)	Reference	971(97.00)	Reference
<b>Good relationships with others and ease in acquiring friends</b>				
No	155(14.18)	0.63(0.24,1.65)	117(11.69)	1.09(0.45,2.65)
Yes	938(85.82)	Reference	884(88.31)	Reference
<b>Good relationships with parents</b>				
No	19(1.74)	1.11(0.09,13.29)	14(1.40)	3.33(0.68,16.25)
Yes	1074(98.26)	Reference	987(98.60)	Reference
<b>Good relationships with neighbors</b>				
No	392(35.86)	1.69(0.78,3.68)	337(33.67)	0.89(0.45,1.76)
Yes	701(64.14)	Reference	664(66.33)	Reference

(Continued)

Variable	Participants without CVD		Participants with CVD	
	<i>N(% of total 1136)</i>	<i>OR(95%CI) for depression</i>	<i>N (% of total 1063)</i>	<i>OR(95%CI) for depression</i>
<b><u>Negative life events</u></b>				
<b>A sudden decline of health status</b>				
No	889(81.34)	1.18(0.51,2.75)	715(71.43)	1.29(0.69,2.39)
Yes	204(18.66)	Reference	286(28.57)	Reference
<b>Serious financial problems</b>				
No	1007(92.13)	0.43(0.13,1.37)	919(91.81)	1.41(0.63,3.16)
Yes	86(7.87)	Reference	82(8.19)	Reference
<b>Death of closely related person</b>				
No	958(87.65)	1.16(0.45,3.02)	832(83.12)	2.07(1.02,4.18) *
Yes	135(12.35)	Reference	169(16.88)	Reference
<b>Something important lost or stolen</b>				
No	1067(97.62)	2.37(0.67,8.37)	971(97.00)	0.16(0.02,1.5)
Yes	26(2.38)	Reference	30(3.00)	Reference
<b>Anything else severely upsetting</b>				
No	1041(95.24)	5.28(1.97,14.15)*	936(93.51)	2.17(0.91,5.15)
Yes	52(4.76)	Reference	65(6.49)	Reference
<b>Unpleasantness with relatives, friends, or neighbors</b>				
No	1082(98.99)	2.38(0.49,11.5)	988(98.70)	5.51(1.45,20.97)*
Yes	11(1.01)	Reference	13(1.30)	Reference
<b><u>ADL</u></b>				
<b>ADL (scores)</b>				
0	863(78.96)	0.7(0.24,2.04)	833(83.22)	0.92(0.38,2.21)
1-4	174(15.92)	1.61(0.5,5.16)	108(10.79)	1.01(0.37,2.74)
5-28	56(5.12)	Reference	60(5.99)	Reference

‡ Adjusted for the gender, age, urban-rurality, education level and main occupation.

\*  $P < 0.05$