

## The Influence of Hormonal Contraception on Vitamin D Supplementation on Serum 25(OH)D3 Status in Premenopausal Women: A Prospective Double-Blind Placebo Random Controlled Trial

Item Type	Journal article
Authors	Wyon, Matthew;Wolman, Roger;Nevill, Alan M.;Barber, Alison;Edwards, Marcia;Bowd, Belinda;Clarke, Frances;Bryant, Janine;Cloak, Ross
Citation	Wyon, M.A. et al. (2017) The Influence of Hormonal Contraception on Vitamin D Supplementation on Serum 25(OH)D3 Status in Premenopausal Women: A Prospective Double-Blind Placebo Random Controlled Trial, Journal of Endocrinology and Metabolism, 7(4), pp. 117-121.
DOI	<a href="https://doi.org/10.14740/jem441w">10.14740/jem441w</a>
Publisher	Elmer Press
Journal	Journal of Endocrinology and Metabolism
Download date	2026-03-12 19:31:58
License	<a href="https://creativecommons.org/licenses/by-nc/4.0/">https://creativecommons.org/licenses/by-nc/4.0/</a>
Link to Item	<a href="http://hdl.handle.net/2436/620636">http://hdl.handle.net/2436/620636</a>

# The Influence of Hormonal Contraception on Vitamin D Supplementation on Serum 25(OH)D<sub>3</sub> Status in Premenopausal Women: A Prospective Double-Blind Placebo Random Controlled Trial

Matthew Alexander Wyon<sup>a, e</sup>, Roger Wolman<sup>b</sup>, Alan M. Nevill<sup>a</sup>, Alison Barber<sup>a</sup>,  
Marcia Edwards<sup>c</sup>, Belinda Bowd<sup>a</sup>, Frances Clarke<sup>d</sup>, Janine Bryant<sup>d</sup>,  
Ross Cloak<sup>a</sup>

## Abstract

**Background:** A number of cross-sectional studies have highlighted a potential benefit of estrogen-containing contraception on serum 25-hydroxyvitamin D (25(OH)D) levels. The purpose of the present prospective study was to determine whether oral vitamin D<sub>3</sub> supplementation significantly increases serum 25(OH)D more for women taking the estrogen-containing oral contraception than those not taking this medication.

**Methods:** Thirty-eight premenopausal adult females aged 18 - 45 years old were recruited from a university campus; exclusion criteria included those presently taking vitamin D supplementation, those who stopped or started taking oral contraception in last 6 months and those taking any other form of contraception. A prospective double-blind placebo design was implemented; the dependent variable was serum 25(OH)D and the independent variables were using or not using oral estrogen-containing contraception, and vitamin D<sub>3</sub> or placebo supplementation. Participants were tested 4 weeks apart, and blood samples were collected using a capillary blood spot sample method and analyzed by liquid chromatography-tandem mass spectrometry. An independent technician prepared the identical supplement bottles with either 100 placebo pills or 100 active vitamin D<sub>3</sub> pills (1,000 IU per pill) and participants randomly selected a supplement bottle.

**Results:** Baseline measurements of 25(OH)D were non-significantly 11% higher in those taking estrogen. ANOVA results revealed a sig-

nificant two-way interaction between supplementation group (treatment vs. placebo) and treatment period (before vs. after) ( $P < 0.001$ ), demonstrating a substantial rise in serum 25(OH)D for the treatment group compared with the placebo group. The results also identified a three-way interaction ( $P = 0.014$ ) on serum 25(OH)D between the three independent variables, with the vitamin D oral contraception group having significantly greater serum 25(OH)D increases (from 45.9 to 98.3 nmol/L) compared with those not taking oral contraception (44.2 - 69.6 nmol/L) ( $P = 0.019$ ).

**Conclusions:** The estrogen-containing oral contraception increases serum 25(OH)D in premenopausal women with a magnified effect in those taking vitamin D supplementation. Future studies need to examine the relationship between estrogen, vitamin D supplementation, serum 25(OH)D, 1,25(OH)D, parathyroid hormone and other markers of bone metabolisms.

**Keywords:** Premenopausal; Vitamin D; Estrogen; Contraception; Female; Serum 25(OH)D; Intervention

## Introduction

There is a high prevalence of vitamin D insufficiency and deficiency across the world [1-3] and living near the equator, with its associated exposure to sunlight, does not necessary equate to adequate serum vitamin D status [4]. Deficiency is associated with a range of medical disorders including metabolic bone disease [3, 5], type 2 diabetes [2], cancer and impaired immune function [6, 7]. It is also associated with muscle weakness, which has been linked to the increased risk of falls in the elderly [5, 8] and to impaired athletic performance [9-11].

It is therefore important to be able to accurately measure vitamin D to identify those people with deficiency. Serum 25-hydroxyvitamin D (25(OH)D) is the most stable circulating biomarker of vitamin D and is therefore measured in routine clinical practice. It exists in two major forms: 25(OH)D<sub>2</sub> and 25(OH)D<sub>3</sub>. The former is mainly plant-derived, whilst 25(OH)D<sub>3</sub> is obtained mainly from sunlight exposure and diet (dairy

Manuscript submitted August 4, 2017, accepted August 15, 2017

<sup>a</sup>Research Centre for Sport Exercise and Performance, University of Wolverhampton, Walsall, UK

<sup>b</sup>Department of Rheumatology and Sport and Exercise Medicine, Royal National Orthopaedic Hospital, Stanmore, UK

<sup>c</sup>Institute of Health, University of Wolverhampton, Walsall, UK

<sup>d</sup>School of Performing Arts, University of Wolverhampton, Walsall, UK

<sup>e</sup>Corresponding Author: Matthew Wyon, Research Centre for Sport Exercise and Performance, University of Wolverhampton, Walsall, WS1 3BD, UK. Email: m.wyon@wlv.ac.uk

doi: <https://doi.org/10.14740/jem441w>

**Table 1.** Participants' Descriptive Data

Group		Age (years)	Height (cm)	Weight (kg)	BMI
Contraception	Supplementation				
Yes	Vit D (n = 10)	24 ± 5.99	165.56 ± 9.74	67.0 ± 9.23	20.18 ± 1.92
	Placebo (n = 9)	30 ± 11.62	161.50 ± 9.03	64.3 ± 5.63	19.88 ± 1.39
No	Vit D (n = 10)	22 ± 4.55	169.22 ± 6.41	70.6 ± 12.33	20.77 ± 2.97
	Placebo (n = 9)	27 ± 10.71	166.33 ± 6.32	68.8 ± 9.28	20.66 ± 2.61

products and fatty fish). Serum 25(OH)D<sub>3</sub> has a higher biopotency than 25(OH)D<sub>2</sub> and therefore food fortification or supplements are mainly given as vitamin D<sub>3</sub> [12]. Serum 1,25-dihydroxyvitamin D is the active form of the vitamin but this is more difficult to measure.

Serum levels of 25(OH)D are influenced by a range of factors. Palaniswamy et al [13] reported that gender, season, diet, obesity and physical activity were all associated with low 25(OH)D concentrations. A number of cross-sectional studies have reported that estrogen-containing hormonal contraception enhances serum 25(OH)D levels in women [10, 13, 14]. Harris and Dawson-Hughes [14] reported that women on contraception had 39% higher serum 25(OH)D levels compared with non-users. They also noted that five participants who ceased taking oral contraception during the following year had decreases in their 25(OH)D levels. The underlying mechanism for this is unclear at present, although there are at least two possible mechanisms: estrogen increases vitamin D 25-hydroxylase activity in the liver [15], and it causes an increased circulating concentration of vitamin D binding protein [16]. Both these mechanisms could increase 25(OH)D levels but it is still unclear whether this has a physiological effect on bone metabolism.

The previous studies have been observational in nature. The purpose of the present prospective study was to determine whether oral vitamin D<sub>3</sub> supplementation significantly increases serum 25(OH)D more for women taking the estrogen-containing oral contraception than those not taking this medication.

## Materials and Methods

### Participants

Premenopausal adult females aged between 18 and 45 years old were recruited from a university campus (latitude 52.58 °N) during the month of March. Participant exclusion criteria included those presently taking vitamin D supplementation, those who stopped or started taking oral contraception in last 6 months and those taking any other form of contraception, e.g. implant. Inclusion criteria for participants taking oral contraception required a minimum of 6 months using estrogen-containing contraception. Required sample size was calculated, based upon data provided by Harris and Dawson-Hughes [14], as six participants per group (oral contraception + vit D supplementation; oral contraception + placebo; no contraception

+ vit D supplementation; no contraception + placebo) based upon magnitude of effect at 90% confidence intervals [17]. To account for blinded nature of the study and potential dropout, it was decided to recruit 60 participants. Fifty-eight female participants volunteered for the study and 20 dropped out (34%) during its course (Table 1).

### Experimental design

A prospective double-blind placebo design was implemented. The dependent variable was serum 25(OH)D and the independent variables were using or not using oral estrogen-containing contraception, and vitamin D<sub>3</sub> or placebo supplementation.

An independent technician prepared the identical supplement bottles with either 100 placebo pills or 100 active vitamin D<sub>3</sub> pills (1,000 IU per pill) that had uniform appearance. Using an online random number generator ([www.sealedenvelope.com](http://www.sealedenvelope.com)), all bottles were allocated a number using a 1:1 allocation. Participants, outcome assessors, and data analysts were kept blinded to the allocation, and the independent researcher unblinded the trial post data analysis. The entire protocol was approved by a university Ethics Committee.

### Protocol

Participants were tested 4 weeks apart. Recorded sunlight during this period was 124.8 h. Each participant completed informed consent before anthropometric measurements and blood collection. Standing height was measured to the nearest 0.1 cm using a Seca stadiometer (Hamburg, Germany), with the participants in bare feet and their heads in Frankfurt horizontal plane. Total body mass was measured to the nearest 0.5 kg with a Seca beam balance 710 (Hamburg, Germany), and participants wore minimal clothing. Using a capillary blood spot sample method, the tester used a single use lancet on the participant's selected finger and the first show of blood was wiped away. Four blood spots were collected on the blood collection card (City Assays, Birmingham, UK) making sure the spots were of sufficient size and had soaked through the paper. The card was then sealed before being sent to an independent laboratory for analysis (City Assays, Pathology Department Sandwell and West Birmingham Hospital NHS Trust, UK). Each participant was asked to select a number from a number grid 1-60. The relevant supplement sample was then issued to the participant with the instructions to complete the

**Table 2.** Descriptive Pre and Post Serum 25(OH)D Data

Group		Serum 25(OH)D	
Contraception	Supplementation	Pre (nmol/L)	Post (nmol/L)
Yes	Vit D (n = 10)	52.3 ± 33.65	98.3 ± 37.02* <sup>†</sup>
	Placebo (n = 9)	43.5 ± 19.00	45.9 ± 2.06
No	Vit D (n = 10)	47.5 ± 24.85	69.6 ± 24.21*
	Placebo (n = 9)	39.0 ± 14.86	44.2 ± 15.79

\*Significant change pre and post ( $P < 0.001$ ). <sup>†</sup>Significant greater increase compared to other groups ( $P = 0.019$ ).

supplementation within 3 days. Blood data collection occurred again 4 weeks later. After the study, the lead author contacted all participants to feedback their individual results and provide participants who had been on the placebo trial with vitamin D<sub>3</sub> supplementation.

### Blood analysis

A liquid chromatography-tandem mass spectrometry (LC-MS/MS) method was applied to dried blood spot samples, utilizing blood spot calibrators. The method is standardized against conventional 25-hydroxyvitamin D<sub>3</sub> and D<sub>2</sub> LC-MS/MS service for serum. Blood spot results showed good comparability to serum/plasma results ( $r^2 = 0.98$ ; intra-assay variation  $< 10\%$ ; inter-assay variation  $< 11\%$ ). The City Assays laboratory participates in the DEQAS external quality assurance scheme.

### Data analysis

A three-way repeated-measures ANOVA was used to detect changes in serum 25(OH)D due to the independent variables, either using or not using oral contraception, and vitamin D<sub>3</sub> or placebo supplementation, before and after being given the vitamin D<sub>3</sub> or placebo supplementation. The repeated-measures factors were: oral contraception (yes or no) and vitamin D<sub>3</sub> supplementation (vitamin D<sub>3</sub> or placebo) as the two between subject factors, and the treatment period (before or after being given the supplementation) as the third within-subject factor.

### Results

There was no difference in the age and anthropometric measurements between the groups (Table 1). The baseline measurements of 25(OH)D were 11% higher in those taking estrogen ( $48.1 \pm 27.33$  vs.  $43.5 \pm 20.64$  nmol/L), but this did not reach statistical significance.

The ANOVA results revealed a significant two-way interaction between supplementation group (treatment vs. placebo) and treatment period (before vs. after) ( $P < 0.001$ ), demonstrating a substantial rise in serum 25(OH)D for the treatment group compared with the placebo group, following the period of supplementation (Table 2). However, the ANOVA results also identified a three-way interaction ( $P = 0.014$ ) on serum


25(OH)D between the three independent variables (Fig. 1).

Clearly, the benefits of being given vitamin D can be seen in both groups (Fig. 1, as described in the two-way interaction above), but the benefit of being given vitamin D is significantly greater in those taking oral contraception compared with those not taking oral contraception ( $P = 0.019$ ). The take-up of serum 25(OH)D by those taking oral contraception is over 52 nmol/L (from 45.9 to 98.3 nmol/L) compared with only 25 nmol/L (44.2 - 69.6 nmol/L) for those not on oral contraception.

### Discussion

The present prospective study showed that both groups on vitamin D<sub>3</sub> supplementation increased serum 25(OH)D levels but the group that was also on estrogen-containing oral contraception reported a further 100% increase compared to those who were not taking this medication. This study supports the observations of previous cross-sectional and longitudinal observation studies that oral contraception has a positive effect on serum 25(OH)D levels [10, 14]. Our study also showed those taking the estrogen-containing oral contraceptive had 11% higher baseline 25(OH)D but this did not reach statistical significance. Studies by Harris and Dawson-Hughes [14] and Harmon et al [18] have both shown significant larger increases in 25(OH)D for those on the oral contraceptive of 20-39%. It is difficult to provide an explanation for these differences, but all three studies demonstrate that 25(OH)D serum levels increase in response to the estrogen-containing oral contraceptive. Our study also shows that this effect is enhanced when taking high-dose vitamin D supplementation.

Two mechanisms have been proposed for the increase in 25(OH)D. Nelson et al [15] suggested estrogen-containing oral contraception increased vitamin D 25-hydroxylase activity in the liver, whilst Moller et al [16] proposed the increase was due to an increase in circulating concentration of vitamin D binding protein (VDBP). However, it is currently unclear whether this biochemical response of 25(OH)D occurs in isolation or whether it is associated with an increase in vitamin D activity on bone metabolism. It is possible that, if the increase in 25(OH)D is due to an increase in the circulating VDBP with an increase in bound vitamin D, this may not be accompanied by an increase in bio-activity. This may be of some concern as we use 25(OH)D as a clinical marker of vitamin D status. An increase in this marker due to estrogen supplementation may mask a potential deficiency in vitamin D bio-activity.



**Figure 1.** Changes in serum 25(OH)D. \*Significant change pre and post ( $P < 0.001$ ). †Significant greater increase compared to other groups ( $P = 0.019$ ).

Future prospective studies should therefore assess whether the increase in 25(OH)D is associated with parallel changes in serum 1,25(OH)D, parathyroid hormone (PTH) and other markers of bone metabolism. Wolman et al [10] noted a significant decrease in PTH for those on oral contraception that was associated with a significant increase in serum 25(OH)D above the levels reported for those not taking oral contraception. The study was limited by low numbers of participants on contraception [5] and therefore the results cannot be generalized further.

The observed vitamin D response to the estrogen-containing oral contraceptive raises the question as to whether there is a vitamin D response to other situations where there is a change in estrogen. In studies on postmenopausal women taking hormone replacement therapy (HRT), one reported a significant increase in vitamin D binding protein [19], whilst another showed no increased benefit on bone mineral content [20]. Heikkinen et al [21] reported a 38.2% increase in serum 25(OH)D for participants taking HRT and vitamin D<sub>3</sub>, but this was only a slightly greater increase than the vitamin D<sub>3</sub> only group (33.5%). It is also possible that vitamin D levels may change in response to those going through the menarche and the menopause. Further studies are needed to address these questions.

In conclusion, the estrogen-containing oral contraception

increases serum 25(OH)D in premenopausal women with a magnified effect in those taking vitamin D supplementation. Future studies need to examine the relationship between estrogen, vitamin D supplementation, serum 25(OH)D, 1,25(OH)D, PTH and other markers of bone metabolism.

## Acknowledgments

We acknowledge SunVit for supply of vitamin D<sub>3</sub> supplementation.

## Conflicts of Interest

The authors have no conflicts of interest.

## Author Contributions

M. A. Wyon and R. Wolman: study concept, study design, and write up. A. M. Nevill: statistical analysis. A. Barber, B. Bowd, F. Clarke, and J. Bryant: participant recruitment and manuscript review. R. Cloak: participant randomization and manuscript review.

## References

1. Dobnig H. A review of the health consequences of the vitamin D deficiency pandemic. *J Neurol Sci.* 2011;311(1-2):15-18.
2. Heaney RP. Vitamin D in health and disease. *Clin J Am Soc Nephrol.* 2008;3(5):1535-1541.
3. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr.* 2008;87(4):1080S-1086S.
4. Hilger J, Friedel A, Herr R, Rausch T, Roos F, Wahl DA, Pierroz DD, et al. A systematic review of vitamin D status in populations worldwide. *Br J Nutr.* 2014;111(1):23-45.
5. Jesudason D, Need AG, Horowitz M, O'Loughlin PD, Morris HA, Nordin BE. Relationship between serum 25-hydroxyvitamin D and bone resorption markers in vitamin D insufficiency. *Bone.* 2002;31(5):626-630.
6. Antico A, Tampoia M, Tozzoli R, Bizzaro N. Can supplementation with vitamin D reduce the risk or modify the course of autoimmune diseases? A systematic review of the literature. *Autoimmun Rev.* 2012;12(2):127-136.
7. Devaux S, Castela A, Archier E, Gallini A, Joly P, Misery L, Aractingi S, et al. Topical vitamin D analogues alone or in association with topical steroids for psoriasis: a systematic review. *J Eur Acad Dermatol Venereol.* 2012;26(Suppl 3):52-60.
8. Gerdhem P, Ringsberg KA, Obrant KJ, Akesson K. Association between 25-hydroxy vitamin D levels, physical activity, muscle strength and fractures in the prospective population-based OPRA Study of Elderly Women. *Osteoporos Int.* 2005;16(11):1425-1431.
9. Cannell JJ, Hollis BW, Sorenson MB, Taft TN, Anderson JJ. Athletic performance and vitamin D. *Med Sci Sports Exerc.* 2009;41(5):1102-1110.
10. Wolman R, Wyon MA, Koutedakis Y, Nevill AM, Eastell R, Allen N. Vitamin D status in professional ballet dancers: winter vs. summer. *J Sci Med Sport.* 2013;16(5):388-391.
11. Wyon MA, Wolman R, Nevill AM, Cloak R, Metsios GS, Gould D, Ingham A, et al. Acute effects of Vitamin D3 supplementation on muscle strength in judoka athletes: a randomized placebo-controlled, double-blind trial. *Clin J Sport Med.* 2016;26(4):279-284.
12. Heaney RP, Recker RR, Grote J, Horst RL, Armas LA. Vitamin D(3) is more potent than vitamin D(2) in humans. *J Clin Endocrinol Metab.* 2011;96(3):E447-452.
13. Palaniswamy S, Hypponen E, Williams DM, Jokelainen J, Lowry E, Keinanen-Kiukaanniemi S, Herzig KH, et al. Potential determinants of vitamin D in Finnish adults: a cross-sectional study from the Northern Finland birth cohort 1966. *BMJ Open.* 2017;7(3):e013161.
14. Harris SS, Dawson-Hughes B. The association of oral contraceptive use with plasma 25-hydroxyvitamin D levels. *J Am Coll Nutr.* 1998;17(3):282-284.
15. Nelson ML, Blum JM, Hollis BW, Rosen C, Sullivan SS. Supplements of 20 microg/d cholecalciferol optimized serum 25-hydroxyvitamin D concentrations in 80% of premenopausal women in winter. *J Nutr.* 2009;139(3):540-546.
16. Moller UK, Streym S, Jensen LT, Mosekilde L, Schoenmakers I, Nigdikar S, Rejnmark L. Increased plasma concentrations of vitamin D metabolites and vitamin D binding protein in women using hormonal contraceptives: a cross-sectional study. *Nutrients.* 2013;5(9):3470-3480.
17. Hopkins W. Estimating sample size for magnitude-based inferences. *Sportscience.* 2006;10:63-70.
18. Harmon QE, Umbach DM, Baird DD. Use of estrogen-containing contraception is associated with increased concentrations of 25-Hydroxy Vitamin D. *J Clin Endocrinol Metab.* 2016;101(9):3370-3377.
19. Rejnmark L, Lauridsen AL, Brot C, Vestergaard P, Heickendorff L, Nexø E, Mosekilde L. Vitamin D and its binding protein Gc: long-term variability in peri- and postmenopausal women with and without hormone replacement therapy. *Scand J Clin Lab Invest.* 2006;66(3):227-238.
20. Christiansen C, Christensen MS, Rodbro P, Hagen C, Transbol I. Effect of 1,25-dihydroxy-vitamin D3 in itself or combined with hormone treatment in preventing postmenopausal osteoporosis. *Eur J Clin Invest.* 1981;11(4):305-309.
21. Heikkinen A, Parviainen MT, Tuppurainen MT, Niskanen L, Komulainen MH, Saarikoski S. Effects of postmenopausal hormone replacement therapy with and without vitamin D3 on circulating levels of 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D. *Calcif Tissue Int.* 1998;62(1):26-30.