

Determinants of Fatigue after First-Ever Ischemic Stroke during Acute Phase

Item Type	Journal article
Authors	Chen, Ruoling;Wang, Jia-Ji;Wang, Pei-Xi;Wang, Shan-Shan
Citation	Wang S-S, Wang J-J, Wang P-X, Chen R (2014) Determinants of Fatigue after First-Ever Ischemic Stroke during Acute Phase. PLoS ONE 9(10): e110037. https://doi.org/10.1371/journal.pone.0110037
DOI	10.1371/journal.pone.0110037
Publisher	PLoS One
Journal	PloS one
Download date	2026-04-15 06:27:31
License	https://creativecommons.org/licenses/by/4.0/
Link to Item	http://hdl.handle.net/2436/620801

Correction

Correction: Determinants of Fatigue after First-Ever Ischemic Stroke during Acute Phase

The PLOS ONE Staff

The affiliation for the fourth author is incorrect. Ruoling Chen is not affiliated with #3 but with #4 Centre for Health and Social Care Improvement (CHSCI), University of Wolverhampton, Wolverhampton, United Kingdom.

Reference

1. Wang S-S, Wang J-J, Wang P-X, Chen R (2014) Determinants of Fatigue after First-Ever Ischemic Stroke during Acute Phase. PLoS ONE 9(10): e110037. doi:10.1371/journal.pone.0110037

Citation: The PLOS ONE Staff (2014) Correction: Determinants of Fatigue after First-Ever Ischemic Stroke during Acute Phase. PLoS ONE 9(12): e116461. doi:10.1371/journal.pone.0116461

Published: December 19, 2014

Copyright: © 2014 The PLOS ONE Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.