

The relationship between vitamin D levels, injury and muscle function in adolescent dancers

Item Type	Journal article
Authors	de Rezende Araújo, Iris Iasmine;Sampaio, Lucas Henrique Ferreira;Bittar, Adriano Jabur;da Silva Hamu, Tânia Cristina Dias;Wyon, Matthew;Formiga, Cibelle Kayenne Martins Roberto
Citation	De Rezende Araújo, I. I., Sampaio, L. H. F., Bittar, A. J., Da Silva Hamu, T. C. D., Wyon, M., Formiga, C. K. M. R. (2020) The relationship between vitamin D levels, injury and muscle function in adolescent dancers, International Journal of Sports Medicine, 41(6), pp. 360-364
DOI	10.1055/a-1087-2130
Publisher	Thieme
Journal	International Journal of Sports Medicine
Download date	2026-04-17 03:37:11
License	https://creativecommons.org/licenses/by-nc-nd/4.0/
Link to Item	http://hdl.handle.net/2436/622967

The relationship between vitamin D levels, injury and muscle function in adolescent dancers.

Iris Iasmine de Rezende Araújo, MSc¹; Lucas Henrique Ferreira Sampaio, PhD²; Adriano Jabur Bittar, PhD¹; Tânia Cristina Dias da Silva Hamu, PhD³; Matthew Wyon, PhD^{4,5}; Cibelle Kayenne Martins Roberto Formiga, PhD³

¹ State University of Goiás, Brazil

² Laboratory of Immunology, State University of Goiás, Brazil

³ Laboratory of Musculoskeletal Research, State University of Goiás, Brazil

⁴ Institute of Human Sciences, University of Wolverhampton, UK

⁵ National Institute of Dance Medicine and Science

Corresponding author

Prof Matthew Wyon

Institute of Human Sciences,

University of Wolverhampton,

Gorway Rd, Walsall

WS1 3BD, UK

Abstract

Vitamin D has been shown to benefit a diverse range of health functions including muscle function. The aim of the present study was to identify serum 25(OH)D₃ levels in a sample of adolescent dancers and compare them to muscle function and injury incidence. We incorporated a cross-sectional design to study 49 pre-professional male and female dancers (17±4.44yrs, 52.1±6.72kg, 1.63±0.07m) in full-time training in Brazil. Serum 25[OH]D₃ was analyzed by Enzyme-linked Immunosorbent Assay; quadriceps and hamstring peak torque and muscle fatigue were measured by isokinetic dynamometer at 60 and 300° s⁻¹. Injury type and location in the previous 6-months were determined by self-report questionnaire. Participants were categorized into 2 groups: normal or insufficient/deficient (>or<30 ng/ml 25[OH]D₃). Results indicated the normal serum 25[OH]D₃ group had significantly lower fatigue rates than the insufficient/deficient group (p<0.05) but not for the other muscle function parameters. Fifty-seven percent of participants reported at least 1 injury. The most common were sprains (33%) and tendinopathies (19%). Injured dancers had significantly lower peak torque at 60°/s. The link between serum 25[OH]D₃ and reduced muscle fatigue resistance has not been shown before, though the underlying mechanisms aren't apparent and the link between muscular strength and injury has been previously evidenced.

Keywords: Muscle Strength Dynamometer, fatigue, dance, serum 25(OH)D.

INTRODUCTION

Pre-professional dance students undertake 3-6 hours of dance training a day; this is generally technique focused but at different times of the year will also include rehearsals and performances [1, 2]. The training regimen is often perceived as exhaustive and intense leading to several health problems and a high injury prevalence [3, 4]. The injury prevalence has been attributed to the long training hours the high relative intensity, the impact from jumping and muscular imbalances [5, 6]. A lean and ectomorphic body is required for classical ballet professionals[7] which has been linked to poor nutritional practices[8].

Vitamin D has several effects on body systems[9] and has an influence on muscle function[10]. Vitamin D is linked to metabolism and protein synthesis in skeletal muscle cells[11]. Dancers have been shown to be vitamin D insufficient/deficient throughout the year, with only slight improvements during the summer months, potentially due to their long hours training indoors[12]. Supplementation has been shown to improve vitamin D levels with subsequent improvements in muscle function and a reduction in injury incidence[13, 14].

There are still few studies relating to classical ballet dancing and vitamin D levels in warm climate countries like Brazil, although studies on the general population in high sunlight countries have indicated similar deficiency incidence[15]. Considering that dancers are often involved in technical classes/rehearsals indoors for several hours a day and that such activities make them more susceptible to musculoskeletal injuries, we hypothesize that, in dancers, insufficient levels of vitamin D are related to a decrease in muscle function and a higher injury incidence. Therefore, the aim was to identify serum 25[OH]D₃, muscle function and injury incidence in adolescent vocational ballet dancers.

MATERIALS AND METHODS

The study, a cross-sectional design, was approved by the lead author's university Ethics and Research Committee[16]. Data collection took place in Goiania at latitude -16.6799, longitude -49.255 during the month of October; the previous three months mean sunlight was 11.5 hours (<http://www.inmet.gov.br/>). The inclusion criteria were as follows: male and female dancers, aged 13 years and older, who have been dancing for more than 5 years and who were duly enrolled in Centro de Educação Profissional em Artes Basileu França (CEPABF). Dancers who had been absent from training for 1 month, were under medical care or injured and younger than 13 years old were excluded. Fifty-eight dancers were initially recruited, 49 dancers presented for blood collection and a further 7 did not attend the muscle strength evaluation; therefore, the final sample of the present study was composed of 42 dancers (Table 1). All participants with the exception of two were white Caucasians.

Table 1: Participant information

	Age (yrs)	Body Mass (kg)	Stature (m)	BMI	Age started dancing (yrs)	Years dancing (yrs)	Dance exposure per week (hrs)
Males (n=5)	17 ±4.01	61.6 ±7.92	1.74 ±0.09	20.4 ±0.76	15 ±4.85	7 ±3.27	5 ±2.06
Females (n=37)	17 ±5.55	51.2 ±5.99	1.62 ±0.06	19.4 ±1.91	7 ±2.90	10 ±3.29	5 ±2.02

Procedures

The dancers' anthropometric and training history data were collected including age, sex, stature, body mass, age they started dancing and daily duration of training. Stature (cm) was measured to the nearest 0.1cm using a SECA 217 stadiometer (Germany) and body mass (kg) to the nearest 0.1kg with SECA 761 scales (Germany) wearing minimal clothing. Age and age they started training were documented as whole years. Daily training was recorded in 30 minute periods. A questionnaire also included questions regarding supplementation consumption (minerals or vitamins) and whether any of the females were taking contraceptive medication.

Participant's vitamin D levels were measured via blood sample. The blood collection was performed by ulnar venipuncture, in accordance with the recommendations of the Brazilian Society of Clinical Pathology. A 5-7 ml sample of whole blood was collected into tubes which did not contain anticoagulants. The blood was left to clot for 30 min at room temperature before being spun in a centrifuge at $1500 \times g$ for 10 min. A 0.5 ml of serum was aliquoted into 2 ml micro-tubes prior to being stored at -80°C . The thawed samples were later analyzed for serum 25(OH)D₃ by using the enzyme-linked immunosorbent assay (ELISA), through the 25-OH Vitamin D[®] kit (DRG, Springfield, NJ, USA), with interassay CV of 2.9%. All the collected blood material was analyzed in the Laboratory of Molecular Immunology at UEG Campus Laranjeiras.

Muscle function was evaluated by isokinetic dynamometry of the knee joint using a Biodex System 4 PRO isokinetic dynamometer (Biodex Medical Systems, Shirley, NY, USA). The isokinetic dynamometer was set up with 5° recline of the chairback from the vertical position and the participant positioned on the chair so that the popliteal fossa of the knee was 5 cm away from the chair edge and the arm was positioned and secured by a belt 5 mm above the lateral malleolus. The range of motion was limited to between 10° and 110° of knee flexion. The participant was stabilized by 4 belts; two diagonals to the thorax, one in the lower abdominal region (in the transverse axis) and one in the lateral femoral condyle region of the femur bone of the evaluated limb (on the transverse axis). Participants carried out a moderate intensity cycle ergometer warm up for 5 minutes prior to a series of self-directed dynamic stretches. The test consisted of 3 sets of 5 repetitions for knee concentric flexion/extension at angular velocities of 60°s^{-1} and 300°s^{-1} (12,13); the velocity order was randomly assigned for each participant using a sealed envelope method.

Injury history was assessed by self-report questionnaire. Participants were asked to recall injuries over the last 6-months that included anatomical location, injury type (dislocation or subluxation, sprain, fracture, contusion and tendinopathy) and time off from dance activities.

Data Analysis

Vitamin D insufficiency was determined by circulating serum 25(OH)D₃ levels of 10–30 ng/ml (25–75 nmol/l) and deficiency as a level below 10 ng/ml (25 nmol/l)[17], participants were grouped according to having either normal levels (>30 ng/ml) or insufficient/deficient levels (<30 ng/ml). Dancers were further grouped according to their injury recall status into either injury or no injury. Fatigue index was calculated from the isokinetic data using the individual's greatest and lowest score over the 15 repetitions for both flexion (hamstrings) and extension (quadriceps). Peak torque was recorded from the isokinetic dynamometer.

Statistical analysis

Descriptive statistics were calculated for all the muscle function and injury data. Multiple Analysis of Variance tests were used to examine the effect of sex, vitamin D group, movement (extension/flexion), leg (left/right) on peak torque and fatigue index. Skin tone was not used as a covariate as only two participants were not white Caucasians. Chi-square tests were used to examine injury data between normal and insufficient/deficient vitamin D groups and vitamin D levels, and muscle function between injury and no injury groups. The Statistical Package for Social Sciences (SPSS) 23.0 was used for all analyses and significance set at $p \leq 0.05$.

RESULTS

None of the participants reported taking mineral or vitamin supplementation and no females reported taking contraceptive medication. Main effects were observed for Sex ($F=5.673$, $p<0.001$), Leg ($F=9.997$, $p<0.001$), Movement ($F=46.18$, $p<0.001$), Sex*Vitamin D group ($F=4.989$, $p<0.001$), Leg* Movement ($F=9.896$, $p<0.001$).

Table 2: Descriptive data (mean±sd) for serum 25(OH)D₃, peak torque and fatigue index according to gender, vitamin D group and leg

	Normal				Insufficient/Deficient			
	Male		Female		Male		Female	
Leg	Right	Left	Right	Left	Right	Left	Right	Left
Serum 25(OH)D ₃ (ng/ml)	48.8 ±7.49		43.3 ±8.74		5.9 ±0.66		17.8 ±7.92	
Peak Torque 60° s ⁻¹								
Extension (n.m)	124.2 ±46.67	136.3 ±49.24	130.1 ±21.92	122.3 ±25.79	159.2 ±16.62	138.8 ±37.83	134. ±34.83	119.4 ±36.47
Flexion (n.m)	72.3 ±14.31	70.4 ±46.39	56.6 ±13.37	57.5 ±14.31	65.7 ±10.75	69.4 ±8.42	62.9 ±25.76	58.3 ±20.68
300° s ⁻¹								
Extension (n.m)	86.6 ±45.11	86.4 ±39.69	69.4 ±11.39	66.5 ±12.74	80.1 ±13.72	65.3 ±21.85	64.9 ±19.56	63.0 ±19.68
Flexion (n.m)	59.4 ±31.79	59.0 ±33.13	47.3 ±9.28	48.3 ±9.87	59.4 ±31.79	54.5 ±4.38	46.1 ±14.89	47.6 ±15.74
Fatigue index 60° s ⁻¹								
Extension (%)	13.3 ±3.16	18.9 ±10.54	20.3 ±11.09	21.2 ±11.93	14.6 ±15.49	-6.9 ±52.12	17.6 ±5.40	19.4 ±13.89
Flexion (%)	10.7 ±15.62	18.7 ±5.75	16.5 ±15.49	19.5 ±10.28	-16.4 ±40.31	10.9 ±20.22	3.9 ±27.75	18.0 ±8.13
300° s ⁻¹								
Extension (%)	22.0 ±7.16	25.7 ±12.54	27.6 ±13.69	28.7 ±11.99	10.3 ±42.85	-0.7 ±52.04	19.5 ±16.98	21.0 ±15.64
Flexion (%)	28.7 ±12.78	68.0 ±7.25	27.9 ±14.17	68.4 ±9.77	9.7 ±56.56	69.9 ±4.67	8.8 ±26.88	39.7 ±37.2

Vitamin D

According to the parameters adopted by the Endocrine Society, 26 dancers were classified as having normal 25(OH)D₃ levels (43.5 ±9.09 ng/ml); 4 were insufficient (25.4 ±2.79 ng/ml) and 12 were deficient (12.2 ±6.14 ng/ml); these were subsequently put into 2 groups, normal and insufficient/deficient (Table 2). Between-subject effects reported a Sex*Group interaction ($F_{1,152}=17.653$; $p<0.001$) with males in the insufficient/deficient group having significantly lower serum 25(OH)D₃ than females; there was no difference between the sexes in the normal group. There were significant between-subject effects for the vitamin D groups for fatigue index at 60°s⁻¹ ($F_{1,152}=7.319$; $p<0.01$) and 300°s⁻¹ ($F_{1,152}=7.440$; $p<0.01$); in both instances the fatigue index was greater in the normal vitamin D group.

Muscle function

Irrespective of vitamin D group the following interactions were observed. Between-subject sex effects reported significantly greater extension than flexion peak torques at 60°s⁻¹ ($F_{1,152}=119.991$; $p<0.001$) and 300°s⁻¹ ($F_{1,152}=27.201$; $p<0.001$). A sex effect was also noted for peak torque at 300°s⁻¹ with males having significantly higher scores ($F_{1,152}=6.334$; $p<0.05$). Females reported a significantly higher fatigue index at 60°s⁻¹ ($F_{1,152}=9.058$; $p<0.01$) than their male counterparts. The left leg also had a significantly higher fatigue index than the right leg ($F_{1,152}=39.375$; $p<0.001$). Flexion fatigue index was significantly greater than extension data at 300°s⁻¹ ($F_{1,152}=38.479$; $p<0.001$)

Table 3: Incidence of injury types for gender, vitamin D group and anatomical site

Vit D status		Dislocation	Sprain	Fracture*	Contusion*	Tendonitis*	Total
Normal	Male	0	0	1	0	0	1
	Female	1	8	3	0	5	17
Insufficient/ Deficient	Male	1	1	0	0	0	2
	Female	4	5	4	3	4	20
Total		6	14	8	3	9	40
Site							
	Toe		2				2
	Foot	3	4	2		1	9
	Ankle		2			3	5
	Calf				1		1
	Knee	1	3		1	3	8
	Thigh	1					1
	Leg		1				1
	Groin					1	1
	Coccyx	1					1
	Hand			1			1
	Finger		2	1			3
	Face			1			1

*missing data for site

Injury

Fifty-seven percent of participants reported an injury during the study period. The most common types were sprain (33%), tendinopathy (19%) and fracture (12%) (Table 3) with the majority of injuries were to the lower limbs. Neither sex nor vitamin D status had a significant effect on injury incidence. Injured dancers had significantly lower extension and flexion peak torque at 60°s^{-1} ($p < 0.05$).

DISCUSSION

The present study aimed to identify levels of vitamin D₃ and to verify the impact of these levels on the parameters of thigh muscle strength and injury status of adolescent classic ballet dance professionals. The data collection occurred during the dry season in Goiânia, Brazil that begins in April and extends until the first half of October that is considered favorable for vitamin D production through solar radiation[18]. Despite the potential solar benefit 43% of the dancers were observed to be insufficient or deficient in serum 25(OH)D₃; this could be due to the length of time spent indoors[14], the use of sunscreen[19] and a diet poor in vitamin D sources[15]. This is similar to the data reported by Wolman et al[12] that the potential of summer sun exposure only has a negligible effect on dancers' serum 25(OH)D₃ as they are indoors between 10am to 6pm and also use sunscreen. The increased prevalence of deficiency in males was also observed by Looker et al[20], though they could not provide any rationale for the sex difference.

Unlike a number of previous studies on dancers[12-14, 21, 22] this study demonstrated no positive link between muscle function and serum 25(OH)D₃ levels. In fact, fatigue resistance (a lower fatigue percentage) was associated with the insufficient/deficient vitamin D group. This result could be due to the increased variance in the male fatigue data, which is affected by small participant numbers, but the phenomenon remains when the male participants are excluded.

Previous studies have negatively linked muscle strength with injury incidence[23-25], with stronger dancers reporting fewer injuries[26]. This phenomenon was also observed in the current study with weaker dancers that produced less torque being significantly more likely to be injured (peak torque at 60°s^{-1}). Muscle function profiles are similar to those previously reported in that extension peak torque and fatigue index are significantly greater than the flexion data, and peak torque at 60°s^{-1} were greater than at 300°s^{-1} [26-28]. The lack of sex differences in peak torque at 60°s^{-1} is unusual in our study; torque production by males is expected to be greater than females. A possible reason for this observation is the age range in the current study (13-21 years) that will encompass differing ages for adolescent pubertal development and its associated hormonal changes[29].

Injury incidence is lower than previously reported[30, 31] but is probably due to a reduced reporting period in the current study. The self-reported injury sites are similar to that previously reported with a focus on the lower limbs[31], particularly the knee and ankle. The main two types of injury were sprains and tendinopathy which amongst a number of mechanisms also have links to muscle underperformance or imbalance, through either weakness or fatigue [32, 33].

Limitations in the current study revolve around the sex imbalance in participants, the use of self-report injury history and the study's cross-sectional nature. Focused recruitment of a larger sex balanced population with equal numbers across the age range, taking into account pubertal development would provide a stronger foundation for a study. The use of prospective injury recording, either through a medical team or self-report[34] would potentially strengthen the injury data. Longitudinal data would provide an insight into vitamin D and muscle function fluctuations over the academic year and holiday periods.

CONCLUSION

The current study has provided a novel insight into vitamin D levels, muscle function and self-report injuries in elite adolescent Brazilian ballet dancers. Thirty-eight percent of the cohort were classified as either insufficient or deficient despite data collection occurring during the dry season that had approximately 12 hours of sun a day. The dancers insufficient/deficient in vitamin D unusually reported greater resistance to muscular fatigue. In the current cohort, there was no link between serum 25(OH)D levels and muscular strength. A negative association between muscle strength and injury incidence highlights the need for supplemental training to rectify this risk.

REFERENCES

1. Ekegren CL, Quested R, Brodrick A. Injuries in pre-professional ballet dancers: incidence, characteristics and consequences. *J Sci Med Sport* 2014; 17: 271-275
2. Caine D, Bergeron G, Goodwin BJ et al. A survey of injuries affecting pre-professional ballet dancers. *J Dance Med Sci* 2016; 20: 115-126
3. Vaisman N, Voet H, Akivis A et al. Weight perception of adolescent dancing school students. *Arch Pediatr Adolesc Med* 1996; 150: 187-190
4. Braisted JR, Mellin L, Gong EJ et al. The adolescent ballet dancer: Nutritional practices and characteristics associated with anorexia nervosa. *J Adolesc Health Care* 1985; 6: 365-371
5. Gamboa JM, Roberts LA, Maring J et al. Injury patterns in elite preprofessional ballet dancers and the utility of screening programs to identify risk characteristics. *J Orthop Sports Phys Ther* 2008; 38: 126-136
6. Luke AC, Kinney SA, D Hemecourt PA et al. Determinants of injuries in young dancers. *Med Probl Perform Artists* 2002; 17: 105-112
7. de Moura UIS, Mendes LR, de Oliveira Silva IP et al. Consumo alimentar, perfil antropométrico e imagem corporal de bailarinas clássicas do Vale do São Francisco. *Revista Brasileira de Nutrição Esportiva* 2015; 9: 237-246
8. Wyon M, Hutchings K, Wells A et al. Body Mass Index, Nutritional Knowledge, and Eating Behaviors in Elite Student and Professional Ballet Dancers. *Clin J Sport Med* 2014; 24: 390-396

9. Dobnig H. A review of the health consequences of the vitamin D deficiency pandemic. *J Neurol Sci* 2011. 15-18
10. Beaudart C, Buckinx F, Rabenda V et al. The effects of vitamin D on skeletal muscle strength, muscle mass, and muscle power: a systematic review and meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab* 2014; 99: 4336-4345. doi:10.1210/jc.2014-1742
11. Pedrosa-Castro MAC, Lazaretti-Castro M. Papel da vitamina D na função neuromuscular. *Arquivos Brasileiros de Endocrinologia & Metabologia* 2005.
12. Wolman R, Wyon M, Allen N et al. The vitamin D status of professional dancers in the winter and in the summer. *J Sci Med Sport* 2013; 16: 388-391. doi:dx.doi.org/10.1016/j.jsams.2012.12.010
13. Wyon M, Koutedakis Y, Wolman R et al. The influence of winter vitamin D supplementation on muscle function and injury occurrence in elite ballet dancers: A controlled study. *J Sci Med Sport* 2014; 17: 8-12. doi:10.1016/j.jsams.2013.03.007
14. Wyon M, Wolman R, Kolokythas N et al. The effect of Vitamin D supplementation in elite adolescent dancers on muscle function and injury incidence: a randomised double-blind study. *Inter J Sports Physiol Perf* 2018. doi:10.1123/ijsp.2018-0084. doi:10.1123/ijsp.2018-0084
15. Wahl D, Cooper C, Ebeling P et al. A global representation of vitamin D status in healthy populations *Archives of Osteoporosis* 2012. doi:10.1007/s11657-012-0093-0. doi:10.1007/s11657-012-0093-0
16. Harriss D, MacSween A, Atkinson G. Ethical Standards in Sport and Exercise Science Research: 2020 Update. *Int J Sports Med* 2019.
17. Ross A, Manson J, Abrams S et al. The 2011 Report on Dietary Reference Intakes for Calcium and Vitamin D from the Institute of Medicine: What Clinicians Need to Know *J Clin Endocrinol Metab* 2011; 96: 53-58
18. Maeda SS, Kunii IS, Hayashi L et al. The effect of sun exposure on 25-hydroxyvitamin D concentrations in young healthy subjects living in the city of São Paulo, Brazil. *Braz J Med Biol Res* 2007; 40: 1653-1659
19. Faurischou A, Beyer D, Schmedes A et al. The relation between sunscreen layer thickness and vitamin D production after ultraviolet B exposure: a randomized clinical trial. *Br J Dermatol* 2012; 167: 391-395
20. Looker AC, Johnson CL, Lacher DA et al. Vitamin D status: United states, 2001–2006. *NCHS data brief* 2011; 59: 1-8
21. Constantini N, Arieli R, Chodick G et al. High prevalence of vitamin D insufficiency in athletes and dancers. *Clin J Sport Med* 2010; 20: 368-371. doi:10.1097/JSM.0b013e3181f207f2
22. Ducher G, Kukuljan S, Hill B et al. Vitamin D status and musculoskeletal health in adolescent male ballet dancers: a pilot study. *Journal of Dance Medicine and Science* 2011; 15: 99-107
23. Khayambashi K, Ghoddosi N, Straub RK et al. Hip muscle strength predicts noncontact anterior cruciate ligament injury in male and female athletes: a prospective study. *Am J Sports Med* 2016; 44: 355-361. doi:10.1177/0363546515616237

24. Lehance C, Binet J, Bury T et al. Muscular strength, functional performances and injury risk in professional and junior elite soccer players. *Scand J Med Sci Sports* 2009; 19: 243-251
25. Nadler S, Malanga G, DePrince M et al. The relationship between lower extremity injury, low back pain, and hip muscle strength in male and female collegiate athletes. *Clin J Sport Med* 2000; 10: 89-97
26. Koutedakis Y, Khalouha M, Pacy P et al. Thigh peak torques and lower-body injuries in dancers. *J Dance Med Sci* 1997; 1: 12-15
27. Koutedakis Y, Agrawal A, Sharp NCC. Isokinetic characteristics of knee flexors and extensors in male dancers, Olympic oarsmen, Olympic bobsleighters, and non-athletes. *Journal of Dance Medicine and Science* 1999; 2: 63-67
28. Koutedakis Y, Frischknecht R, Murthy M. Knee flexion to extension peak torque ratios and low-back injuries in highly active individuals. *Int J Sports Med* 1997; 18: 290-295. doi:10.1055/s-2007-972636
29. Quatman CE, Ford KR, Myer GD et al. Maturation leads to gender differences in landing force and vertical jump performance: a longitudinal study. *The American journal of sports medicine* 2006; 34: 806-813
30. Kenny SJ, Whittaker JL, Emery CA. Risk factors for musculoskeletal injury in preprofessional dancers: a systematic review. *Br J Sports Med* 2016; 50: 997-1003
31. Laws H. *Fit to Dance 2 - Report of the second national inquiry into dancers' health and injury in the UK*. London: Newgate Press; 2005
32. Niemuth PE, Johnson RJ, Myers MJ et al. Hip muscle weakness and overuse injuries in recreational runners. *Clin J Sport Med* 2005; 15: 14-21
33. Ribeiro F, Oliveira J. Efeito da fadiga muscular local na propriocepção do joelho. *Fisioterapia em Movimento* 2017; 21
34. Kenny SJ, Palacios-Derflinger L, Whittaker JL et al. The influence of injury definition on injury burden in preprofessional ballet and contemporary dancers. *J Orthop Sports Phys Ther* 2018; 48: 185-193. doi:10.2519/jospt.2018.7542