

Chemical cues of identity and reproductive status in Japanese macaques

Item Type	Journal article
Authors	Rigaille, Lucie;Vaglio, Stefano;Setchell, Joanna M;Suda-Hashimoto, Naoko;Furuichi, Takeshi;Garcia, Cécile
Citation	Rigaille, L., Vaglio, S., Setchell, J. M., Suda-Hashimoto, N., et al (2022) Chemical cues of identity and reproductive status in Japanese macaques, American Journal of Primatology, 84(8), e23411.
DOI	10.1002/ajp.23411
Publisher	Wiley-Blackwell
Journal	American Journal of Primatology
Download date	2026-04-18 07:22:56
License	https://authorservices.wiley.com/author-resources/Journal-Authors/licensing/self-archiving.html
Link to Item	http://hdl.handle.net/2436/624781

1 **Chemical cues of identity and reproductive status in Japanese macaques**

2

3 Lucie Rigaill^{1,2}, Stefano Vaglio³, Joanna M. Setchell⁴, Naoko Suda-Hashimoto², Takeshi

4 Furuichi², Cécile Garcia⁵

5 ¹ EthoS (Éthologie animale et humaine), CNRS, Université de Rennes 1, Normandie Université,

6 Rennes, France

7 ² Center for the Evolutionary Origins of Human Behavior, Kyoto University, Inuyama, Japan

8 ³ School of Sciences, University of Wolverhampton, UK

9 ⁴ Department of Anthropology & Behaviour, Ecology and Evolution Research Centre, Durham

10 University, Durham, UK

11 ⁵ Eco-anthropologie (EA), Muséum national d'Histoire naturelle, CNRS, Université Paris Cité,

12 Paris, France

13

14 Short title: Chemical signaling in Japanese macaques

15

16 Corresponding author: Lucie Rigaill, EthoS - Ethologie animale et humaine, Campus de

17 Beaulieu - Bâtiment 25, 263, 35042 Rennes Cedex, France. lucie.rigaill@gmail.com

18

19

20 **Abstract**

21 Olfactory communication plays an important role in the regulation of socio-sexual interactions
22 in mammals. There is growing evidence that both human and non-human primates rely on odors
23 to inform their mating decisions. Nevertheless, studies of primate chemical ecology remain
24 scarce due to the difficulty of obtaining and analyzing samples. We analyzed 67 urine samples
25 from 5 captive female Japanese macaques (*Macaca fuscata*) and 30 vaginal swabs from 3 of
26 these females using gas chromatography–mass spectrometry and examined the relationship
27 between odor (compounds identified, richness, intensity, and diversity) and female identity as
28 well as cycle phase. We found a total of 36 urine compounds of which we identified 31, and 68
29 vaginal compounds of which we identified 37. Our results suggest that urine and vaginal odor
30 varied more between individuals than within cycle phases. However, we found that within a
31 female cycle, urine samples from similar phases may cluster more than samples from different
32 phases. Our results suggest that female odor may encode information about identity (vaginal
33 and urine odor) and reproductive status (urine odor). The question of how conspecifics use
34 female urine and vaginal odor remains open and could be tested using bioassays. Our results
35 and their interpretation are constrained by our limited sample size and our study design.
36 Nonetheless, our study provides insight into the potential signaling role of female odor in sexual
37 communication in Japanese macaques and contributes to our understanding of how odors may
38 influence mating strategies in primates.

39
40 Keywords: olfactory communication, gas chromatography–mass spectrometry, sexual
41 signaling, urine and vaginal odor, *Macaca fuscata*

42 **Introduction**

43 Communication plays a crucial role in mating decisions. Individuals are expected to choose or
44 avoid potential mates based on the costs (e.g., disease transmission, energy investment, sexual
45 competition, parental investment) and benefits (e.g., good genes, social support, increased
46 fitness) associated with mating (Beltran-Bech & Richard, 2014; Møller et al., 1999; Thrall et
47 al., 2000; Trivers, 1972). As a result, individuals exchange information about their attributes,
48 including social rank, genotype, health, and fertility (e.g., sexual maturity, reproductive status)
49 (Candolin, 2003; Snowdon, 2004), directly and indirectly benefitting both emitters and
50 receivers. Emitters may attract potential mates and enhance intra-sexual competition, thus
51 simultaneously increasing their mating opportunities and chances of producing attractive and
52 strong offspring (*sexy-sons/daughters* and *good genes* hypotheses R. A. Fisher, 1915; Hamilton
53 & Zuk, 1982). Meanwhile, receivers may balance the cost of mating by focusing their mating
54 effort on mates that are fertile and capable of dealing with the costs of reproduction and by
55 using information about their rivals' state to adjust their mating strategies to the level of
56 competition.

57 Like many other animals, primates communicate through multiple sensory channels.
58 Females display multiple traits that are attractive to males and correlate with their fixed (e.g.,
59 social rank, age) or variable (e.g., reproductive status) attributes (Tables 1-2). Most studies of
60 these traits have focused on female visual traits such as sexual swellings (enlargement of the
61 anogenital area) and skin coloration in relation to fertility, probably because the sense of vision
62 is particularly well developed in primates (Jacobs, 2008; Osorio & Vorobyev, 2005). Females
63 can also inform conspecifics about their fertility through auditory (vocalizations or voice),
64 behavioral (proceptive behaviors), and olfactory (odors) traits (Table 1).

Table 1. Examples of the potential ovulatory signaling function of female behavioral, visual, auditory, and olfactory traits in primates.

Species	Female trait	Reproductive status	Evidence
common marmosets, <i>Callithrix jacchus</i>	Anogenital odor	Correlates with the fertile phase.	Encoded information and receivers' response (Kücklich et al., 2019)
pygmy marmosets, <i>Cebuella pygmaea</i>	Anogenital gland or secretion odor	Correlates with the fertile phase.	Receivers' response (Converse et al., 1995)
tufted capuchins, <i>Cebus apella</i>	Proceptive behaviors	Correlate with the fertile phase.	Encoded information (Carosi & Visalberghi, 2002)
crested macaques, <i>Macaca nigra</i>	Sexual swelling (size)	Correlates with the fertile phase and conception ¹ and swellings are not expressed during pregnancy.	Encoded information (Higham et al., 2012)
	Proceptive behaviors	Correlate with the fertile phase and conception.	
	Copulation calls	Correlate with the fertile phase.	
Barbary macaques, <i>Macaca sylvanus</i>	Sexual swelling (size)	Correlates with the fertile phase and swellings are expressed during early pregnancy.	Encoded information (Brauch et al., 2007; Möhle et al., 2005; Young et al., 2013)
	Proceptive behaviors	Not clear.	Encoded information (Brauch et al., 2007; Young et al., 2013)
	Copulation calls	Do not correlate with the fertile phase.	Encoded information (Pfefferle et al., 2008, 2011; Semple & McComb, 2000) Receivers' response (Semple & McComb, 2000)
stumptailed macaques, <i>Macaca arctoides</i>	Vaginal secretions	May indicate the periovulatory period and stimulate male sexual arousal	Receivers' response (Cerdeña-Molina et al., 2006)

long-tailed macaques, <i>Macaca fascicularis</i>	Sexual swelling (size)	Does not correlate with the fertile phase and swellings are expressed during early pregnancy.	Encoded information (Engelhardt et al., 2005, 2007)
	Proceptive behaviors	Correlate with the fertile phase and are expressed during the early pregnancy.	
	Copulation calls	Do not correlate with the fertile phase.	Encoded information (Engelhardt et al., 2012)
Japanese macaques, <i>Macaca fuscata</i>	Red skin coloration (face and hindquarters)	Correlates with the pregnancy period (face coloration) but not with the fertile phase or conception.	Encoded information (Rigaill et al., 2015; Rigaill, MacIntosh, et al., 2017; Rigaill et al., 2019) Receivers' response (Rigaill & Garcia, 2021)
	Proceptive behaviors	Correlate with the pregnancy period but not with the fertile phase.	Encoded information (Garcia et al., 2009; O'Neill et al., 2004; Rigaill et al., 2015)
	Copulation and 'estrus' calls	Do not correlate with the fertile phase, and may correlate with pregnancy.	
	Urine odor	May not correlate with the fertile phase.	Receivers' response (Rigaill, Suda-Hashimoto, et al., 2017)
rhesus macaques, <i>Macaca mulatta</i>	Red skin coloration (face)	Correlates with the fertile phase.	Encoded information (Dubuc et al., 2009; Higham et al., 2010) Receivers' response (Higham et al., 2011)
olive baboons, <i>Papio anubis</i>	Sexual swelling (size or shape)	Correlates with the fertile phase and pregnancy.	Encoded information (Altmann, 1970; Higham, Heistermann, et al., 2008; Higham, MacLarnon, et al., 2008; Rigaill et al., 2013)
	Red skin coloration (sexual swelling)	Correlates with pregnancy status but not with the fertile phase.	Encoded information (Higham, MacLarnon, et al., 2008; Rigaill et al., 2013)
	Proceptive behaviors	May correlate with the fertile phase.	Encoded information (Higham et al., 2009; Rigaill et al., 2013)
	Vaginal secretions	Correlate with the fertile phase	Encoded information (Vaglio et al., 2021) Receivers' response (Rigaill et al., 2013)

yellow baboons, <i>Papio cynocephalus</i>	Copulation calls	Do not correlate with the fertile phase.	Encoded information and receivers' response (Semple, 2001; Semple et al., 2002)
chacma baboons, <i>Papio ursinus</i>	Vaginal odor	Correlates with the fertile phase	Receivers' response (P. M. R. Clarke et al., 2009)
mandrills, <i>Mandrillus sphinx</i>	Sexual swelling (size or shape)	May not correlate with conception.	Encoded information (Huchard et al., 2009; Setchell & Wickings, 2004a, 2004b)
	Red skin coloration (sexual swelling)	May not correlate with conception.	Encoded information (Setchell, Charpentier, et al., 2006; Setchell & Wickings, 2004a)
	Red skin coloration (face)	Correlates with the pregnancy status, but not with the fertile phase or conception.	Encoded information (Setchell et al., 2009; Setchell, Wickings, et al., 2006)
sooty mangbeys, <i>Cercocebus torquatus atys</i>	Sexual swelling (size)	Correlates with the fertile phase but not with pregnancy.	Encoded information (Gordon et al., 1991)
howler monkeys, <i>Alouatta pigra</i>	Proceptive behaviors	Correlate with the fertile phase.	Encoded information (Van Belle et al., 2009)
bonobos, <i>Pan paniscus</i>	Sexual swelling (size)	Does not correlate with the fertile phase or pregnancy.	Encoded information (Douglas et al., 2016; Furuichi, 1987)
chimpanzees, <i>Pan troglodytes</i>	Sexual swelling (size)	Correlates with the fertile phase but not with pregnancy.	Encoded information (Deschner et al., 2003, 2004; Emery & Whitten, 2003; Wallis & Lemmon, 1986)
	Copulation calls	Do not correlate of the fertile phase.	Encoded information (Townsend et al., 2011)
	Vaginal secretions	May correlate with the fertile phase	Encoded information (Matsumoto-Oda et al., 2003)
humans, <i>Homo sapiens</i>	Red skin coloration (cheeks and lips)	Does not correlate with the fertile phase.	Encoded information (Burriss et al., 2015; Rigaill, 2020)
	Proceptive behaviors	Not clear.	Encoded information (Gangestad et al., 2002; Haselton et al., 2007)

	Voice pitch	Correlates with the fertile phase.	Encoded information and receivers' response (Pipitone & Gallup, 2008, 2012; Pisanski et al., 2018; Shoup-Knox et al., 2019)
	Body odor	Pleasantness may correlate with the period of highest fertility.	Receivers' response (Gildersleeve et al., 2012; Haselton et al., 2007; Kuukasjärvi et al., 2004; Singh & Bronstad, 2001) but see (Mei et al., 2022)

66 ¹ difference between conceptive and non-conceptive cycles.

67 In comparison to visual and auditory traits, little is known about the potential role of
68 olfactory traits in primate sexual communication. This is probably because primates were long
69 regarded as microsmatic (having a reduced sense of olfaction, Heymann, 2006; T. D. Smith &
70 Bhatnagar, 2004). However, there is now good evidence that odors encode information about
71 an individual's state and may regulate primate social interactions (Table 2). Therefore, there is
72 ample reason to suspect that odors also influence mating decisions in both human and non-
73 human primates. Indeed, males attend more to female odors during the period of highest fertility
74 than at other times (Table 1).
75

76 **Table 2.** Examples of individual characteristics encoded by odors in primates.

Species	Olfactory traits	Individual characteristics	Evidence
crowned lemurs, <i>Eulemur coronatus</i>	Anogenital odor (males: scent marks, females: secretion)	Correlates with sex.	Encoded information (Elwell et al., 2021)
ring-tailed lemurs, <i>Lemur catta</i>	Labial and scrotal secretions	Correlate with kinship, genetic compatibility and genetic quality. Are affected by health status.	Encoded information (Boulet et al., 2009, 2010; Charpentier et al., 2008; Grogan et al., 2019) Receivers' response (Charpentier et al., 2010; Grogan et al., 2019; Harris et al., 2018)
emperor tamarins, <i>Saguinus imperator</i>	Anogenital scent marks, scent gland secretions, and skin odor	Correlate with sex and differ between reproductive and non-reproductive individuals.	Encoded information (Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021)
Weddell's saddleback tamarins, <i>Leontocebus weddelli</i>	Anogenital scent marks, scent gland secretions, and skin odor	Correlate with sex and differ between reproductive and non-reproductive individuals.	Encoded information (Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021)
common marmosets, <i>Callithrix jacchus</i>	Circumgenital scent marks	Correlate with female familiarity, identity, age, and parity.	Encoded information (T. E. Smith et al., 1997; T. E. Smith, 2006; Kücklich et al., 2019) Receivers' response (Kücklich et al., 2019)
owl monkeys, <i>Aotus nancymaae</i>	Perianal gland secretions	Correlate with sex, age, and kinship.	Encoded information (Macdonald et al., 2008; Spence-Aizenberg et al., 2018)
rhesus macaques, <i>Macaca mulatta</i>	Body odor (genital area)	Correlates with familiarity or group membership.	Receivers' response (Henkel et al., 2015)
mandrills, <i>Mandrillus sphinx</i>	Sternal gland secretions and hairs	Correlate with sex, age, social rank in males, group identity, individual genetic quality, genetic compatibility, and health	Encoded information (Poirotte et al., 2017; Setchell et al., 2010, 2011; Vaglio et al., 2016) Receivers' response (Poirotte et al., 2017)

chimpanzees, <i>Pan troglodytes</i>	Urine odor	Correlates with group membership and kinship.	Receivers' response (Henkel & Setchell, 2018) ⁷⁷
humans, <i>Homo sapiens</i>	Axillary or body odor	Is affected by health but does not correlate with human leucocyte antigen genotype.	Receivers' response (Olsson et al., 2014; Probst et al., 2017)

78 Variation in the chemical composition of female odor, such as the presence or absence
79 of compounds and their relative abundance, across and within individuals may inform
80 conspecifics about a female's fertility. For example, sex hormones can influence the
81 composition of female odor (humans: Michael et al., 1974; rhesus macaques: Michael &
82 Keverne, 1970). Across primate species, the chemical composition of female odor varies
83 between mating and non-mating seasons (ring-tailed lemurs: Scordato & Drea, 2007; Greene
84 & Drea, 2014; and potentially Milne-Edwards' sifakas, *Propithecus edwardsi*: Morelli et al.,
85 2013), across cycle phases (common marmosets: Kücklich et al., 2019; olive baboons: Vaglio
86 et al., 2021; but see for chimpanzees: Fox, 1982), and between reproductive and non-
87 reproductive individuals (tamarins: Poirier, Waterhouse, Dunn, et al., 2021; Poirier,
88 Waterhouse, Watsa, et al., 2021; owl monkeys: Spence-Aizenberg et al., 2018). However, only
89 one study of naturally cycling female catarrhines (Afroeurasian monkeys and apes, including
90 humans) has yet found evidence that fine-scale intra-cycle variation in vaginal odor intensity
91 correlates with ovulation (Vaglio et al., 2021). This limits our understanding of whether and
92 how female odors modulate male and female mating strategies across primates.

93 Here, we aimed to determine the potential signaling function (i.e., encoded information)
94 of female odor in relation to sexual communication in Japanese macaques (*Macaca fuscata*). In
95 this seasonal species, female behaviors, vocalizations, and skin coloration vary with female
96 reproductive status, both between cycle phases (Garcia et al., 2009; O'Neill et al., 2004; Rigaill
97 et al., 2019), and from pre- to post-conception periods (Rigaill et al., 2015), but do not
98 accurately signal ovulation (Rigaill & Garcia, 2021). It is not clear yet if males have access to
99 other information about ovulation. Males biased their mating effort to the fertile phase in one
100 study (Garcia et al., 2009) but not in others (O'Neill et al., 2004; Rigaill & Garcia, 2021).
101 However, females exhibit a strong vaginal or urinary odor during the mating season and males
102 frequently inspect females' genital areas by sniffing or touching them (Garcia & Rigaill,

103 unpublished observations of Wakasa captive group housed at Kyoto University Primate
104 Research Institute during the mating seasons in 2011-12 and 2015-16, and of Main Group of
105 Kojima Island, Wildlife Research Center, Kyoto University during the mating season in 2013-
106 14). Female odor may thus vary between individuals and cycle phases, as shown in other
107 primates (Table 1), and guide male mating behaviors.

108 We aimed to determine if female odor contains information about a female's timing of
109 ovulation. Here, we describe female urine and vaginal odor in Japanese macaques, focusing on
110 their chemical composition, complexity, intensity, and diversity, and examine how odor varies
111 between individuals and cycles (i.e., with a female's cycle) and within cycles (i.e., between pre-
112 fertile, fertile, and post-fertile phases). Our sample size is small (67 samples from 5 females for
113 urine, 30 samples from 3 females for vaginal odor), due to the difficulty inherent in obtaining
114 samples, but our study informs two hypotheses:

- 115 1. Female vaginal or urine odor conveys information about individual characteristics. If
116 so, we predict that the chemical composition, complexity, intensity, or diversity of
117 female urine or vaginal odor will vary across individuals. We also predict greater
118 variation across than within individuals.
- 119 2. Female odor conveys information about the timing of ovulation. If so, we predict that
120 the chemical composition, complexity, intensity, or diversity of vaginal and urine odor
121 will vary between a given cycle's cycle phases, particularly between the fertile phase
122 and the pre- and post-fertile phases.

123

124 **Methods**

125 **Study subjects**

126 We sampled 5 adult and naturally cycling female Japanese macaques ($12.2 \pm \text{SD } 3.0$ years old,
127 range = 8–16 years) housed at the Kyoto University Center for the Evolutionary Origins of

128 Human Behavior (EHUB, Inuyama, Japan; previously the Primate Research Institute). During
129 the training period, females were housed in individual cages (W90 cm, D76 cm, H85 cm) in
130 the same room (size: W4.0 m, D6.0 m, temperature: 27°C). Females were moved to a different
131 room (individual cage size: W78 cm, D65 cm, H80 cm, temperature: 20°C, window to the
132 external environment) for the data collection period. In both periods, females were housed with
133 3 adult males with whom they had visual, auditory, and olfactory communication but no
134 physical contact. Animals were fed twice daily (monkey pellets and sweet potatoes). Water was
135 supplied *ad libitum*.

136 Our research protocol was reviewed and approved by the EHUB Center for Human
137 Evolution Modeling Research (research protocol 2014-082-10). Our methods comply with the
138 Guidelines for the Care and Use of Nonhuman Primates of Kyoto University EHUB and with
139 the American Society of Primatologists Principles for the Ethical Treatment of Non-Human
140 Primates.

141

142 **Animal training**

143 We used positive reinforcement (Fernström et al., 2009) to train female macaques to present
144 their anogenital area and allow us to collect vaginal swabs for odor sampling (methods
145 developed by NSH for EHUB). We trained females for 26 weeks from May to October 2014.
146 Each female was trained during a 2-min session between 10.30 and 11.30 AM, 3 to 5 times a
147 week. One trainer (NSH or LR) conducted the training session with one observer (LR, NSH, or
148 LD). We used 1 cm³ diced apples and peanuts as the primary reinforcer, and a clicker as a
149 secondary reinforcer. We carried out training step by step following a predetermined order of
150 defined behaviors (see supplementary material, Table S1 adapted from Fernström et al., 2009).
151 One female failed to reach cooperation level 2 after 60 training sessions. We thus excluded her
152 from our training protocol and replaced her with another cycling adult female. We conducted

153 429 training sessions in total (mean per female = 86.8 sessions, range = 28–103 sessions).
154 Success rates varied across individuals. All females achieved anogenital presentation training
155 level 1 but only 2 allowed vaginal sampling. These 2 females already had experience with
156 positive reinforcement training for similar or other purposes at EHUB. For vaginal sampling,
157 we sampled these 2 females along with a third female, who did not complete training, but
158 spontaneously presented her hindquarters to the experimenter and accepted odor sampling
159 during the data collection period.

160

161 **Collection of vaginal and urine samples**

162 We collected data during the 2014–2015 mating season, from early November to late January
163 (85 days in total). One experimenter (LR) collected vaginal and urine samples between 08.30
164 AM and 11.00 AM. We aimed to collect samples every 2 days starting from the end of the first
165 observed menstruation, which we determined as the day when no fresh blood was observed on
166 the female anogenital area. Mean menstrual cycle length is $27.3 \pm \text{SD } 5.1$ days in both wild and
167 captive populations (Enomoto et al., 1979; Fooden & Aimi, 2005; Garcia et al., 2009; Nigi,
168 1975; O’Neill et al., 2004). If the experimenter failed to collect a vaginal or urine sample on a
169 designated day, they tried again the following day. The experimenter collected vaginal samples
170 by gently rubbing a sterile cotton swab (Rikaken, Japan) 5 times on the vaginal walls (N = 3
171 females). The experimenter started vaginal sample collection at least 10 min after entering the
172 experimental room and after any observed micturition. The experimenter collected
173 environmental controls by exposing sterile swabs to the air in the room to later identify volatiles
174 that did not derive from the sampled females. To sample urine, the experimenter placed a plastic
175 container, covered with mesh to prevent contamination from feces, under each female’s
176 individual cage to collect urine samples (approximately 2 ml) directly after micturition (N = 5
177 females). Our use of plastic containers to collect urine and the lack of matching controls such

178 as water collected from the same plastic containers to test for any effects of using plastic
179 containers or vials on our urine results may affect subsequent chemical analyses (Drea et al.,
180 2013).

181 We stored all samples in sterile sealed glass vials at $-80\text{ }^{\circ}\text{C}$ at EHUB to prevent chemical
182 degradation of the volatile compounds (Drea et al., 2013). We used screw-capped clear glass
183 vials of different diameters (D) and heights (H) (19x 10 ml vials of D 18 mm X H 50 mm, 94x
184 20 ml vials D 25 mm X H 50 mm, 8x 50ml vials of D 30 mm X H 80 mm). We used different
185 sizes of vial to store different types of sample. This may have affected our results because
186 evaporation of samples depends on the available air space, and may be confounded with our
187 predictor variables. In parallel, we kept a fraction of each urine sample at $-20\text{ }^{\circ}\text{C}$ for hormonal
188 analyses to determine the female's reproductive status. In total, we collected 35 vaginal samples
189 (mean per female = $11.7 \pm \text{SD } 4.5$, $N = 3$ females), 169 urine samples (mean per female = 33.8
190 $\pm \text{SD } 6.7$, $N = 5$ females), and 45 environmental control samples over two consecutive
191 menstrual cycles for each female.

192

193 **Determination of ovulation periods**

194 We determined sex hormone concentrations in a subset of 157 urine samples selected to
195 determine ovulation dates (2 consecutive menstrual cycles per female, 10 menstrual cycles in
196 total, mean per female = $31.4 \pm \text{SD } 5.9$ samples). This sample is slightly smaller than the total
197 because we excluded samples collected on consecutive days, and those collected just before or
198 after menstruation. We analyzed urine samples for estrone conjugates (E1C) and pregnanediol-
199 3-glucuronide (PdG) using previously described and validated enzyme immunoassays (EIA)
200 (Rigaill, Suda-Hashimoto, et al., 2017). We estimated that ovulation (day 0) occurred the day
201 following a urinary E1C peak associated with a continuous raise in PdG concentrations (Fujita
202 et al., 2001). We labelled the day directly preceding the estimated ovulation day as day -1, the

203 day directly following it as day +1, and so on. We defined the fertile phase as the 5-day period
204 covering day -2 to day +2 to account for the life span of sperm in the female tract (macaques:
205 Behboodi et al., 1991). The pre-fertile and post-fertile phases covered the 5 days preceding (day
206 -7 to day -3) and following (day +3 to day +7) the fertile phase. Hormonal profiles showed that
207 one of the 5 females (female 1) had abnormal hormonal patterns after her first cycle, so we
208 excluded her second cycle from the analyses. The cycle length of the 8 cycles for which we
209 observed menstrual bleeding was $29.4 \pm \text{SD } 6.9$ days.

210

211 **Chemical analyses**

212 We used 67 urine samples (mean per female = $7.4 \pm \text{SD } 0.7$ samples, 2 consecutive cycles per
213 female except for one female, Table 3) and 30 vaginal samples (mean per female = $7.5 \pm \text{SD}$
214 0.6 samples, one menstrual cycle for two females and two consecutive menstrual cycles for the
215 remaining female, Table 4) collected during the pre-fertile, fertile, and post-fertile phases for
216 odor analyses, along with 24 corresponding environmental control samples collected on the
217 same day as the test samples. We did not analyze the remaining 102 urine samples (mean per
218 female = $21.4 \pm \text{SD } 8.0$ samples) and 5 vaginal samples (5 samples from female 1, second
219 abnormal cycle) collected outside the 15-day period of interest.

220 We carried out laboratory analyses of odor at the Laboratory for Analysis and Research
221 in Environmental Chemistry - Italian National Research Council, Florence, Italy. We shipped
222 samples via airmail, using dry ice to keep them cold (-30°C). We investigated the volatile
223 compounds found in the Japanese macaque odor using solid-phase microextraction and gas
224 chromatography-mass spectrometry, applying the same methods used in other work on primate
225 odor signals (reviewed in Walker & Vaglio, 2021).

226 We introduced a $65 \mu\text{m}$ polydimethylsiloxane/divinylbenzene solid-phase
227 microextraction syringe needle through the vial septum and exposed the fiber to the headspace

228 above the sample in the vial for 15 minutes at 40°C. We analyzed the adsorbed volatile analytes
229 using a 5975C mass spectrometer (Agilent Technologies, Santa Clara, CA) EI, 70 eV, coupled
230 directly to a 7890B gas chromatograph (Agilent Technologies) equipped with a fused silica
231 HP5-MS UI capillary column (Agilent Technologies) 30 m x 0.25 mm crossbonded 5%-phenyl-
232 95%-dimethylpolysiloxane, film thickness 0.25µm. We maintained the injector temperature at
233 270°C and transfer line temperature at 280°C. We made injections in splitless mode (purge valve
234 opened after 1 minute) with a constant flow of helium carrier gas of 1 ml per minute. We started
235 the oven temperature program at 45°C for 2 minutes, then raised it by 4°C per minute to 170
236 °C and then by 20°C per minute to 300°C.

237 We assessed potential contamination from the laboratory environment using blank
238 analyses of an empty 10 ml vial (Supelco) following the same procedure used for the samples
239 (laboratory controls). We conditioned the fiber at 260°C for 5 minutes pre-injection and 20
240 minutes post-injection to avoid any possible carry-over effects. We conducted these
241 conditioning steps every day before analyzing samples.

242

243 **Identification of compounds**

244 From visual inspection of chromatograms (see examples of typical chromatograms in the
245 supplementary material, Figure S1), we detected 36 peaks from urine samples and 68 peaks
246 from vaginal samples that were absent in both environmental and laboratory controls. We found
247 25 compounds that did not derive from the animals and removed these from the swab results.

248 We standardized peak retention times using an internal standard (alpha pinene). We integrated
249 chromatograms to obtain retention time and peak area data using ChemStation software
250 (Agilent Technologies). We tentatively identified the eluted compounds by comparing the
251 experimental spectra with the spectra provided by the National Institute of Standards and
252 Technology (NIST) mass spectral database, version MSDF.01.01.2317 (Agilent Technologies).

253 We considered the identification valid when the minimum matching factor exceeded 80%. If
254 more than one compound was a good match for the same chromatographic peak, we compared
255 Kovats' retention index with values reported in the literature for the same chromatographic
256 column type to minimize the chances of misidentification. We also checked whether more than
257 one compound co-eluted below the same detected peak by checking the mass fragments using
258 ChemStation software.

259 We determined the relative amounts of compounds by integrating the areas of the
260 corresponding peaks in the total ion current profile and calculated percentages with respect to
261 the total area. We retained peaks that comprised at least 0.05% of the total area of the
262 chromatogram to avoid problems associated with unreliable quantification at very low relative
263 amounts (i.e., to exclude the background noise), although this may mean that we missed trace
264 chemicals (T. E. Smith et al., 2001). We analyzed all samples over a few days to minimize
265 interassay variability.

266 We describe the chemical composition of the 67 urine and 30 vaginal samples according
267 to female identity as well as cycle phases, including the percentage of samples in each category
268 that included each compound detected for urine (N = 36 compounds) and vaginal (N = 68
269 compounds) samples (Tables 3-4). For the following analyses, we excluded all compounds that
270 co-eluted as we cannot be sure about the measure of their contribution to the total ion current
271 (Drea et al., 2013). Two urine samples only contained co-eluted compounds, this may be due
272 to a mistake occurring at any step between data collection and chemical evaluation. We thus
273 removed these samples from the data set. Overall, we excluded 1 of 36 (3%) urine compounds
274 (Table 3) and 10 of 68 (15%) vaginal compounds (Table 4).

275

276 **Data analyses**

277 We carried out analyses using R software (version 4.0.3). We conducted our analyses on 35
278 compounds detected across 65 urine and 58 compounds detected across 30 vaginal samples.

279 We investigated the relationship between female odor and identity as well as cycle
280 phase. Several quantitative or qualitative analytic tools can be used to investigate chemical data.
281 The choice of one approach over another largely depends on the nature of the data (and the
282 methods used for chemical analyses), the species (e.g., whether chemical profiles are composed
283 of single or multiple compounds), the number of samples, and the research question (Drea et
284 al., 2013). Some studies assess the presence/absence of one or few compounds or categories of
285 compounds of interest (e.g., fatty acids or alcohols), while others reduce the dimensionality of
286 the data set (Drea et al., 2013). However, these methods may underestimate or overestimate
287 variance in chemical composition between samples by analyzing only a subset of the data (Drea
288 et al., 2013).

289 We opted for a conservative approach, keeping all compounds of the animal swab
290 results in the analyses and avoiding subjective thresholds such as retaining compounds or
291 variables based on *a priori* assumptions about these compounds' roles or on mathematical
292 rather than biological reasons. We used Nonmetric Multidimensional Scaling (NMDS), a rank-
293 based approach relying on distance or dissimilarity matrix, to represent pairwise dissimilarity
294 between samples in a dimensional space.

295 We tested for differences in both urine and vaginal samples according to female identity
296 and cycle phases. We computed a distance matrix for the data with the Bray-Curtis dissimilarity
297 index using the `vegdist` function in the *vegan* package (Oksanen et al., 2020). We then
298 calculated the 2-dimensional NMDS coordinates from these Bray-Curtis indices using the
299 `metaMDS` function in the *vegan* package and plotted the outcomes (Wickham, 2016). The stress
300 factors, which roughly represent the goodness of fit, associated with the coordinates were 0.19
301 for urine and 0.18 for vaginal data, and are thus considered fair (K. R. Clarke, 1993). We then

302 performed a permutational multivariate analysis of variance (PERMANOVA) using distance
303 matrices by using the *Adonis* function in (*vegan* package). This analysis tests whether the
304 centroids and dispersion of the groups tested as defined by measure space are equivalent for all
305 groups (Anderson, 2017). Because females contributed the data for one or two consecutive
306 menstrual cycles, we used “a female’s cycle” in our analyses as a proxy for both “menstrual
307 cycle” and “female identity”. Here we thus tested the effect of the interaction between a
308 female’s cycle and cycle phase (i.e., *Are samples from the same cycle phase more or less similar*
309 *than samples from another cycle phase within a female given cycle?*).

310 To facilitate comparison with other studies of similar research questions and substrates
311 (e.g., vaginal secretions, Vaglio et al., 2021), we also calculated three diversity indexes to
312 graphically investigate variation in sample richness (total number of detected peaks per
313 sample), Shannon’s H (which accounts for the relative abundance of each compound within a
314 sample) and odor intensity (total area of detected peaks per sample). Richness and Shannon H
315 of both urine and vaginal samples followed a normal distribution while urine and vaginal
316 intensity followed a lognormal distribution. We explored variation across females and cycle
317 phases graphically by plotting urine and vaginal richness, Shannon H, and intensity against
318 either female identity (categorical variable) or cycle phase (categorical variable).

319

320 **Data availability statement**

321 The data that support the findings of this study are openly available in figshare,
322 doi:10.6084/m9.figshare.19649766.

323

324 **Results**

325 **Identity of volatile compounds**

326 We detected 36 volatile compounds from the analysis of 67 urine samples. We proposed a
327 tentative identification for 31 of these compounds, including one that may result from chromatic
328 co-elution (Table 3). We detected 68 volatile compounds from the analysis of 30 vaginal
329 samples. We proposed a tentative identification for 37 of these vaginal compounds, with nine
330 compounds that may result from chromatic co-elution (Table 4). Only five volatile compounds
331 were found in both urine and vaginal samples (2-pentanone, 2-hexanone, dimethyl disulfide,
332 toluene, 2-heptanone, Tables 3-4). This represents 5% (5/104) of all the compounds detected in
333 urine and vaginal samples.

334

335 **Inter-individual variation in female odor**

336 Nineteen of 36 urine compounds (53%) were unique to one of the five females (Table 3).
337 Although the two dimensions extracted from the NMDS did not clearly cluster samples per
338 female identity (Fig. 1), the results of the PERMANOVA suggest that a female's cycle explains
339 25% of the variance in the data (Table 5). Urine richness, and odor intensity varied across
340 females (Fig. 2), with lower richness values in female 2 (Fig. 2A) and higher odor intensity in
341 female 1 (Fig. 2C). Shannon H index showed lower inter-individual variation (Fig. 2).

342 Twenty-eight of 68 vaginal compounds (41%) appeared to be unique to one of the three
343 females (Table 4). The two dimensions extracted from NMDS and the results of the
344 PERMANOVA suggest that a female's cycle explains 29% of the variance in the data and the
345 grouping patterns of samples (Fig. 3, Table 5). Vaginal richness, Shannon's H index, and odor
346 intensity varied across females (Fig. 2). One female (female 1) showed lower vaginal richness,
347 Shannon H, and intensity values than the two other females sampled (Fig. 2D, E, F).

348

349 **Intra-cycle variation in female odor**

350 Ten urine compounds (28%) were only found during the fertile phase (Table 3). The cycle phase
351 explained only 3% of the variance in urine data (Table 5). However, the interaction between a
352 female's cycle and cycle phase explained most of the variance in the data (27%, Table 5). Thus,
353 within a given female cycle, samples from same phases may cluster more than samples from
354 different phases. There was little intra-cycle variation in urine richness, Shannon H, or intensity
355 (Fig. 4). Urine richness, but not Shannon H (Fig. 4B), values tended to decrease from pre- to
356 post-fertile phase (Fig. 4A), while urine intensity increased slightly during the fertile phase
357 compared to pre- and post-fertile phases (Fig. 4C).

358 Six vaginal compounds (9%) were only found during the fertile phase (Table 4). The
359 cycle phase or the interaction between a female's cycle and cycle phase explained respectively
360 6% and 17% of the variance in vaginal data (Table 5). Vaginal richness, Shannon H, or intensity
361 did not appear to vary across the menstrual cycle (Fig. 4).

362 Individual trajectories are presented in the supplementary materiel (Fig. S2-3).

363

364 **Table 3.** The 36 chemical compounds retrieved from 65 urine samples, their tentative
365 identification, their mean peak area, and their presence in all samples and in samples from the
366 fertile phase. Data are presented for all females and for each female's menstrual cycle.
367 Compounds marked * co-eluted and are not included in graphical exploration of the data. ¹
368 Compounds often used as lab solvents (Drea et al., 2013) but kept in the analyses for
369 comparison with previously published data set (Delbarco-Trillo et al., 2013; Vaglio et al.,
370 2021).

371 *(Table 4 is provided as a separate file as it didn't fit in the manuscript)*

372

373 **Table 4.** The 68 chemical compounds retrieved from 30 vaginal samples, their tentative
374 identification, their mean peak area, and their presence in all samples and in samples from the

375 fertile phase. Data are presented for all females and for each female's menstrual cycle.
 376 Compounds marked * co-eluted and are not included in graphical exploration of the data. ¹
 377 Compounds often used as lab solvents (Drea et al., 2013) but kept in the analyses for
 378 comparison with previously published data set (Vaglio et al., 2021).

379 *(Table 4 is provided as a separate file as it didn't fit in the manuscript)*

380

381 **Table 5.** PERMANOVA results based on Bray-Curtis dissimilarities using abundance data for
 382 female odors from urine and vaginal samples

	df	Sum of Squares	Mean of Squares	F	R²
<i>Urine samples</i>					
Female's cycle	8	4.88	0.61	2.62	0.25
Cycle phase	2	0.63	0.31	1.35	0.03
Interaction	16	5.39	0.34	1.45	0.27
Residuals	38	8.84	0.23		0.45
Total	64	19.74			1.00
<i>Vaginal samples</i>					
Female's cycle	3	3.15	1.05	3.59	0.29
Cycle phase	2	0.62	0.31	1.06	0.06
Interaction	6	1.90	0.32	1.08	0.17
Residuals	18	5.27	0.29		0.48
Total	29	10.94			1.00

383

384 **Discussion**

385 We analyzed inter-individual and intra-cycle variation in urine and vaginal samples in female
 386 Japanese macaques based on identified compounds, odor richness, diversity and intensity. We
 387 identified 31 urine and 37 vaginal compounds of potential semiochemical importance in this
 388 species that may be of interest for future work. Although our limited sample size restricts the
 389 interpretation of our results, our analyses suggest that urine and vaginal odors varied more
 390 between females than between cycle phases. However, we also found that, within a female's
 391 cycle, urine samples from the same cycle phase may cluster more than samples from different
 392 phases.

393 We tentatively identified 31 (86%) of 36 volatile compounds found in urine samples
394 and 37 (54%) of 68 volatile compounds detected in female vaginal secretions. Some of the
395 volatile compounds we identified are known to play a role in plant or insect communication or
396 metabolism (source: <https://pubchem.ncbi.nlm.nih.gov>). Some of the compounds we detected
397 may be from the diced apples we used as a food reward during data collection (e.g., aldehydes:
398 hexanal, nonanal, decanal, benzaldehyde; alcohols: ethanol, 1-butanol, 3-methyl-; esters: ethyl
399 acetate, (Espino-Díaz et al., 2016)). However, it is also possible that these compounds are
400 widespread across plant and animal species due to their role in both intra- and inter-species
401 communication (e.g., pollination, alarm signals, mate attraction: (Das et al., 2013; Kelliher,
402 2007; Leonhardt et al., 2016). Indeed, 16 (51%) and 13 (35%) of the volatile compounds we
403 identified in urine and vaginal samples are found in female odor profiles (urine, vaginal
404 secretions, glands, hairs) of other primate species, from strepsirrhines to hominins (Tables 3-
405 4). Despite these general similarities across species, vaginal secretions have very dissimilar
406 odor profiles in Japanese macaques (Table 4) and olive baboons (Vaglio et al., 2021). This is
407 consistent with the hypothesis that (dis)similarities in odor profiles across primates relate to
408 socio-ecological and phylogenetic factors (Delbarco-Trillo et al., 2011; delBarco-Trillo &
409 Drea, 2014; Heymann, 2006; Jänig et al., 2019; Ueno, 1994).

410 We found that the relative abundance of compounds across urine and vaginal samples
411 varied across individuals in female Japanese macaques. Urine is known to play a role in the
412 individual recognition mechanisms of several species of mammals, including some platyrrhines
413 (mice, *Mus domesticus*: J. L. Hurst et al., 2001; common marmosets: T. E. Smith, 2006). To
414 our knowledge, our study is the first to show that urine odor, along with vaginal odor, may
415 convey information about identity in a catarrhine species.

416 Exchanging information about identity through odors may benefit Japanese macaques,
417 a species that lives in large cohesive groups (Fooden & Aimi, 2005). As proposed by Henkel

418 and colleagues for rhesus macaques (Henkel et al., 2015), the ability to identify individuals –
419 and thus potentially kin, familiar, or higher-ranking conspecifics – may be crucial to coordinate
420 movement and regulate social interactions. Males may pay particular attention to female odor
421 during the mating season. Dominant or central males, which have a closer access to females,
422 could use female anogenital (vaginal and urine) odor to follow and mate-guard fertile females
423 during group movement and to frustrate females' attempts to escape male monopolization.
424 Subordinate or peripheral males may also use female odor to find mating opportunities. Urine
425 odor may be of particular interest in this latter case: since the information can be decoupled
426 from the emitters, receivers may follow such fingerprints in the environment to find receptive
427 and fertile females.

428 Our study provides some evidence that urine, but not vaginal, odor can be informative
429 about a female's reproductive status in Japanese macaques. This may explain why in a previous
430 study, male Japanese macaques did not investigate (i.e., smell or taste) urine samples from the
431 fertile phase more than samples from other phases, as they were exposed to unknown rather
432 than familiar females' odors (Rigaill, Suda-Hashimoto, et al., 2017). Across animals, urine is
433 known to play a role in sexual signaling (Rothschild's giraffes, *Giraffa camelopardalis*
434 *rothschildi*: Bercovitch et al., 2006; common squirrel monkeys, *Saimiri sciureus*: Candland et
435 al., 1980; woolly spider monkeys, *Brachyteles arachnoides*: Milton, 1985; capuchin monkeys,
436 *Cebus apella*: Phillips et al., 2011; Asian elephants, *Elephas maximus*: Rasmussen et al., 1982;
437 giant pandas, *Ailuropoda melanoleuca*: Swaisgood et al., 2002). Our results suggest that this
438 communicative process through urine may also be present in catarrhines. To investigate this
439 hypothesis further, studies should assess male response to familiar female urine odor across
440 different cycle phases using bioassays.

441 The fact that vaginal odor does not appear to encode information about the fertile period
442 is surprising considering previous results in other primate species (Table 1 and Introduction).

443 It is possible that our small sample size prevented us from detecting any intra-cycle variation.
444 Moreover, variation may occur within a few days of ovulation rather than across the broader
445 cycle phases we studied. More and finer-scaled data are needed to better understand the
446 relationship between vaginal odor and ovulatory signaling in Japanese macaques and other
447 primate species.

448 Exciting research questions remain to be tested. If female odor varies in relation to sex
449 hormone concentrations, then does odor vary with reproductive status in the same individual:
450 e.g., between mating and non-mating periods, from prepubescent to sexually mature, between
451 cycling and non-cycling phases, and from pre- to post-conception phases? Do female age,
452 genetic profile, and health status influence this relationship and how might this affect female
453 reproductive capacity and success? And, finally, does female odor modulate socio-sexual
454 interactions, especially in relation to the level of female-female competition, and if so, how?
455 This last question is particularly interesting as, besides advertising their fertility, females would
456 also benefit from receiving information about their rivals' fertility status to increase their own
457 mating success. They may intensify competition with other fertile females through agonistic
458 interactions (Baniel et al., 2018; A. C. Hurst et al., 2017) or by exaggerating their own
459 attractivity (e.g., behavioral or vocal solicitations (Fallon et al., 2016)). However, whether and
460 how female odor modulates the level of intra-sexual competition is little studied in mammals,
461 including primates (M. L. Fisher & Burch, 2021; Stockley et al., 2013). Investigating these
462 research questions will provide valuable information about primate olfactory communication.
463 However, doing so will be challenging in both captive and wild populations as such studies
464 require longitudinal observations and data collection, a considerable budget for data storage
465 and analyses, and may conflict with population management plans (e.g., use of contraceptives
466 in breeding programs preclude analyses of female cycles).

467 In conclusion, our study identifies 31 volatile urine compounds and 37 volatile vaginal
468 compounds of possible importance in olfactory communication in Japanese macaques. We
469 found evidence for inter-individual (urine and vaginal samples) and intra-cycle (urine samples)
470 differences in female odors. While we cannot draw clear conclusions about the role of female
471 odors in Japanese macaque sexual communication, our results contribute to studies
472 investigating how olfaction mediates socio-sexual interactions in human and non-human
473 primates. To assess whether female odor has an adaptive sexual signaling function, further work
474 is needed to increase sampling effort and determine if males and females can perceive inter-
475 individual and intra-cycle differences in female odors, and whether female odors modify the
476 receiver's behavior.

477

478 **Acknowledgments**

479 We thank the Center for Human Evolution Modeling Research of the Kyoto University Center
480 for the Evolutionary Origins of Human Behavior (EHUB) (previously Primate Research
481 Institute) for the use of the primate facilities. We also thank Keiko Mouri from EHUB for her
482 help in the lab and Louise Ducroix (LD) for her help with animal training. Finally, we thank
483 Prof. Luca Calamai and Dr Marco Michelozzi for their assistance with chemical analyses at the
484 Laboratory for Analysis and Research in Environmental Chemistry – Italian National Research
485 Council, Florence, Italy. We are thankful to two anonymous reviewers for their comments on
486 earlier versions of this manuscript. This work was financially supported by the Japanese Society
487 for the Promotion of Science (JSPS) to LR (Kakenhi 15J01067) and by the CNRS to CG (PICS
488 N°7258 - Multimodal advertisement of reproductive status in Japanese macaques (*Macaca*
489 *fuscata*)). The authors declare no conflict of interest.

490

491 **Figures**

492 **Fig 1.** Nonmetric multidimensional scaling plots of similarity in urine sample chemical
493 composition across cycles in relation to female identity and menstrual cycle phases.

494 **Fig 2.** Variation in urine (A, B, C) and vaginal (D, E, F) richness (A, D), Shannon H (B, E),
495 and intensity (C, F) across females. Plots show the median (black horizontal line), first and third
496 quartiles (top and bottom of the box), and the range (upper and lower whiskers) values.

497 **Fig 3.** Nonmetric multidimensional scaling plots of similarity in vaginal sample chemical
498 composition in relation to female identity and menstrual cycle phases.

499 **Fig 4.** Variation in urine (A, B, C) and vaginal (D, E, F) richness (A, D), Shannon H (B, E),
500 and intensity (C, F) across cycle phases. Plots show the median (black horizontal line), the first
501 and third quartiles (top and bottom of the box), and the range (upper and lower whiskers) values.

502

503 **Fig S1.** Example of typical chromatographs from urine (A) and vaginal (B) samples during the
504 fertile phase.

505 **Fig S2.** Individual variation in urine richness (A), Shannon H (B), and intensity (C) across cycle
506 phases. Plots show the median (black horizontal line), first and third quartiles (top and bottom
507 of the box), and the range (upper and lower whiskers).

508 **Fig S3.** Individual variation in vaginal richness (A), Shannon H (B), and intensity (C) across
509 cycle phase. Plots show the median (black horizontal line), first and third quartiles (top and
510 bottom of the box), and the range (upper and lower whiskers).

511

512 **References**

513 Altmann, S. A. (1970). The pregnancy sign in savannah baboons. *The Journal of Zoo Animal*
514 *Medicine*, 4(2), 8-12. <https://doi.org/10.2307/20094180>

- 515 Anderson, M. J. (2017). Permutational Multivariate Analysis of Variance (PERMANOVA). In
516 *Wiley StatsRef : Statistics Reference Online* (p. 1-15). John Wiley & Sons, Ltd.
517 <https://doi.org/10.1002/9781118445112.stat07841>
- 518 Baniel, A., Cowlshaw, G., & Huchard, E. (2018). Jealous females? Female competition and
519 reproductive suppression in a wild promiscuous primate. *Proc. R. Soc. B*, *285*(1886),
520 20181332. <https://doi.org/10.1098/rspb.2018.1332>
- 521 Behboodi, E., Katz, D., Samuels, S., Tell, L., Hendrickx, A., & Lasley, B. (1991). The use of a
522 urinary estrone conjugates assay for detection of optimal mating time in the
523 cynomolgus macaque (*Macaca fascicularis*). *Journal of Medical Primatology*, *20*(5),
524 229-234.
- 525 Beltran-Bech, S., & Richard, F.-J. (2014). Impact of infection on mate choice. *Animal*
526 *Behaviour*, *90*, 159-170. <https://doi.org/10.1016/j.anbehav.2014.01.026>
- 527 Bercovitch, F. B., Bashaw, M. J., & del Castillo, S. M. (2006). Sociosexual behavior, male
528 mating tactics, and the reproductive cycle of giraffe *Giraffa camelopardalis*.
529 *Hormones and Behavior*, *50*(2), 314-321.
530 <https://doi.org/10.1016/j.yhbeh.2006.04.004>
- 531 Boulet, M., Charpentier, M. J., & Drea, C. M. (2009). Decoding an olfactory mechanism of kin
532 recognition and inbreeding avoidance in a primate. *BMC Evolutionary Biology*, *9*, 281.
533 <https://doi.org/10.1186/1471-2148-9-281>
- 534 Boulet, M., Crawford, J. C., Charpentier, M. J. E., & Drea, C. M. (2010). Honest olfactory
535 ornamentation in a female-dominant primate. *Journal of Evolutionary Biology*, *23*(7),
536 1558-1563. <https://doi.org/10.1111/j.1420-9101.2010.02007.x>
- 537 Brauch, K., Pfefferle, D., Hodges, K., Möhle, U., Fischer, J., & Heistermann, M. (2007). Female
538 sexual behavior and sexual swelling size as potential cues for males to discern the

- 539 female fertile phase in free-ranging Barbary macaques (*Macaca sylvanus*) of
540 Gibraltar. *Hormones and Behavior*, 52(3), 375-383.
541 <https://doi.org/10.1016/j.yhbeh.2007.06.001>
- 542 Burriss, R. P., Troscianko, J., Lovell, P. G., Fulford, A. J. C., Stevens, M., Quigley, R., Payne, J.,
543 Saxton, T. K., & Rowland, H. M. (2015). Changes in women's facial skin color over the
544 ovulatory cycle are not detectable by the human visual system. *PLoS ONE*, 10(7),
545 e0130093. <https://doi.org/10.1371/journal.pone.0130093>
- 546 Candland, D. K., Blumer, E. S., & Mumford, M. D. (1980). Urine as a communicator in a New
547 World primate, *Saimiri sciureus*. *Animal Learning & Behavior*, 8(3), 468-480.
548 <https://doi.org/10.3758/BF03199636>
- 549 Candolin, U. (2003). The use of multiple cues in mate choice. *Biological Reviews*, 78(4),
550 575-595. <https://doi.org/10.1017/S1464793103006158>
- 551 Carosi, M., & Visalberghi, E. (2002). Analysis of tufted capuchin (*Cebus apella*) courtship and
552 sexual behavior repertoire : Changes throughout the female cycle and female
553 interindividual differences. *American Journal of Physical Anthropology*, 118(1), 11-24.
554 <https://doi.org/10.1002/ajpa.10083>
- 555 Cerda-Molina, A. L., Hernández-López, L., Rojas-Maya, S., Murcia-Mejía, C., & Mondragón-
556 Ceballos, R. (2006). Male-induced sociosexual behavior by vaginal secretions in
557 *Macaca arctoides*. *International Journal of Primatology*, 27(3), 791-807.
558 <https://doi.org/10.1007/s10764-006-9045-0>
- 559 Charpentier, M. J. E., Boulet, M., & Drea, C. M. (2008). Smelling right : The scent of male
560 lemurs advertises genetic quality and relatedness. *Molecular Ecology*, 17(14),
561 3225-3233. <https://doi.org/10.1111/j.1365-294X.2008.03831.x>

- 562 Charpentier, M. J. E., Crawford, J. C., Boulet, M., & Drea, C. M. (2010). Message 'scent' :
563 Lemurs detect the genetic relatedness and quality of conspecifics via olfactory cues.
564 *Animal Behaviour*, *80*(1), 101-108. <https://doi.org/10.1016/j.anbehav.2010.04.005>
- 565 Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community
566 structure. *Australian Journal of Ecology*, *18*(1), 117-143.
567 <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>
- 568 Clarke, P. M. R., Barrett, L., & Henzi, S. P. (2009). What role do olfactory cues play in chacma
569 baboon mating? *American Journal of Primatology*, *71*(6), 493-502.
570 <https://doi.org/10.1002/ajp.20678>
- 571 Converse, L. J., Carlson, A. A., Ziegler, T. E., & Snowdon, C. T. (1995). Communication of
572 ovulatory state to mates by female pygmy marmosets, *Cebuella pygmaea*. *Animal*
573 *Behaviour*, *49*(3), 615-621. [https://doi.org/10.1016/0003-3472\(95\)80194-4](https://doi.org/10.1016/0003-3472(95)80194-4)
- 574 Das, A., Lee, S.-H., Hyun, T. K., Kim, S.-W., & Kim, J.-Y. (2013). Plant volatiles as method of
575 communication. *Plant Biotechnology Reports*, *7*(1), 9-26.
576 <https://doi.org/10.1007/s11816-012-0236-1>
- 577 Delbarco-Trillo, J., Burkert, B. A., Goodwin, T. E., & Drea, C. M. (2011). Night and day : The
578 comparative study of strepsirrhine primates reveals socioecological and phylogenetic
579 patterns in olfactory signals. *Journal of Evolutionary Biology*, *24*(1), 82-98.
580 <https://doi.org/10.1111/j.1420-9101.2010.02145.x>
- 581 delBarco-Trillo, J., & Drea, C. M. (2014). Socioecological and phylogenetic patterns in the
582 chemical signals of strepsirrhine primates. *Animal Behaviour*, *97*, 249-253.
583 <https://doi.org/10.1016/j.anbehav.2014.07.009>
- 584 Delbarco-Trillo, J., Harelimana, I. H., Goodwin, T. E., & Drea, C. M. (2013). Chemical
585 differences between voided and bladder urine in the aye-aye (*Daubentonia*

- 586 *madagascariensis*) : Implications for olfactory communication studies. *American*
587 *Journal of Primatology*, 75(7), 695-702. <https://doi.org/10.1002/ajp.22083>
- 588 Deschner, T., Heistermann, M., Hodges, K., & Boesch, C. (2003). Timing and probability of
589 ovulation in relation to sex skin swelling in wild West African chimpanzees, *Pan*
590 *trogodytes verus*. *Animal Behaviour*, 66(3), 551-560.
591 <https://doi.org/10.1006/anbe.2003.2210>
- 592 Deschner, T., Heistermann, M., Hodges, K., & Boesch, C. (2004). Female sexual swelling size,
593 timing of ovulation, and male behavior in wild West African chimpanzees. *Hormones*
594 *and Behavior*, 46(2), 204-215. <https://doi.org/10.1016/j.yhbeh.2004.03.013>
- 595 Douglas, P. H., Hohmann, G., Murtagh, R., Thiessen-Bock, R., & Deschner, T. (2016). Mixed
596 messages : Wild female bonobos show high variability in the timing of ovulation in
597 relation to sexual swelling patterns. *BMC Evolutionary Biology*, 16, 140.
598 <https://doi.org/10.1186/s12862-016-0691-3>
- 599 Drea, C. M., Boulet, M., Delbarco-Trillo, J., Greene, L. K., Sacha, C. R., Goodwin, T. E., &
600 Dubay, G. R. (2013). The « secret » in secretions : Methodological considerations in
601 deciphering primate olfactory communication. *American Journal of Primatology*,
602 75(7), 621-642. <https://doi.org/10.1002/ajp.22143>
- 603 Dubuc, C., Brent, L. J. N., Accamando, A. K., Gerald, M. S., MacLarnon, A., Semple, S.,
604 Heistermann, M., & Engelhardt, A. (2009). Sexual skin color contains information
605 about the timing of the fertile phase in free-ranging *Macaca mulatta*. *International*
606 *Journal of Primatology*, 30(6), 777-789. <https://doi.org/10.1007/s10764-009-9369-7>
- 607 Elwell, E. J., Walker, D., & Vaglio, S. (2021). Sexual dimorphism in crowned lemur scent-
608 marking. *Animals: An Open Access Journal from MDPI*, 11(7), 2091.
609 <https://doi.org/10.3390/ani11072091>

- 610 Emery, M. A., & Whitten, P. L. (2003). Size of sexual swellings reflects ovarian function in
611 chimpanzees (*Pan troglodytes*). *Behavioral Ecology and Sociobiology*, 54(4), 340-351.
612 <https://doi.org/10.1007/s00265-003-0648-6>
- 613 Engelhardt, A., Fischer, J., Neumann, C., Pfeifer, J.-B., & Heistermann, M. (2012). Information
614 content of female copulation calls in wild long-tailed macaques (*Macaca fascicularis*).
615 *Behavioral Ecology and Sociobiology*, 66(1), 121-134.
616 <https://doi.org/10.1007/s00265-011-1260-9>
- 617 Engelhardt, A., Hodges, J. K., & Heistermann, M. (2007). Post-conception mating in wild long-
618 tailed macaques (*Macaca fascicularis*) : Characterization, endocrine correlates and
619 functional significance. *Hormones and Behavior*, 51(1), 3-10.
620 <https://doi.org/10.1016/j.yhbeh.2006.06.009>
- 621 Engelhardt, A., Hodges, J. K., Niemitz, C., & Heistermann, M. (2005). Female sexual behavior,
622 but not sex skin swelling, reliably indicates the timing of the fertile phase in wild long-
623 tailed macaques (*Macaca fascicularis*). *Hormones and Behavior*, 47(2), 195-204.
624 <https://doi.org/10.1016/j.yhbeh.2004.09.007>
- 625 Enomoto, T., Seiki, K., & Haruki, Y. (1979). On the correlation between sexual behavior and
626 ovarian hormone level during the menstrual cycle in captive Japanese monkeys.
627 *Primates*, 20(4), 563-570. <https://doi.org/10.1007/BF02373438>
- 628 Espino-Díaz, M., Sepúlveda, D. R., González-Aguilar, G., & Olivas, G. I. (2016). Biochemistry of
629 apple aroma : A review. *Food Technology and Biotechnology*, 54(4), 375-397.
630 <https://doi.org/10.17113/ftb.54.04.16.4248>
- 631 Fallon, B. L., Neumann, C., Byrne, R. W., & Zuberbühler, K. (2016). Female chimpanzees
632 adjust copulation calls according to reproductive status and level of female

- 633 competition. *Animal Behaviour*, 113, 87-92.
- 634 <https://doi.org/10.1016/j.anbehav.2015.12.028>
- 635 Fernström, A.-L., Fredlund, H., Spångberg, M., & Westlund, K. (2009). Positive reinforcement
636 training in rhesus macaques—Training progress as a result of training frequency.
637 *American Journal of Primatology*, 71(5), 373-379. <https://doi.org/10.1002/ajp.20659>
- 638 Fisher, M. L., & Burch, R. L. (2021). Female-Female Competition. In T. K. Shackelford & V. A.
639 Weekes-Shackelford (Éds.), *Encyclopedia of Evolutionary Psychological Science* (p.
640 3043-3046). Springer International Publishing. [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-19650-3_1408)
641 [19650-3_1408](https://doi.org/10.1007/978-3-319-19650-3_1408)
- 642 Fisher, R. A. (1915). The evolution of sexual preference. *The Eugenics Review*, 7(3), 184-192.
- 643 Fooden, J., & Aimi, M. (2005). Systematic review of Japanese macaques, *Macaca fuscata*
644 (Gray, 1870). *Fieldiana Zoology*, 104, 1-198.
- 645 Fox, G. J. (1982). Potentials for pheromones in chimpanzee vaginal fatty acids. *Folia*
646 *Primatologica*, 37(3-4), 255-266. <https://doi.org/10.1159/000156036>
- 647 Fujita, S., Mitsunaga, F., Sugiura, H., & Shimizu, K. (2001). Measurement of urinary and fecal
648 steroid metabolites during the ovarian cycle in captive and wild Japanese macaques,
649 *Macaca fuscata*. *American Journal of Primatology*, 53(4), 167-176.
650 <https://doi.org/10.1002/ajp.3>
- 651 Furuichi, T. (1987). Sexual swelling, receptivity, and grouping of wild pygmy chimpanzee
652 females at Wamba, Zaïre. *Primates*, 28(3), 309-318.
653 <https://doi.org/10.1007/BF02381014>
- 654 Gangestad, S. W., Thornhill, R., & Garver, C. E. (2002). Changes in women's sexual interests
655 and their partners' mate-retention tactics across the menstrual cycle : Evidence for

- 656 shifting conflicts of interest. *Proceedings. Biological Sciences / The Royal Society*,
657 269(1494), 975-982. <https://doi.org/10.1098/rspb.2001.1952>
- 658 Garcia, C., Shimizu, K., & Huffman, M. (2009). Relationship between sexual interactions and
659 the timing of the fertile phase in captive female Japanese macaques (*Macaca*
660 *fuscata*). *American Journal of Primatology*, 71(10), 868-879.
661 <https://doi.org/10.1002/ajp.20717>
- 662 Gildersleeve, K. A., Haselton, M. G., Larson, C. M., & Pillsworth, E. G. (2012). Body odor
663 attractiveness as a cue of impending ovulation in women : Evidence from a study
664 using hormone-confirmed ovulation. *Hormones and Behavior*, 61(2), 157-166.
665 <https://doi.org/10.1016/j.yhbeh.2011.11.005>
- 666 Gordon, T. P., Gust, D. A., Busse, C. D., & Wilson, M. E. (1991). Hormones and sexual
667 behavior associated with postconception perineal swelling in the sooty mangabey
668 (*Cercocebus torquatus atys*). *International Journal of Primatology*, 12(6), 585-597.
669 <https://doi.org/10.1007/BF02547671>
- 670 Greene, L. K., & Drea, C. M. (2014). Love is in the air : Sociality and pair bondedness
671 influence sifaka reproductive signalling. *Animal Behaviour*, 88, 147-156.
672 <https://doi.org/10.1016/j.anbehav.2013.11.019>
- 673 Grogan, K. E., Harris, R. L., Boulet, M., & Drea, C. M. (2019). Genetic variation at MHC class II
674 loci influences both olfactory signals and scent discrimination in ring-tailed lemurs.
675 *BMC Evolutionary Biology*, 19(1), 171. <https://doi.org/10.1186/s12862-019-1486-0>
- 676 Hamilton, W. D., & Zuk, M. (1982). Heritable true fitness and bright birds : A role for
677 parasites? *Science (New York, N.Y.)*, 218(4570), 384-387.

- 678 Harris, R. L., Boulet, M., Grogan, K. E., & Drea, C. M. (2018). Costs of injury for scent
679 signalling in a strepsirrhine primate. *Scientific Reports*, *8*(1), 9882.
680 <https://doi.org/10.1038/s41598-018-27322-3>
- 681 Haselton, M. G., Mortezaie, M., Pillsworth, E. G., Bleske-Rechek, A., & Frederick, D. A. (2007).
682 Ovulatory shifts in human female ornamentation : Near ovulation, women dress to
683 impress. *Hormones and Behavior*, *51*(1), 40-45.
684 <https://doi.org/10.1016/j.yhbeh.2006.07.007>
- 685 Henkel, S., Lambides, A. R., Berger, A., Thomsen, R., & Widdig, A. (2015). Rhesus macaques
686 (*Macaca mulatta*) recognize group membership via olfactory cues alone. *Behavioral*
687 *Ecology and Sociobiology*, *69*(12), 2019-2034. [https://doi.org/10.1007/s00265-015-](https://doi.org/10.1007/s00265-015-2013-y)
688 [2013-y](https://doi.org/10.1007/s00265-015-2013-y)
- 689 Henkel, S., & Setchell, J. M. (2018). Group and kin recognition via olfactory cues in
690 chimpanzees (*Pan troglodytes*). *Proceedings of the Royal Society B: Biological*
691 *Sciences*, *285*(1889), 20181527. <https://doi.org/10.1098/rspb.2018.1527>
- 692 Heymann, E. W. (2006). The neglected sense-olfaction in primate behavior, ecology, and
693 evolution. *American Journal of Primatology*, *68*(6), 519-524.
694 <https://doi.org/10.1002/ajp.20249>
- 695 Higham, J. P., Brent, L. J. N., Dubuc, C., Accamando, A. K., Engelhardt, A., Gerald, M. S.,
696 Heistermann, M., & Stevens, M. (2010). Color signal information content and the eye
697 of the beholder : A case study in the rhesus macaque. *Behavioral Ecology*, *21*(4),
698 739-746. <https://doi.org/10.1093/beheco/arq047>
- 699 Higham, J. P., Heistermann, M., Ross, C., Semple, S., & Maclarnon, A. (2008). The timing of
700 ovulation with respect to sexual swelling detumescence in wild olive baboons.
701 *Primates*, *49*(4), 295-299. <https://doi.org/10.1007/s10329-008-0099-9>

- 702 Higham, J. P., Heistermann, M., Saggau, C., Agil, M., Perwitasari-Farajallah, D., & Engelhardt,
703 A. (2012). Sexual signalling in female crested macaques and the evolution of primate
704 fertility signals. *BMC Evolutionary Biology*, *12*, 89. [https://doi.org/10.1186/1471-](https://doi.org/10.1186/1471-2148-12-89)
705 [2148-12-89](https://doi.org/10.1186/1471-2148-12-89)
- 706 Higham, J. P., Hughes, K. D., Brent, L. J. N., Dubuc, C., Engelhardt, A., Heistermann, M.,
707 Maestriperi, D., Santos, L. R., & Stevens, M. (2011). Familiarity affects the assessment
708 of female facial signals of fertility by free-ranging male rhesus macaques. *Proceedings*
709 *of the Royal Society of London B: Biological Sciences*, *278*(1723), 3452-3458.
710 <https://doi.org/10.1098/rspb.2011.0052>
- 711 Higham, J. P., MacLarnon, A. M., Ross, C., Heistermann, M., & Semple, S. (2008). Baboon
712 sexual swellings : Information content of size and color. *Hormones and Behavior*,
713 *53*(3), 452-462. <https://doi.org/10.1016/j.yhbeh.2007.11.019>
- 714 Higham, J. P., Semple, S., MacLarnon, A., Heistermann, M., & Ross, C. (2009). Female
715 reproductive signaling, and male mating behavior, in the olive baboon. *Hormones*
716 *and Behavior*, *55*(1), 60-67. <https://doi.org/10.1016/j.yhbeh.2008.08.007>
- 717 Huchard, E., Benavides, J. A., Setchell, J. M., Charpentier, M. J. E., Alvergne, A., King, A. J.,
718 Knapp, L. A., Cowlshaw, G., & Raymond, M. (2009). Studying shape in sexual signals :
719 The case of primate sexual swellings. *Behavioral Ecology and Sociobiology*, *63*(8),
720 1231-1242. <https://doi.org/10.1007/s00265-009-0748-z>
- 721 Hurst, A. C., Alquist, J. L., & Puts, D. A. (2017). Women's fertility status alters other women's
722 jealousy and mate guarding. *Personality and Social Psychology Bulletin*, *43*(2),
723 191-203. <https://doi.org/10.1177/0146167216678859>
- 724 Hurst, J. L., Payne, C. E., Nevison, C. M., Marie, A. D., Humphries, R. E., Robertson, D. H.,
725 Cavaggioni, A., & Beynon, R. J. (2001). Individual recognition in mice mediated by

- 726 major urinary proteins. *Nature*, 414(6864), 631-634.
- 727 <https://doi.org/10.1038/414631a>
- 728 Jacobs, G. H. (2008). Primate color vision : A comparative perspective. *Visual Neuroscience*,
- 729 25(5-6), 619-633. <https://doi.org/10.1017/S0952523808080760>
- 730 Jänig, S., Weiß, B. M., Birkemeyer, C., & Widdig, A. (2019). Comparative chemical analysis of
- 731 body odor in great apes. *American Journal of Primatology*, 81(6), e22976.
- 732 <https://doi.org/10.1002/ajp.22976>
- 733 Kelliher, K. R. (2007). The combined role of the main olfactory and vomeronasal systems in
- 734 social communication in mammals. *Hormones and Behavior*, 52(5), 561-570.
- 735 <https://doi.org/10.1016/j.yhbeh.2007.08.012>
- 736 Kücklich, M., Weiß, B. M., Birkemeyer, C., Einspanier, A., & Widdig, A. (2019). Chemical cues
- 737 of female fertility states in a non-human primate. *Scientific Reports*, 9(1), 1-12.
- 738 <https://doi.org/10.1038/s41598-019-50063-w>
- 739 Kuukasjärvi, S., Eriksson, C. J. P., Koskela, E., Mappes, T., Nissinen, K., & Rantala, M. J. (2004).
- 740 Attractiveness of women's body odors over the menstrual cycle : The role of oral
- 741 contraceptives and receiver sex. *Behavioral Ecology*, 15(4), 579-584.
- 742 <https://doi.org/10.1093/beheco/arh050>
- 743 Leonhardt, S. D., Menzel, F., Nehring, V., & Schmitt, T. (2016). Ecology and evolution of
- 744 communication in social insects. *Cell*, 164(6), 1277-1287.
- 745 <https://doi.org/10.1016/j.cell.2016.01.035>
- 746 Macdonald, E. A., Fernandez-Duque, E., Evans, S., & Hagey, L. R. (2008). Sex, age, and family
- 747 differences in the chemical composition of owl monkey (*Aotus nancymaae*)
- 748 subcaudal scent secretions. *American Journal of Primatology*, 70(1), 12-18.
- 749 <https://doi.org/10.1002/ajp.20450>

- 750 Matsumoto-Oda, A., Oda, R., Hayashi, Y., Murakami, H., Maeda, N., Kumazaki, K., Shimizu, K.,
751 & Matsuzawa, T. (2003). Vaginal fatty acids produced by chimpanzees during
752 menstrual cycles. *Folia Primatologica*, 74(2), 75-79.
- 753 Mei, M., Grillot, R. L., Abbey, C. K., Emery Thompson, M., & Roney, J. R. (2022). Does scent
754 attractiveness reveal women's ovulatory timing? Evidence from signal detection
755 analyses and endocrine predictors of odour attractiveness. *Proceedings of the Royal
756 Society B: Biological Sciences*, 289(1970), 20220026.
757 <https://doi.org/10.1098/rspb.2022.0026>
- 758 Michael, R. P., Bonsall, R. W., & Warner, P. (1974). Human vaginal secretions : Volatile fatty
759 acid content. *Science (New York, N.Y.)*, 186(4170), 1217-1219.
- 760 Michael, R. P., & Keverne, E. B. (1970). Primate sex pheromones of vaginal origin. *Nature*,
761 225(5227), 84-85. <https://doi.org/10.1038/225084a0>
- 762 Milton, K. (1985). Urine washing behavior in the woolly spider monkey (*Brachyteles*
763 *arachnoides*). *Zeitschrift Für Tierpsychologie*, 67(1-4), 154-160.
764 <https://doi.org/10.1111/j.1439-0310.1985.tb01384.x>
- 765 Möhle, U., Heistermann, M., Dittami, J., Reinberg, V., Wallner, B., & Hodges, J. K. (2005).
766 Patterns of anogenital swelling size and their endocrine correlates during ovulatory
767 cycles and early pregnancy in free-ranging barbary macaques (*Macaca sylvanus*) of
768 Gibraltar. *American Journal of Primatology*, 66(4), 351-368.
769 <https://doi.org/10.1002/ajp.20161>
- 770 Møller, A. P., Christe, P., & Lux, E. (1999). Parasitism, host immune function, and sexual
771 selection. *The Quarterly Review of Biology*, 74(1), 3-20.

- 772 Morelli, T. L., Hayes, R. A., Nahrung, H. F., Goodwin, T. E., Harelimana, I. H., Macdonald, L. J.,
773 & Wright, P. C. (2013). Relatedness communicated in lemur scent. *Die*
774 *Naturwissenschaften*, 100(8), 769-777. <https://doi.org/10.1007/s00114-013-1074-x>
- 775 Nigi, H. (1975). Menstrual cycle and some other related aspects of Japanese monkeys
776 (*Macaca fuscata*). *Primates*, 16(2), 207-216. <https://doi.org/10.1007/BF02381418>
- 777 Oksanen, J., Blanchet, G. F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
778 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.
779 (2020). *vegan : Community ecology package*. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
780 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan)
- 781 Olsson, M. J., Lundström, J. N., Kimball, B. A., Gordon, A. R., Karshikoff, B., Hosseini, N.,
782 Sorjonen, K., Olgart Höglund, C., Solares, C., Soop, A., Axelsson, J., & Lekander, M.
783 (2014). The scent of disease : Human body odor contains an early chemosensory cue
784 of sickness. *Psychological Science*, 25(3), 817-823.
785 <https://doi.org/10.1177/0956797613515681>
- 786 O'Neill, A. C., Fedigan, L. M., & Ziegler, T. E. (2004). Relationship between ovarian cycle
787 phase and sexual behavior in female Japanese macaques (*Macaca fuscata*). *American*
788 *Journal of Physical Anthropology*, 125(4), 352-362.
789 <https://doi.org/10.1002/ajpa.20024>
- 790 Osorio, D., & Vorobyev, M. (2005). Photoreceptor spectral sensitivities in terrestrial animals :
791 Adaptations for luminance and colour vision. *Proceedings of the Royal Society of*
792 *London B: Biological Sciences*, 272(1574), 1745-1752.
793 <https://doi.org/10.1098/rspb.2005.3156>
- 794 Pfefferle, D., Brauch, K., Heistermann, M., Hodges, J. K., & Fischer, J. (2008). Female Barbary
795 macaque (*Macaca sylvanus*) copulation calls do not reveal the fertile phase but

- 796 influence mating outcome. *Proceedings. Biological Sciences / The Royal Society*,
797 275(1634), 571-578. <https://doi.org/10.1098/rspb.2007.1499>
- 798 Pfefferle, D., Heistermann, M., Pirow, R., Hodges, J. K., & Fischer, J. (2011). Estrogen and
799 progesterone correlates of the structure of female copulation calls in semi-free-
800 ranging Barbary macaques (*Macaca sylvanus*). *International Journal of Primatology*,
801 32(4), 992-1006. <https://doi.org/10.1007/s10764-011-9517-8>
- 802 Phillips, K. A., Buzzell, C. A., Holder, N., & Sherwood, C. C. (2011). Why do capuchin monkeys
803 urinate wash? An experimental test of the sexual communication hypothesis using
804 fMRI. *American Journal of Primatology*, 73(6), 578-584.
805 <https://doi.org/10.1002/ajp.20931>
- 806 Pipitone, R. N., & Gallup, G. G. (2008). Women's voice attractiveness varies across the
807 menstrual cycle. *Evolution and Human Behavior*, 29(4), 268-274.
808 <https://doi.org/10.1016/j.evolhumbehav.2008.02.001>
- 809 Pipitone, R. N., & Gallup, G. G. (2012). The unique impact of menstruation on the female
810 voice : Implications for the evolution of menstrual cycle cues. *Ethology*, 118(3),
811 281-291. <https://doi.org/10.1111/j.1439-0310.2011.02010.x>
- 812 Pisanski, K., Oleszkiewicz, A., Plachetka, J., Gmiterek, M., & Reby, D. (2018). Voice pitch
813 modulation in human mate choice. *Proceedings of the Royal Society B: Biological*
814 *Sciences*, 285(1893), 20181634. <https://doi.org/10.1098/rspb.2018.1634>
- 815 Poirier, A. C., Waterhouse, J. S., Dunn, J. C., & Smith, A. C. (2021). Scent marks signal species,
816 sex and reproductive status in tamarins (*Saguinus spp.*, neotropical primates).
817 *Chemical Senses*, bjab008. <https://doi.org/10.1093/chemse/bjab008>
- 818 Poirier, A. C., Waterhouse, J. S., Watsa, M., Erkenwick, G. A., Moreira, L. A. A., Tang, J.,
819 Dunn, J. C., Melin, A. D., & Smith, A. C. (2021). On the trail of primate scent signals : A

- 820 field analysis of callitrichid scent-gland secretions by portable gas chromatography-
821 mass spectrometry. *American Journal of Primatology*, 83(3), e23236.
822 <https://doi.org/10.1002/ajp.23236>
- 823 Poirotte, C., Massol, F., Herbert, A., Willaume, E., Bomo, P. M., Kappeler, P. M., &
824 Charpentier, M. J. E. (2017). Mandrills use olfaction to socially avoid parasitized
825 conspecifics. *Science Advances*, 3(4), e1601721.
826 <https://doi.org/10.1126/sciadv.1601721>
- 827 Probst, F., Fischbacher, U., Lobmaier, J. S., Wirthmüller, U., & Knoch, D. (2017). Men's
828 preferences for women's body odours are not associated with human leucocyte
829 antigen. *Proceedings of the Royal Society B: Biological Sciences*, 284(1864),
830 20171830. <https://doi.org/10.1098/rspb.2017.1830>
- 831 Rasmussen, L. E., Schmidt, M. J., Henneous, R., Groves, D., & Daves, G. D. (1982). Asian bull
832 elephants : Flehmen-like responses to extractable components in female elephant
833 estrous urine. *Science*, 217(4555), 159-162. <https://doi.org/10.1126/science.7089549>
- 834 Rigaill, L. (2020). Fine-scale variation in lip and cheek colour according to the timing of
835 ovulation in women. *Behavioral Ecology and Sociobiology*, 74(6), 71.
836 <https://doi.org/10.1007/s00265-020-02851-y>
- 837 Rigaill, L., & Garcia, C. (2021). Does male mate choice select for female coloration in a
838 promiscuous primate species? *Animal Behaviour*, 177, 171-181.
839 <https://doi.org/10.1016/j.anbehav.2021.05.002>
- 840 Rigaill, L., Higham, J. P., Lee, P. C., Blin, A., & Garcia, C. (2013). Multimodal sexual signaling
841 and mating behavior in olive baboons (*Papio anubis*). *American Journal of*
842 *Primatology*, 75(7), 774-787. <https://doi.org/10.1002/ajp.22154>

- 843 Rigaill, L., Higham, J. P., Winters, S., & Garcia, C. (2019). The redder the better? Information
844 content of red skin coloration in female Japanese macaques. *Behavioral Ecology and*
845 *Sociobiology*, 73(8), 103. <https://doi.org/10.1007/s00265-019-2712-x>
- 846 Rigaill, L., MacIntosh, A. J. J., Higham, J. P., Winters, S., Shimizu, K., Mouri, K., Furuichi, T., &
847 Garcia, C. (2015). Multimodal advertisement of pregnancy in free-ranging female
848 Japanese macaques (*Macaca fuscata*). *PLoS ONE*, 10(8), e0135127.
849 <https://doi.org/10.1371/journal.pone.0135127>
- 850 Rigaill, L., MacIntosh, A. J. J., Higham, J. P., Winters, S., Shimizu, K., Mouri, K., Suzumura, T.,
851 Furuichi, T., & Garcia, C. (2017). Testing for links between face color and age,
852 dominance status, parity, weight, and intestinal nematode infection in a sample of
853 female Japanese macaques. *Primates*, 58(1), 83-91. [https://doi.org/10.1007/s10329-](https://doi.org/10.1007/s10329-016-0575-6)
854 [016-0575-6](https://doi.org/10.1007/s10329-016-0575-6)
- 855 Rigaill, L., Suda-Hashimoto, N., Ducroix, L., Mouri, K., Furuichi, T., & Garcia, C. (2017). Testing
856 for links between female urine odor and male sexual behaviors in Japanese
857 macaques (*Macaca fuscata*). *International Journal of Primatology*, 38(5), 823-837.
858 <https://doi.org/10.1007/s10764-017-9980-y>
- 859 Scordato, E. S., & Drea, C. M. (2007). Scents and sensibility : Information content of olfactory
860 signals in the ringtailed lemur, *Lemur catta*. *Animal Behaviour*, 73(2), 301-314.
861 <https://doi.org/10.1016/j.anbehav.2006.08.006>
- 862 Semple, S. (2001). Individuality and male discrimination of female copulation calls in the
863 yellow baboon. *Animal Behaviour*, 61(5), 1023-1028.
864 <https://doi.org/10.1006/anbe.2001.1692>

- 865 Semple, S., & McComb, K. (2000). Perception of female reproductive state from vocal cues in
866 a mammal species. *Proceedings of the Royal Society of London. Series B: Biological*
867 *Sciences*, 267(1444), 707-712. <https://doi.org/10.1098/rspb.2000.1060>
- 868 Semple, S., McComb, K., Alberts, S., & Altmann, J. (2002). Information content of female
869 copulation calls in yellow baboons. *American Journal of Primatology*, 56(1), 43-56.
870 <https://doi.org/10.1002/ajp.1062>
- 871 Setchell, J. M., Charpentier, M. J. E., Abbott, K. M., Wickings, E. J., & Knapp, L. A. (2009). Is
872 brightest best? Testing the Hamilton-Zuk hypothesis in mandrills. *International*
873 *Journal of Primatology*, 30(6), 825-844. <https://doi.org/10.1007/s10764-009-9371-0>
- 874 Setchell, J. M., Charpentier, M. J. E., Bedjabaga, I.-B., Reed, P., Wickings, E. J., & Knapp, L. A.
875 (2006). Secondary sexual characters and female quality in primates. *Behavioral*
876 *Ecology and Sociobiology*, 61(2), 305-315. [https://doi.org/10.1007/s00265-006-0260-](https://doi.org/10.1007/s00265-006-0260-7)
877 7
- 878 Setchell, J. M., Vaglio, S., Abbott, K. M., Moggi-Cecchi, J., Boscaro, F., Pieraccini, G., & Knapp,
879 L. A. (2011). Odour signals major histocompatibility complex genotype in an Old
880 World monkey. *Proceedings. Biological Sciences / The Royal Society*, 278(1703),
881 274-280. <https://doi.org/10.1098/rspb.2010.0571>
- 882 Setchell, J. M., Vaglio, S., Moggi-Cecchi, J., Boscaro, F., Calamai, L., & Knapp, L. A. (2010).
883 Chemical composition of scent-gland secretions in an old world monkey (*Mandrillus*
884 *sphinx*) : Influence of sex, male status, and individual identity. *Chemical Senses*, 35(3),
885 205-220. <https://doi.org/10.1093/chemse/bjp105>
- 886 Setchell, J. M., Wickings, E. J., & Knapp, L. A. (2006). Signal content of red facial coloration in
887 female mandrills (*Mandrillus sphinx*). *Proceedings of the Royal Society B: Biological*
888 *Sciences*, 273(1599), 2395-2400. <https://doi.org/10.1098/rspb.2006.3573>

- 889 Setchell, J. M., & Wickings, J. E. (2004a). Sexual swelling in mandrills (*Mandrillus sphinx*) : A
890 test of the reliable indicator hypothesis. *Behavioral Ecology*, *15*(3), 438-445.
891 <https://doi.org/10.1093/beheco/arh027>
- 892 Setchell, J. M., & Wickings, J. E. (2004b). Social and seasonal influences on the reproductive
893 cycle in female mandrills (*Mandrillus sphinx*). *American Journal of Physical*
894 *Anthropology*, *125*(1), 73-84. <https://doi.org/10.1002/ajpa.10375>
- 895 Shoup-Knox, M. L., Ostrander, G. M., Reimann, G. E., & Pipitone, R. N. (2019). Fertility-
896 dependent acoustic variation in women's voices previously shown to affect listener
897 physiology and perception. *Evolutionary Psychology*, *17*(2), 1474704919843103.
898 <https://doi.org/10.1177/1474704919843103>
- 899 Singh, D., & Bronstad, P. M. (2001). Female body odour is a potential cue to ovulation.
900 *Proceedings of the Royal Society B: Biological Sciences*, *268*(1469), 797-801.
901 <https://doi.org/10.1098/rspb.2001.1589>
- 902 Smith, T. D., & Bhatnagar, K. P. (2004). Microsmatic primates : Reconsidering how and when
903 size matters. *The Anatomical Record Part B: The New Anatomist*, *279B*(1), 24-31.
904 <https://doi.org/10.1002/ar.b.20026>
- 905 Smith, T. E. (2006). Individual olfactory signatures in common marmosets (*Callithrix jacchus*).
906 *American Journal of Primatology*, *68*(6), 585-604. <https://doi.org/10.1002/ajp.20254>
- 907 Smith, T. E., Abbott, D. H., Tomlinson, A. J., & Mlotkiewicz, J. A. (1997). Differential display of
908 investigative behavior permits discrimination of scent signatures from familiar and
909 unfamiliar socially dominant female marmoset monkeys (*Callithrix jacchus*). *Journal*
910 *of Chemical Ecology*, *23*(11), 2523-2546.
911 <https://doi.org/10.1023/B:JOEC.0000006664.38169.e5>

- 912 Smith, T. E., Tomlinson, A. J., Mlotkiewicz, J. A., & Abbott, D. H. (2001). Female marmoset
913 monkeys (*Callithrix jacchus*) can be identified from the chemical composition of their
914 scent marks. *Chemical Senses*, 26(5), 449-458.
915 <https://doi.org/10.1093/chemse/26.5.449>
- 916 Snowdon, C. T. (2004). Sexual selection and communication. In C. P. van Schaik & P. M.
917 Kappeler (Éds.), *Sexual Selection in Primates : New and Comparative Perspectives* (p.
918 57-70). Cambridge University Press.
919 <https://doi.org/10.1017/CBO9780511542459.006>
- 920 Spence-Aizenberg, A., Kimball, B. A., Williams, L. E., & Fernandez-Duque, E. (2018). Chemical
921 composition of glandular secretions from a pair-living monogamous primate : Sex,
922 age, and gland differences in captive and wild owl monkeys (*Aotus spp.*). *American*
923 *Journal of Primatology*, 80(2), e22730. <https://doi.org/10.1002/ajp.22730>
- 924 Stockley, P., Bottell, L., & Hurst, J. L. (2013). Wake up and smell the conflict : Odour signals in
925 female competition. *Philosophical Transactions of the Royal Society B: Biological*
926 *Sciences*, 368(1631), 20130082. <https://doi.org/10.1098/rstb.2013.0082>
- 927 Swaisgood, R. R., Lindburg, D. G., & Zhang, H. (2002). Discrimination of oestrous status in
928 giant pandas (*Ailuropoda melanoleuca*) via chemical cues in urine. *Journal of Zoology*,
929 257(3), 381-386. <https://doi.org/10.1017/S0952836902000985>
- 930 Thrall, P. H., Antonovics, J., & Dobson, A. P. (2000). Sexually transmitted diseases in
931 polygynous mating systems : Prevalence and impact on reproductive success.
932 *Proceedings of the Royal Society B: Biological Sciences*, 267(1452), 1555-1563.
- 933 Townsend, S. W., Deschner, T., & Zuberbühler, K. (2011). Copulation calls in female
934 chimpanzees (*Pan troglodytes schweinfurthii*) convey identity but do not accurately


- 935 reflect fertility. *International Journal of Primatology*, 32(4), 914-923.
- 936 <https://doi.org/10.1007/s10764-011-9510-2>
- 937 Trivers, R. L. (1972). Parental investment and sexual selection. In *Sexual Selection and the*
- 938 *Descent of Man, 1871–1971*. B. Campbell.
- 939 Ueno, Y. (1994). Olfactory discrimination of urine odors from five species by tufted capuchin
- 940 (*Cebus apella*). *Primates*, 35(3), 311-323. <https://doi.org/10.1007/BF02382728>
- 941 Vaglio, S., Minicozzi, P., Kessler, S. E., Walker, D., & Setchell, J. M. (2021). Olfactory signals
- 942 and fertility in olive baboons. *Scientific Reports*, 11(1), 8506.
- 943 <https://doi.org/10.1038/s41598-021-87893-6>
- 944 Vaglio, S., Minicozzi, P., Romoli, R., Boscaro, F., Pieraccini, G., Moneti, G., & Moggi-Cecchi, J.
- 945 (2016). Sternal gland scent-marking signals sex, age, rank, and group identity in
- 946 captive mandrills. *Chemical Senses*, 41(2), 177-186.
- 947 <https://doi.org/10.1093/chemse/bjv077>
- 948 Van Belle, S., Estrada, A., Ziegler, T. E., & Strier, K. B. (2009). Sexual behavior across ovarian
- 949 cycles in wild black howler monkeys (*Alouatta pigra*) : Male mate guarding and
- 950 female mate choice. *American Journal of Primatology*, 71(2), 153-164.
- 951 <https://doi.org/10.1002/ajp.20635>
- 952 Walker, D., & Vaglio, S. (2021). Sampling and analysis of animal scent signals. *Journal of*
- 953 *Visualized Experiments: JoVE*, 168. <https://doi.org/10.3791/60902>
- 954 Wallis, J., & Lemmon, W. B. (1986). Social behavior and genital swelling in pregnant
- 955 chimpanzees (*Pan troglodytes*). *American Journal of Primatology*, 10(2), 171-183.
- 956 <https://doi.org/10.1002/ajp.1350100207>
- 957 Wickham, H. (2016). *ggplot2 : Elegant graphics for data analysis*. Springer-Verlag New York.
- 958 <https://ggplot2.tidyverse.org>

959 Young, C., Majolo, B., Heistermann, M., Schülke, O., & Ostner, J. (2013). Male mating
960 behaviour in relation to female sexual swellings, socio-sexual behaviour and
961 hormonal changes in wild Barbary macaques. *Hormones and Behavior*, 63(1), 32-39.
962 <https://doi.org/10.1016/j.yhbeh.2012.11.004>
963


Highlights




- We provide tentative identification of 31 urine and 37 vaginal compounds.
- Urine and vaginal odor vary more between than within female.
- Urine odor may convey information about a female reproductive status as within a female cycle, urine samples from similar phases cluster more than samples from different phases.
- Female odor may play an important role in the mating strategies of both sexes in Japanese macaques.

A. Urine sample from the fertile phase (female 3, sample ID: 63)






B. Vaginal sample from the fertile phase (female 3, sample ID: 94)










A**B****C**


Female 1
 Female 2
 Female 3
 Female 4
 Female 5







A**B****C**

Female 1
 Female 2
 Female 3









Retention time (mins)	Tentative compound ID	Mean peak area of compounds and number of samples (total or fertile phase) containing a compound, N=30 samples, N=65 samples											Literature review
		Mean peak area	total (fertile phase)	female 1 cycle 1 (N=7)	female 2 cycle 1 (N=8)	female 2 cycle 2 (N=8)	female 3 cycle 1 (N=8)	female 3 cycle 2 (N=8)	female 4 cycle 1 (N=8)	female 4 cycle 2 (N=5)	female 5 cycle 1 (N=6)	female 5 cycle 2 (N=7)	
1.329 - 1.340	Benzeneethanamine, 3-fluoro-beta, 5-dihydroxy-N-methyl-	2048534	3 (1)	-	-	-	-	1 (1)	2 (0)	-	-	-	
1.510 - 1.557	Acetone ¹	9121479	35 (8)	6 (2)	7 (1)	1 (0)	6 (2)	4 (1)	2 (1)	2 (1)	4 (0)	3 (0)	aye-eyes (del Barco et al., 2013)
1.615	Unknown 01	846614	1 (0)	-	-	-	-	-	1 (0)	-	-	-	
1.852 - 1.858	5,9-Dodecadien-2-one, 6,10-dimethyl-(E,E)-	2944550	2 (2)	1 (1)	-	-	-	1 (1)	-	-	-	-	
2.557 - 2.588	2-Pentanone	5598740	45 (12)	7 (3)	3 (1)	2 (0)	1 (1)	7 (1)	7 (2)	5 (1)	6 (1)	7 (2)	
3.245	2-Hexanone	850499	1 (1)	-	-	-	-	1 (1)	-	-	-	-	
3.322 - 3.338	Dimethyl disulfide	846330	14 (4)	1 (1)	2 (0)	1 (0)	2 (0)	4 (1)	1 (1)	1 (0)	1 (0)	1 (1)	olive baboons (Vaglio et al., 2021)
3.342 - 3.467	2-Pentanone, 3-methyl-	1371808	12 (2)	-	1 (0)	-	-	5 (2)	-	1 (0)	2 (0)	3 (0)	
3.703 - 3.72	Toluene ¹	1199682	31 (7)	5 (2)	4 (1)	3 (0)	1 (0)	5 (0)	8 (2)	3 (1)	-	2 (1)	olive baboons (Vaglio et al., 2021)
4.35	Hexanal	427641	1 (1)	1 (1)	-	-	-	-	-	-	-	-	aye-eyes (del Barco et al., 2013), mandrills (Vaglio et al., 2016), red ruffed lemrus (Janda et al., 2019), tamarins (Poirier, Waterhouse, Watsa, et al., 2021), olive baboons (Vaglio et al., 2021)
6.467 - 6.475	2-Heptanone	861478	3 (1)	1 (0)	-	-	-	1 (0)	1 (1)	-	-	-	
6.591	3-Cyclohexen-1-ol-methyl-	694526	1 (1)	-	-	-	-	-	1 (1)	-	-	-	

6.820 - 7.091	Oxime-, methoxy-phenyl- *	5918850	61 (17)	7 (7)	7 (2)	8 (2)	7 (2)	7 (2)	8 (2)	4 (1)	6 (1)	7 (2)	mandrills (Setchell et al., 2010), olive baboons (Vaglio et al., 2021)
8.252	Bicyclo[3.1.0]hex-2-ene, 4-methylene-1-(1-methylethyl)-	590611	1 (0)	-	-	-	-	-	-	-	-	1 (0)	
8.499 - 8.588	Benzaldehyde	5593836	19 (4)	6 (2)	-	-	3 (0)	4 (2)	-	1 (0)	4 (1)	1 (0)	marmosets (Smith et al., 2001; Kücklich et al., 2019), aye-ayes (del Barco et al., 2013), mandrills (Vaglio et al., 2016), red ruffed lemur (Janda et al., 2019), tamarins (Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021), olive baboons (Vaglio et al., 2021)
10.357 - 10.367	p-Cymene	30146684	1 (1)	-	-	-	-	-	1 (1)	-	-	-	owl monkeys (MacDonald et al., 2008)
10.504 - 10.525	D-Limonene	773712	5 (0)	2 (0)	1 (0)	-	-	-	-	1 (1)	-	1 (0)	mandrills (Setchell et al., 2010), marmosets (Kücklich et al., 2019), olive baboons (Vaglio et al., 2021)
10.518	1-Hexanol, -2-ethyl-	629876	2 (1)	-	-	-	2 (1)	-	-	-	-	-	mandrills (Setchell et al., 2010; Vaglio et al., 2016), red ruffed lemurs (Janda et al., 2019)
10.589	Eucalyptol	9438479	1 (0)	-	1 (0)	-	-	-	-	-	-	-	
12.337	Benzene, 1-methyl-4-(1-methylethenyl)-	9180801	1 (1)	-	-	-	-	-	1 (1)	-	-	-	
12.795 - 12.801	Nonanal	916644	16 (4)	3 (1)	-	-	1 (0)	1 (0)	8 (2)	3 (1)	-	-	mandrills (Setchell et al., 2010; Vaglio et al., 2016), aye-ayes (del Barco et al., 2013)
13.411 - 13.421	Benzyl methyl ketone	2843039	2 (1)	1 (0)	-	-	-	1 (0)	-	-	-	-	
14.004 - 14.008	cis-Verbenol	483251	2 (0)	-	-	-	-	-	-	-	1 (0)	1 (0)	red ruffed lemurs (Janda et al., 2019)
14.128	Bicyclo[3.1.0]hexane, 1,5-dimethyl-	8350650	1 (1)	-	-	-	-	-	1 (1)	-	-	-	
14.352	Bicyclo[3.1.0]hexan-2-one, 5-(1-methylethyl)-	440692	1 (1)	-	-	-	-	-	1 (1)	-	-	-	

14.566 - 14.676	Phenol, 4-ethyl-	12386393	21 (4)	3 (2)	-	2 (0)	2 (0)	6 (1)	4 (1)	2 (0)	-	2 (0)	red ruffed lemurs (Janda et al., 2019)
14.950	Cyclohexanol, 5-methyl-2-(1-methylethyl)-, (1.alpha.,2.beta.,5.alpha.)-(./-)-	266716	1 (0)	1 (0)	-	-	-	-	-	-	-	-	mandrills (Vaglio et al., 2016)
15.217	Phenol, 2,3,5,6-tetramethyl-	715590	1 (1)	-	-	-	-	-	1 (1)	-	-	-	
15.455 - 15.462	alpha-Terpineol	1399674	1 (1)	-	-	-	-	-	1 (1)	-	-	-	owl monkeys (MacDonald et al., 2008), red ruffed lemurs (Janda et al., 2019), olive baboons (Vaglio et al., 2021)
15.773 - 15.787	Decanal	701992	3 (1)	-	-	-	-	-	2 (1)	1 (0)	-	-	mandrills (Vaglio et al., 2016)
19.669 - 19.685	Pentanoic acid, 2,2,4-trimethyl-3-hydroxy-, isobutyl ester	877286	30 (9)	3 (1)	2 (1)	3 (1)	3 (1)	5 (1)	6 (2)	3 (1)	2 (0)	3 (1)	
20.311 - 20.328	Propanoic acid, 2-methyl-, 3-hydroxy-2,2,4-trimethylpentyl ester	1326768	31 (11)	3 (1)	2 (1)	4 (2)	4 (2)	5 (2)	6 (2)	3 (1)	2 (0)	2 (0)	mandrills (Vaglio et al., 2016)
25.667 - 25.682	2,2,4-Trimethyl-1,3-pentanediol diisobutyrate	950000	44 (14)	4 (2)	2 (1)	4 (2)	5 (2)	8 (2)	6 (1)	5 (1)	4 (1)	6 (2)	
29.884	2-Butenoic acid, 3-[[[(Ethylamino)methoxy phosphinothioyl]oxy]-, 1-methylethyl ester, (E)-	1858935	1 (0)	-	-	-	-	-	1 (0)	-	-	-	
31.087	Unknown 04	330128	1 (0)	1 (0)	-	-	-	-	-	-	-	-	
31.382	Ethane, 1-(2,3-xylyl)-1-(3,4-xylyl)-	356324	1 (0)	1 (0)	-	-	-	-	-	-	-	-	
Compounds unique to a female				4	1	2		10		2			

Retention time (mins)	Tentative compound ID	Mean peak area of compounds and number of samples (total or fertile phase) containing a compound, N=30 samples						Literature review
		Mean peak area	total (fertile phase)	female 1 cycle 1 (N=7)	female 2 cycle 1 (N=8)	female 2 cycle 2 (N=8)	female 3 cycle 1 (N=7)	
1.348 - 1.360	Unknown 01	116464	7 (3)	1 (0)	1 (1)	4 (2)	1 (0)	
1.397 - 1.418	Methanethiol	1792778	15 (3)	5 (2)	4 (1)	4 (0)	2 (0)	
1.420 - 1.445	Ethanol	1570304	13 (3)	1 (0)	5 (1)	4 (0)	3 (2)	marmosets (Kücklich et al., 2019)
1.592 - 1.613	Acetic acid, methyl ester	2148595	9 (3)	3 (1)	2 (1)	2 (1)	2 (0)	
1.680 - 1.707	Propanal, 2-methyl-	980390	8 (3)	-	3 (1)	4 (1)	1 (1)	
1.833 - 1.870	2-Butanone	1788558	14 (3)	1 (0)	7 (2)	2 (0)	4 (1)	aye-ayes (del Barco et al., 2013)
1.960 - 2.005	Ethyl acetate	799353	4 (0)	1 (0)	1 (0)	1 (0)	1 (0)	aye-ayes (del Barco et al., 2013)
2.287 - 2.345	Butanal, 3-methyl-	5712304	15 (6)	-	3 (2)	6 (2)	6 (2)	
2.368 - 2.374	Butanal, 2-methyl-	963903	15 (5)	-	4 (2)	6 (1)	5 (2)	
2.480 - 2.521	2-Propanol, 1-methoxy-	197917	2 (2)	1 (1)	1 (1)	-	-	
2.544 - 2.577	2-Pentanone	3615417	18 (4)	2 (1)	7 (2)	5 (1)	4 (0)	
2.857 - 2.870	Unknown 02	2958386	6 (2)	-	-	1 (0)	5 (2)	
2.924 - 3.013	Unknown 03	2118855	7 (3)	1 (0)	-	1 (1)	5 (2)	
3.147 - 3.189	Unknown 04	834708	6 (2)	-	-	5 (1)	1 (0)	
3.203 - 3.220	1-Butanol, 3-methyl-	2845956	13 (3)	-	4 (1)	3 (0)	6 (2)	
3.247 - 3.278	Methyl isobutyl ketone*	531969	6 (0)	-	3 (0)	3 (0)	-	olive baboons (Vaglio et al., 2021)
3.326 - 3.454	Dimethyl disulphide	12499870	13 (2)	1 (0)	7 (2)	2 (0)	3 (0)	
3.705 - 3.745	Toluene ¹	734202	14 (2)	6 (1)	2 (0)	6 (1)	-	olive baboons (Vaglio et al., 2021)
4.099 - 4.122	2-Hexanone	264372	4 (0)	-	2 (0)	1 (0)	1 (0)	
4.400	Unknown 05	773848	1 (1)	-	-	1 (1)	-	
5.399	Unknown 06	43547	1 (0)	-	-	1 (0)	-	
5.704	Unknown 07	17726	1 (1)	-	-	1 (1)	-	
5.955 - 5.967	1-Hexanol	987535	5 (0)	-	1 (0)	4 (0)	-	
6.314 - 6.324	Pyridine, 2,3-dimethyl-	944530	5 (0)	-	-	5 (0)	-	
6.444 - 6.465	2-Heptanone	2274192	16 (2)	1 (0)	6 (1)	6 (1)	3 (0)	

6.741 - 6.749	3-Methylcyclopentyl acetate	348834	12 (3)	1 (0)	1 (1)	5 (0)	5 (2)	marmosets (Kücklich et al., 2019)
6.848 - 6.971	Unknown 08*	8730763	29 (7)	7 (2)	8 (2)	7 (1)	7 (2)	
7.000 - 7.108	Pyrazine, 2,5-Dimethyl-	15406824	6 (2)	1 (0)	3 (38%)	2 (1)	-	marmosets (Smith et al., 2001; Kücklich et al., 2019)
7.662 - 7.663	Unknown 09	81985	2 (1)	-	2 (1)	-	-	
7.914 - 7.918	Unknown 10	1195235	4 (3)	-	2 (1)	-	2 (2)	
8.057 - 8.101	2(5H)-Furanone, 5,5-Dimethyl-	810258	8 (1)	-	1 (1)	5 (0)	2 (0)	
8.476 - 8.669	Unknown 11	16014736	11 (2)	-	5 (1)	3 (0)	3 (1)	
9.038 - 9.069	1-Octen-3-ol*	3016803	9 (0)	1 (0)	4 (0)	4 (0)	-	olive baboons (Vaglio et al., 2021)
9.320 - 9.345	Furan, 2-pentyl-*	2604348	5 (1)	-	-	5 (1)	-	
9.664	Pyrazine, trimethyl-	451271	1 (0)	-	-	1 (0)	-	marmosets (Smith et al., 2001)
10.002 - 10.023	Acetic acid, hexyl ester	645740	3 (1)	-	-	3 (1)	-	marmosets (Smith et al., 2001)
10.131	Butanal, 2-ethyl-	242333	1 (1)	-	-	-	1 (1)	
10.249 - 10.253	Unknown 12	686034	2 (0)	-	-	2 (0)	-	
10.585 - 10.593	Eucalyptol*	917434	3 (1)	1 (0)	2 (1)	-	-	
10.687 - 10.709	2(3H)-Furanone, 5-ethenyldihydro-5-methyl-	1597813	9 (0)	-	2 (0)	5 (0)	2 (0)	
10.830	Unknown 13	964162	1 (0)	-	-	1 (0)	-	
11.136 - 11.186	2(3H)-Furanone, 5-ethyldihydro-	2663385	8 (2)	-	-	8 (2)	-	
11.390 - 11.398	Unknown 14	332284	4 (0)	-	-	4 (0)	-	
11.568 - 11.599	Acetophenone	5883222	6 (0)	1 (0)	3 (0)	2 (0)	-	aye-ayes (del Barco et al., 2013), tamarins (Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021)
11.757	Ethyl 2-(5-methyl-5-vinyltetrahydrofuran-2-yl)propan-2-yl carbonate*	2561160	17 (4)	-	4 (1)	7 (2)	6 (1)	
11.902	Pyrazine, 3-ethyl-2,5-dimethyl-*	9055134	10 (4)	1 (1)	7 (2)	2 (1)	-	
12.175 - 12.194	Unknown 15	761371	12 (2)	-	2 (0)	6 (1)	4 (1)	
12.236 - 12.252	Unknown 16*	1709345	10 (1)	-	2 (0)	6 (1)	2 (0)	
12.366	Unknown 17	595762	1 (0)	-	1 (0)	-	-	

12.727 - 12.795	Unknown 18*	1080522	6 (1)	2 (1)	-	4 (0)	-	
13.266 - 13.303	Unknown 19	1229052	9 (2)	2 (0)	3 (1)	4 (1)	-	
13.444 - 13.453	Unknown 20*	878941	15 (3)	-	5 (1)	6 (1)	4 (1)	
14.016 - 14.029	Camphor	535695	3 (1)	-	1 (1)	1 (0)	1 (0)	
14.761 - 14.763	endo-Borneol	1581987	2 (1)	-	2 (1)	-	-	
15.321 - 15.335	Unknown 21	107094	2 (0)	-	-	2 (0)	-	
15.455	Unknown 22	228335	1 (0)	1 (0)	-	-	-	
16.685	Unknown 23	64839	1 (0)	-	-	-	1 (0)	
16.818	2-Propanol, 1-(2-butoxy-1-methylethoxy)-	938632	1 (0)	-	-	-	1 (0)	
16.874	Unknown 24	1362729	1 (0)	-	-	1 (0)	-	
17.027	Dodecane	999273	1 (0)	-	-	1 (0)	-	mandrills (Setchell et al., 2010; Vaglio et al., 2016)
17.369 - 17.371	Unknown 25	198053	2 (2)	-	1 (1)	1 (1)	-	
18.165 - 18.217	Unknown 26	1192500	2 (0)	-	2 (0)	-	-	
20.088 - 20.110	Unknown 27	1274496	18 (5)	6 (2)	3 (1)	6 (2)	3 (0)	
23.689 - 23.693	.alpha.-Farnesene	242211	2 (0)	-	-	2 (0)	-	ring-tailed lemurs (Scordato et al., 2013)
26.729 - 26.731	Unknown 28	496909	2 (1)	2 (1)	-	-	-	
28.056 - 28.068	Unknown 29	823238	7 (3)	3 (2)	3 (1)	1 (0)	-	
31.456 - 31.462	Unknown 30	394494	5 (3)	3 (2)	2 (1)	-	-	
34.489	Unknown 31	11432	1 (1)	1 (1)	-	-	-	
Compounds unique to a female				3		22	3	

Chemical cues of identity and reproductive status in Japanese macaques

Lucie Rigaiil, Stefano Vaglio, Joanna M. Setchell, Naoko Suda-Hashimoto, Takeshi Furuichi, Cécile Garcia

Table S1. Definitions of training levels for the five behaviors macaques were trained to perform during the study.

Name	Definition
Bridge	The female is conditioned and associates the sound clicker with a food reward.
Cooperation level 1	The female comes in front of the cage and takes the food reward in the food box, trainer stands within 1m distance from the cage.
Cooperation level 2	The female comes in front of the cage and takes the food reward directly from the trainer's hand.
Cooperation level 3	The female comes in front of the cage and takes the food reward directly from the trainer's hand. The female stays in front of the cage.
Hand training level 1	The female touches the trainer's hand that holds the food reward.
Hand training level 2	The female touches the trainer's hand in the absence of the reward.
Hand training level 3	The female puts her hand on the trainer's hand and hold still for 3 seconds.
Target training level 1	The female takes the food reward on the target stick presented before her.
Target training level 2	The female touches the target stick presented before her.
Target training level 3	The female moves toward the target stick and puts her hand on it.
Target training level 4	The female puts her hand on the target stick and hold still for 3 seconds.
Target training level 5	The female touches the hanging target stick with her hand, the target is hanging in the front of the cage.
Target training level 6	The female puts her hand on the hanging target stick and hold still for 3 seconds, the target is hanging in the front of the cage.
Target training level 7	The female puts her hand on the hanging target stick and hold still for 3 seconds, the target is hanging in the back of the cage.
Presentation training level 1	The female turns around, shows her hindquarters to the trainer, and touches the hanging target stick that hangs on the back of the cage.
Presentation training level 2	The female turns around, shows her hindquarters to the trainer, and touches the hanging target stick that hangs on the back of the cage. The trainer touches the female hindquarters with another target stick.
Presentation training level 3	The female presents her hindquarters to the trainer and hold still for 3 seconds while the trainer touches her hindquarters with another target stick.
Vaginal sampling training level 1	The female presents her hindquarters to the trainer, the trainer touches her hindquarters with a cotton swab.
Vaginal sampling training level 2	The female presents her hindquarters to the trainer and hold still for 3 seconds while the trainer touches her hindquarters with a cotton swab.
Vaginal sampling training level 3	The female presents her hindquarters to the trainer and accepts vaginal sampling with a cotton swab.